



Digital Receipt

This receipt acknowledges that Turnitin received your paper. Below you will find the receipt information regarding your submission.

The first page of your submissions is displayed below.

Submission author: Herry Suryawan
Assignment title: Quick Submit
Submission title: Turnitin AIP 2020
File name: AIP_Desember_2020.pdf
File size: 389.25K
Page count: 10
Word count: 4,395
Character count: 18,907
Submission date: 15-Apr-2023 10:18AM (UTC+0700)
Submission ID: 2065003659

AIP Conference Proceedings

A white noise approach to subfractional Brownian motion

Cite as: AIP Conference Proceedings **2286**, 020003 (2020); <https://doi.org/10.1063/5.0029753>
Published Online: 03 December 2020

Herry Pribawanto Suryawan

 

ARTICLES YOU MAY BE INTERESTED IN

Fractional Brownian motion - Some recent results and generalizations
AIP Conference Proceedings **2286**, 020001 (2020); <https://doi.org/10.1063/5.0029699>

From Markovian to non-Markovian stochastic processes: Diffusion coefficients with power law time dependence
AIP Conference Proceedings **2286**, 020005 (2020); <https://doi.org/10.1063/5.0029883>

Anomalous diffusion in the one-dimensional linear chain driven internally by one-parameter Mittag-Leffler noise
AIP Conference Proceedings **2286**, 020004 (2020); <https://doi.org/10.1063/5.0030081>

 

Your Qubits. Measured.
Meet the next generation of quantum analyzers

- Readout for up to 16 qubits
- Operation at up to 18.5 GHz, mixer-calibration free
- Signal optimization with minimal latency

[Find out more](#)

AIP Conference Proceedings **2286**, 020003 (2020); <https://doi.org/10.1063/5.0029753> **2286**, 020003
© 2020 Author(s).

Turnitin AIP 2020

by Suryawan Herry

Submission date: 15-Apr-2023 10:18AM (UTC+0700)

Submission ID: 2065003659

File name: AIP_Desember_2020.pdf (389.25K)

Word count: 4395

Character count: 18907

A white noise approach to subfractional Brownian motion

13

Cite as: AIP Conference Proceedings **2286**, 020003 (2020); <https://doi.org/10.1063/5.0029753>

Published Online: 03 December 2020

Herry Pribawanto Suryawan



View Online



Export Citation

ARTICLES YOU MAY BE INTERESTED IN

12

[Fractional Brownian motion - Some recent results and generalizations](#)
AIP Conference Proceedings **2286**, 020001 (2020); <https://doi.org/10.1063/5.0029699>

14

[From Markovian to non-Markovian stochastic processes: Diffusion coefficients with power law time dependence](#)

8

AIP Conference Proceedings **2286**, 020005 (2020); <https://doi.org/10.1063/5.0029883>

3

[Anomalous diffusion in the one-dimensional linear chain driven internally by one-parameter Mittag-Leffler noise](#)
AIP Conference Proceedings **2286**, 020004 (2020); <https://doi.org/10.1063/5.0030081>

Your Qubits. Measured.

Meet the next generation of quantum analyzers

- Readout for up to 64 qubits
- Operation at up to 8.5 GHz, mixer-calibration-free
- Signal optimization with minimal latency

[Find out more](#)


23

AIP Conference Proceedings **2286**, 020003 (2020); <https://doi.org/10.1063/5.0029753>**2286**, 020003

© 2020 Author(s).

A White Noise Approach to Subfractional Brownian Motion

Herry Pribawanto Suryawan^{a)}

Department of Mathematics, Sanata Dharma University, Yogyakarta, 55282, Indonesia.

^{a)}Corresponding author: herrypribs@usd.ac.id

Abstract. In this paper we study the subfractional Brownian motion by using white noise analysis. First we recall a representation of subfractional Brownian motion on the white noise space and prove the existence of the Donsker delta function of subfractional Brownian motion. We also solve the Langevin equation and a Wick-type linear stochastic differential equation driven by subfractional Brownian motion by using the S-transform method.

INTRODUCTION

The fractional Gaussian processes have been intensively studied due to the fact that the theory is interesting in itself and at the same time it has a broad range of applications. Fractional Brownian motion (fBm) seems to be the simplest fractional Gaussian process. It was introduced by Kolmogorov in [1] with the definition as follows. The fBm with Hurst parameter $H \in (0, 1)$ is a centered Gaussian process $B^H = (B^H(t))_{t \geq 0}$ with $B^H(0) = 0$ a.s. and covariance function

$$\text{Cov}(B^H(t), B^H(s)) = \frac{1}{2} (t^{2H} + s^{2H} - |t-s|^{2H}), \quad s, t \geq 0.$$

The case $H = \frac{1}{2}$ corresponds to the standard Brownian motion (Bm). Mandelbrot and Van Ness in [2] further studied this process and, in particular, gained a representation of fBm as a stochastic integral with respect to Bm. It is well-known that fBm satisfies self-similarity, long/short-range dependence, Hölder continuity of the sample paths, and stationarity of increments. Furthermore, fBm is the only continuous Gaussian process which is self-similar and has stationary increments. Due to these properties fBm has been used as an important tool for stochastic modeling in hydrology, telecommunication, turbulence, image processing and segmentation, medical image analysis, network traffic analysis, financial mathematics, etc. A comprehensive study of fBm can be found in [3, 4, 5, 6, 7] and references therein.

Another generalization of the Brownian motion is the so-called subfractional Brownian motion (sfBm). It was introduced by Bojdecki *et al* in [8] in connection with the study of occupation time fluctuations of branching particle systems with Poisson initial condition. The sfBm with parameter $H \in (0, 1)$ is a centered Gaussian process $S^H = (S^H(t))_{t \geq 0}$ with $S^H(0) = 0$ a.s. and covariance function

$$\text{Cov}(S^H(t), S^H(s)) = s^{2H} + t^{2H} - \frac{1}{2} ((s+t)^{2H} + |t-s|^{2H}), \quad s, t \geq 0. \quad (1)$$

When $H = \frac{1}{2}$ one recovers the standard Bm. One way to establish the existence of sfBm is by considering the process $\xi^H = (\xi^H(t))_{t \geq 0}$ defined by

$$\xi^H(t) := \frac{B^H(t) + B^H(-t)}{\sqrt{2}}, \quad t \geq 0,$$

where $(B^H(t))_{t \in \mathbb{R}}$ is a two-sided fBm. One can easily check that the covariance function of ξ^H coincides with (1). For details we refer to [6, 8].

The sfBm is intermediate between Bm and fBm in the sense that it has properties analogous to those of fBm, but the increments on non-overlapping intervals are more weakly correlated, and their covariance decays polynomially at a higher rate. In some applications, such as turbulence phenomena in hydromechanics, fBm is an adequate model for small increments, but it seems to be inadequate for large increments. For this reason, sfBm may be an alternative to fBm in some stochastic models.

The sfBm is neither a semimartingale nor a Markov process unless $H = \frac{1}{2}$. This implies that the powerful techniques from classical stochastic calculus are not available when dealing with S^H . sfBm shares many properties to those of fBm such as self-similarity and long-range dependence. On the contrary the increments of sfBm are not stationary.

It is known that sfBm is a quasi-helix in the sense of Kahane, that is it satisfies the inequalities, for any $s, t \geq 0$ with $t > s$,

$$((2 - 2^{2H-1}) \wedge 1) (t-s)^{2H} \leq \mathbb{E} (S^H(t) - S^H(s))^2 \leq ((2 - 2^{2H-1}) \vee 1) (t-s)^{2H}.$$

As a consequence, sample paths of sfBm is Hölder continuous of order γ for any $\gamma < H$. As a conclusion sfBm is suitable for modeling of random phenomenon which possesses self-similarity, long/short dependence, continuous sample paths but non-stationary increments. Some works on sfBm, including basic and advanced properties, stochastic calculus, local times, can be found in [9, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20], among others.

In this paper we realize the realization of sfBm on the white noise probability space and prove that the Donsker delta function of sfBm is a white noise distribution in the Hida sense. We also construct solutions to a subfractional version of the Ornstein-Uhlenbeck and a linear stochastic differential with Wick-multiplicative subfractional white noise in a suitable white noise distribution space.

BASICS ON WHITE NOISE ANALYSIS

In this section we give background on the white noise theory used throughout this paper. For a more comprehensive discussions including various applications of white noise theory we refer to [21, 22, 23] and references therein. We start with the Gelfand triple

$$\mathcal{S}_d(\mathbb{R}) \hookrightarrow L^2_d(\mathbb{R}) \hookrightarrow \mathcal{S}'_d(\mathbb{R}),$$

where $\mathcal{S}_d(\mathbb{R})$ is the space of \mathbb{R}^d -valued Schwartz test function, $\mathcal{S}'_d(\mathbb{R})$ is the space of \mathbb{R}^d -valued tempered distributions, and $L^2_d(\mathbb{R})$ is the real Hilbert space of \mathbb{R}^d -valued Lebesgue square-integrable functions. Next, we construct a probability space $(\mathcal{S}'_d(\mathbb{R}), \mathcal{C}, \mu)$ where \mathcal{C} is the Borel σ -algebra generated by weak topology on $\mathcal{S}'_d(\mathbb{R})$ and the probability measure μ is uniquely determined through the Bochner-Minlos theorem by fixing the characteristic function

$$C(\vec{f}) := \int_{\mathcal{S}'_d(\mathbb{R})} \exp(i\langle \vec{\omega}, \vec{f} \rangle) d\mu(\vec{\omega}) = \exp\left(-\frac{1}{2} |\vec{f}|_0^2\right)$$

for all $\vec{f} \in \mathcal{S}_d(\mathbb{R})$. Here $|\cdot|_0$ denotes the usual norm in the $L^2_d(\mathbb{R})$, and $\langle \cdot, \cdot \rangle$ denotes the dual pairing between $\mathcal{S}'_d(\mathbb{R})$ and $\mathcal{S}_d(\mathbb{R})$. The dual pairing is considered as the bilinear extension of the inner product on $L^2_d(\mathbb{R})$, i.e.

$$\langle \vec{g}, \vec{f} \rangle = \sum_{j=1}^d \int_{\mathbb{R}} g_j(x) f_j(x) dx,$$

for all $\vec{g} = (g_1, \dots, g_d) \in L^2_d(\mathbb{R})$ and $\vec{f} = (f_1, \dots, f_d) \in \mathcal{S}_d(\mathbb{R})$. This probability space is known as the \mathbb{R}^d -valued white noise space since it contains the sample paths of the d -dimensional Gaussian white noise. In this setting a d -dimensional Brownian motion can be represented by a continuous modification of the stochastic process $B = (B_t)_{t \geq 0}$ with

$$B(t) := (\langle \cdot, \mathbf{1}_{[0,t]} \rangle, \dots, \langle \cdot, \mathbf{1}_{[0,t]} \rangle),$$

such that independent d -tuples of Gaussian white noise $\vec{\omega} = (\omega_1, \dots, \omega_d) \in \mathcal{S}'_d(\mathbb{R})$

$$B(t, \vec{\omega}) = (\langle \omega_1, \mathbf{1}_{[0,t]} \rangle, \dots, \langle \omega_d, \mathbf{1}_{[0,t]} \rangle),$$

where $\mathbf{1}_A$ denotes the indicator function of a set $A \subset \mathbb{R}$.

In the sequel we will use the Gelfand triple

$$(\mathcal{S}) \hookrightarrow L^2(\mu) := L^2(\mathcal{S}'_d(\mathbb{R}), \mathcal{C}, \mu) \hookrightarrow (\mathcal{S})^*$$

where (\mathcal{S}) is the space of white noise test functions obtained by taking the intersection of a family of Hilbert subspaces of $L^2(\mu)$. The space of white noise distributions $(\mathcal{S})^*$ is defined as the topological dual space of (\mathcal{S}) .

Elements of (\mathcal{S}) and $(\mathcal{S})^*$ are known as *Hida test functions* and *Hida distributions*, respectively. An important tool in white noise analysis is the S-transform which can be considered as the Laplace transform with respect to the Gaussian measure. The S-transform of $\Phi \in (\mathcal{S})^*$ is defined as

$$(S\Phi)(\vec{f}) := \left\langle \left\langle \Phi, : \exp \left(\left\langle \cdot, \vec{f} \right\rangle \right) : \right\rangle \right\rangle, \quad \vec{f} \in \mathcal{S}_d(\mathbb{R}),$$

where

$$: \exp \left(\left\langle \cdot, \vec{f} \right\rangle \right) ::= C(\vec{f}) \exp \left(\left\langle \cdot, \vec{f} \right\rangle \right),$$

is the so-called Wick exponential and $\langle \cdot, \cdot \rangle$ denotes the dual pairing between $(\mathcal{S})^*$ and (\mathcal{S}) . We define this dual pairing as the bilinear extension of the sesquilinear inner product on $L^2(\mu)$. The S-transform provides a convenient way to identify a Hida distribution $\Phi \in (\mathcal{S})^*$, in particular, when it is hard to find the explicit form for the Wiener-Itô chaos decomposition of Φ . We now state some properties of the S-transform.

Theorem 1 [22]

1. The S-transform is injective, i.e. if $S\Phi(\varphi) = S\Psi(\varphi)$ for all $\varphi \in \mathcal{S}_1(\mathbb{R})$, then $\Phi = \Psi$.
2. Let a stochastic distribution process $X : I \rightarrow (\mathcal{S})^*$ is differentiable. Then, $S \frac{d}{dt} X(t)(\varphi) = \frac{d}{dt} SX(t)(\varphi)$ for all $\varphi \in \mathcal{S}_1(\mathbb{R})$.

In white noise analysis the Wick product of two distributions using the S-transform. Let $\Phi, \Psi \in (\mathcal{S})^*$. The Wick product of Φ and Ψ is defined as

$$\Phi \diamond \Psi := S^{-1}(S\Phi \cdot S\Psi).$$

We will need also the Wick exponential function \exp^\diamond which is defined for $\Phi \in (\mathcal{S})^*$ by

$$S \exp^\diamond(\Phi)(\varphi) = \exp^\diamond(S\Phi(\varphi)), \quad \varphi \in \mathcal{S}_1(\mathbb{R}).$$

See [24] for details on Wick product and calculus based on it.

Finally in this section we state a sufficient condition on the Bochner integrability of a family of Hida distributions which depend on an additional parameter.

Theorem 2 [25] Let $(\Omega, \mathcal{A}, \nu)$ be a measure space and $\lambda \mapsto \Phi_\lambda$ be a mapping from Ω to $(\mathcal{S})^*$. If

- (1) the mapping $\lambda \mapsto S(\Phi_\lambda)(\vec{f})$ is measurable for all $\vec{f} \in \mathcal{S}_d(\mathbb{R})$, and
- (2) there exist $C_1(\lambda) \in L^1(\Omega, \mathcal{A}, \nu)$, $C_2(\lambda) \in L^\infty(\Omega, \mathcal{A}, \nu)$ and a continuous seminorm $\|\cdot\|$ on $\mathcal{S}_d(\mathbb{R})$ such that for all $z \in \mathbb{C}$, $\vec{f} \in \mathcal{S}_d(\mathbb{R})$

$$\left| S(\Phi_\lambda)(z\vec{f}) \right| \leq C_1(\lambda) \exp \left(C_2(\lambda) |z|^2 \|\vec{f}\|^2 \right),$$

then Φ_λ is Bochner integrable with respect to some Hilbertian norm which topologizing $(\mathcal{S})^*$. Hence $\int_\Omega \Phi_\lambda d\nu(\lambda) \in (\mathcal{S})^*$, and furthermore

$$S \left(\int_\Omega \Phi_\lambda d\nu(\lambda) \right) = \int_\Omega S(\Phi_\lambda) d\nu(\lambda).$$

WHITE NOISE ANALYSIS OF SFBM

In this section we recall a representation of sfBm on the white noise space and show the existence of the Donsker delta function of sfBm. In order to represent sub-fBm on the white noise space, we use the following operator

$$M_-^H f := \begin{cases} C_H D_-^{-(H-\frac{1}{2})} f & \text{if } 0 < H < \frac{1}{2} \\ f & \text{if } H = \frac{1}{2} \\ C_H I_-^{H-\frac{1}{2}} f & \text{if } \frac{1}{2} < H < 1, \end{cases}$$

where $C_H = \sqrt{2H \sin(\pi H) \Gamma(2H)}$ dan Γ denotes the gamma function. Here $I_-^\beta f$, $0 < \beta < 1$, is the Weyl type fractional integral operator defined by

$$(I_-^\beta f)(x) := \frac{1}{\Gamma(\beta)} \int_x^\infty f(t) (t-x)^{\beta-1} dt \quad 28$$

if the integrals exist for almost all $x \in \mathbb{R}$ and $D_-^\beta f$, $0 < \beta < 1$ is the Marchaud type fractional derivative operator defined by

$$(D_-^\beta f)(x) := \lim_{h \rightarrow 0^+} \frac{\beta}{\Gamma(1-\beta)} \int_h^\infty \frac{f(x) - f(x+t)}{t^{1+\beta}} dt, \quad 17$$

if the limit exists in $L^p(\mathbb{R})$ for some $p > 1$. For any Borel function f on $[0, \infty)$ we define its odd extension f° by

$$f^\circ(x) = \begin{cases} f(x) & \text{if } x \geq 0 \\ -f(-x) & \text{if } x < 0. \end{cases} \quad 36$$

Note that sfBm S^H can be written as a Volterra process with the following moving average representation (see e.g. [8]):

$$S^H(t) = K_H \int_{\mathbb{R}} \left((t-s)_+^{H-\frac{1}{2}} + (t+s)_-^{H-\frac{1}{2}} - 2(-s)_+^{H-\frac{1}{2}} \right) dB(s), \quad 41 \quad (2)$$

where

$$K_H = \frac{1}{\sqrt{2}} \left(\int_0^\infty \left((1+x)^{H-\frac{1}{2}} - x^{H-\frac{1}{2}} \right)^2 dx + \frac{1}{2H} \right)^{-1/2} = \frac{\sqrt{H \sin(\pi H) \Gamma(2H)}}{\Gamma(H + \frac{1}{2})}, \quad 33$$

$x_+ = \max\{x, 0\}$, $x_- = \max\{-x, 0\}$, and $(B(t))_{t \in \mathbb{R}}$ is a two-sided Brownian motion. For more information on a white noise approach to Volterra process see [26]. Based on the moving average representation (2) one can show the following.

Proposition 1 [17] It holds that $M_-^H \mathbf{1}_{[0,t]}^\circ \in L^2(\mathbb{R})$ and

$$S^H(t) = \frac{1}{\sqrt{2}} \int_{\mathbb{R}} (M_-^H \mathbf{1}_{[0,t]}^\circ)(s) dB_s = \left\langle \cdot, \frac{1}{\sqrt{2}} M_-^H \mathbf{1}_{[0,t]}^\circ \right\rangle, \quad 39$$

where $B = (B(t))_{t \in \mathbb{R}}$ is a one-dimensional Brownian motion on the real line and $\mathbf{1}_A^\circ$ denotes the odd extension of the indicator function of a set $A \subset \mathbb{R}$. 68

The generalized stochastic process $W^H = (W(t)^H)_{t \geq 0}$ defined by

$$W^H(t) := \frac{d}{dt} (S^H(t)) = \lim_{h \rightarrow 0} \frac{S^H(t+h) - S^H(t)}{h}, \quad 63$$

where the convergence takes place in the Hida distribution space $(\mathcal{S})^*$, is called the one-dimensional subfractional white noise. Note that for any $t \geq 0$ $S^H(t) \in L^2(\mu)$ and $W^H(t) \in (\mathcal{S})^*$. The d -dimensional sfBm can be represented on the white noise space by a continuous modification of the stochastic process $S^H = (S^H(t))_{t \geq 0}$ with

$$S^H(t) := \left(\left\langle \cdot, \frac{1}{\sqrt{2}} M_-^H \mathbf{1}_{[0,t]}^\circ \right\rangle, \dots, \left\langle \cdot, \frac{1}{\sqrt{2}} M_-^H \mathbf{1}_{[0,t]}^\circ \right\rangle \right), \quad 37$$

such that for independent d -tuples of Gaussian white noise $\vec{\omega} = (\omega_1, \dots, \omega_d) \in \mathcal{S}'_d(\mathbb{R})$

$$S^H(t, \vec{\omega}) = \left(\left\langle \omega_1, \frac{1}{\sqrt{2}} M_-^H \mathbf{1}_{[0,t]}^\circ \right\rangle, \dots, \left\langle \omega_d, \frac{1}{\sqrt{2}} M_-^H \mathbf{1}_{[0,t]}^\circ \right\rangle \right). \quad 15$$

Apart from the operator M_-^H above we will need the following operator:

$$M_+^H f := \begin{cases} C_H D_+^{-(H-\frac{1}{2})} f & \text{if } 0 < H < \frac{1}{2} \\ f & \text{if } H = \frac{1}{2} \\ C_H I_+^{H-\frac{1}{2}} f & \text{if } \frac{1}{2} < H < 1, \end{cases}$$

where, for $0 < \beta < 1$, $I_+^\beta f$ is the Weyl fractional integral operator defined by

$$(I_+^\beta f)(x) := \frac{1}{\Gamma(\beta)} \int_{-\infty}^x f(t) (x-t)^{\beta-1} dt$$

if the integrals exist for almost all $x \in \mathbb{R}$ and $D_+^\beta f$ is the Marchaud fractional derivative operator defined by

$$(D_+^\beta f)(x) := \lim_{h \rightarrow 0^+} \frac{\beta}{\Gamma(1-\beta)} \int_h^\infty \frac{f(x) - f(x-t)}{t^{1+\beta}} dt,$$

if the limit exists in $L^p(\mathbb{R})$ for some $p > 1$. One can show that M_-^H and M_+^H are dual operators in the sense that

$$(f, M_-^H g)_0 = (M_+^H f, g)_0$$

for any function f and g satisfying some regularity condition, for example indicator function or Schwartz test function. For details we refer to [27]. The following result shows that the operator M_\pm^H interchanges with the S-transform.

Lemma 1 [28] Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space. If $M_\pm^H X$ exist for some $X \rightarrow L^2(\Omega, \mathcal{F}, \mathbb{P})$, then

$$\mathbb{E}((M_\pm^H X)_t, \Phi) = M_\pm^H(\mathbb{E}(X_t, \Phi))$$

for all $\Phi \in L^2(\Omega, \mathcal{F}, \mathbb{P})$. For $H < \frac{1}{2}$ the convergence of the fractional derivative on the right-hand side is in the $L^p(\mathbb{R})$ sense, if $M_\pm^{-(H-\frac{1}{2})} X \in L^p(\mathbb{R}; \Omega, \mathcal{F}, \mathbb{P})$. In particular, the operator M_\pm^H interchange with the S-transform.

The S-transform of sfBm and subfractional white noise can be computed explicitly, see [17, 28] for proofs and details.

Lemma 2 Let $(S^H(t))_{t \geq 0}$ dan $(W^H(t))_{t \geq 0}$ be one-dimensional sfBm and subfractional white noise, respectively. Then, for any $\varphi \in \mathcal{S}_1(\mathbb{R})$ it holds

1.

$$SS^H(t)(\varphi) = \left\langle \varphi, \frac{1}{\sqrt{2}} M_-^H \mathbf{1}_{[0,t]}^\circ \right\rangle = \frac{1}{\sqrt{2}} \left(\int_0^t M_+^H \varphi(s) ds - \int_{-t}^0 M_+^H \varphi(s) ds \right).$$

2.

$$SW^H(t)(\varphi) = \frac{1}{\sqrt{2}} ((M_+^H \varphi)(t) - (M_+^H \varphi)(-t)).$$

Now we show that the Donsker delta function of a sfBm is a well-defined element in the space of Hida distributions.

Theorem 3 Let $S^H = (S_1^H, \dots, S_d^H)$ be a d -dimensional sfBm and $\vec{x} \in \mathbb{R}^d$. The Bochner integral

$$\delta(S^H(t) - \vec{x}) := \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} \exp(i\langle \vec{\lambda}, S^H(t) - \vec{x} \rangle) d\vec{\lambda}$$

is a Hida distribution with the S-transform given by

$$S\delta(S^H(t) - \vec{x})(\vec{\varphi}) = \left(\frac{1}{2\pi(2-2^{2H-1})t^{2H}} \right)^{\frac{d}{2}} \exp\left(-\frac{1}{2(2-2^{2H-1})t^{2H}} \sum_{j=1}^d \left(x_j - \left\langle \varphi_j, \frac{1}{\sqrt{2}} M_-^H \mathbf{1}_{[0,t]}^\circ \right\rangle\right)^2\right),$$

for any $\vec{\varphi} \in \mathcal{S}_d(\mathbb{R})$.

Proof. For any $\vec{\varphi} \in \mathcal{S}_d(\mathbb{R})$ we have

$$\begin{aligned}
\text{Sexp}\left(i\langle \vec{\lambda}, S^H(t) - \vec{x} \rangle\right)(\vec{\varphi}) &= \left\langle \left\langle \exp\left(i\langle \vec{\lambda}, S^H(t) - \vec{x} \rangle\right), \mathbb{C}(\vec{\varphi}) \exp(\langle \cdot, \vec{\varphi} \rangle)\right\rangle \right\rangle \\
&= \exp\left(i\langle \vec{\lambda}, \vec{x} \rangle - \frac{1}{2}|\vec{\varphi}|_0^2\right) \int_{\mathcal{S}'_d(\mathbb{R})} \exp\left(\left\langle \vec{\omega}, \vec{\varphi} - \frac{i\vec{\lambda}}{\sqrt{2}} M^H \mathbf{1}_{[0,t]}^{\circ} \right\rangle\right) d\mu(\vec{\omega}) \\
&= \exp\left(i\langle \vec{\lambda}, \vec{x} \rangle - \frac{1}{2}|\vec{\varphi}|_0^2\right) \exp\left(\frac{1}{2}\left|\vec{\varphi} - \frac{i\vec{\lambda}}{\sqrt{2}} M^H \mathbf{1}_{[0,t]}^{\circ}\right|_0^2\right) \\
&= \exp\left(i\langle \vec{\lambda}, \vec{x} - \langle \vec{\varphi}, \frac{1}{\sqrt{2}} M^H \mathbf{1}_{[0,t]}^{\circ} \rangle\right) - \frac{1}{2}|\vec{\lambda}|^2 \left|\frac{1}{\sqrt{2}} M^H \mathbf{1}_{[0,t]}^{\circ}\right|_0^2,
\end{aligned}$$

a measurable function of $\vec{\lambda}$. Moreover, for $\vec{\varphi} \in \mathcal{S}_d(\mathbb{R})$ and $z \in \mathbb{C}$ we obtain

$$\begin{aligned}
\left|\text{Sexp}\left(i\langle \vec{\lambda}, S^H(t) - \vec{x} \rangle\right)(z\vec{\varphi})\right| &\leq \prod_{j=1}^d \exp\left(-\frac{1}{2}\lambda_j^2 \text{Var}(S^H(t))\right) \exp\left(|z|\lambda_j \left|\left\langle \varphi_j, \frac{1}{\sqrt{2}} M^H \mathbf{1}_{[0,t]}^{\circ} \right\rangle\right|\right) \\
&\leq \prod_{j=1}^d \exp\left(-\frac{1}{2}\lambda_j^2 \text{Var}(S^H(t))\right) \exp\left(\frac{1}{4}\lambda_j^2 \text{Var}(S^H(t))\right) \exp(|z|^2|\varphi_j|^2) \\
&= \exp\left(-\frac{2-2^{2H-1}}{4}t^{2H}|\vec{\lambda}|^2\right) \exp(|z|^2|\vec{\varphi}|_0^2).
\end{aligned}$$

Note that the first factor is an integrable function of $\vec{\lambda}$ and the second factor is independent of $\vec{\lambda}$. Thus, Theorem 2 implies that $\delta(S^H(t) - \vec{x}) \in (\mathcal{S})^*$. Finally, by calculating the Gaussian integral we get

$$\begin{aligned}
\text{S}\delta(S^H(t) - \vec{x})(\vec{\varphi}) &= \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} \text{Sexp}\left(i\langle \vec{\lambda}, S^H(t) - \vec{x} \rangle\right)(\vec{\varphi}) d\vec{\lambda} \\
&= \prod_{j=1}^d \frac{1}{2\pi} \int_{\mathbb{R}} \exp\left(i\lambda_j \left(x_j - \left\langle \varphi_j, \frac{1}{\sqrt{2}} M^H \mathbf{1}_{[0,t]}^{\circ} \right\rangle\right) - \frac{1}{2}\lambda_j^2 \text{Var}(S^H(t))\right) d\lambda_j \\
&= \prod_{j=1}^d \frac{1}{\sqrt{2\pi \text{Var}(S^H(t))}} \exp\left(-\frac{1}{2(2-2^{2H-1})t^{2H}} \left(x_j - \left\langle \varphi_j, \frac{1}{\sqrt{2}} M^H \mathbf{1}_{[0,t]}^{\circ} \right\rangle\right)^2\right) \\
&= \left(\frac{1}{2\pi(2-2^{2H-1})t^{2H}}\right)^{\frac{d}{2}} \exp\left(-\frac{1}{2(2-2^{2H-1})t^{2H}} \sum_{j=1}^d \left(x_j - \left\langle \varphi_j, \frac{1}{\sqrt{2}} M^H \mathbf{1}_{[0,t]}^{\circ} \right\rangle\right)^2\right).
\end{aligned}$$

Corollary 1 The transition probability function of a particle in a system described by a d -dimensional sfBm $S^H = (S^H(t))_{t \geq 0}$ to move from $\vec{x}_0 \in \mathbb{R}^d$ to an endpoint $\vec{x}_T \in \mathbb{R}^d$ at a later time $t = T$ is given by

$$p(\vec{x}_0, \mathbf{0}; \vec{x}_T, T) = \left(\frac{1}{2\pi(2-2^{2H-1})t^{2H}}\right)^{\frac{d}{2}} \exp\left(-\frac{1}{2(2-2^{2H-1})t^{2H}} |\vec{x}_0 - \vec{x}_T|_{\mathbb{R}^d}^2\right).$$

Proof. The generalized expectation of the Donsker delta function can be obtained from Theorem 3 by evaluating the value of the S-transform at $\vec{\varphi} = \vec{0}$:

$$\mathbb{E}_{\mu}(\delta(S^H(t) - \vec{c})) = \text{S}\delta(S^H(t) - \vec{c})(\vec{0}) = \left(\frac{1}{2\pi(2-2^{2H-1})t^{2H}}\right)^{\frac{d}{2}} \exp\left(-\frac{1}{2(2-2^{2H-1})t^{2H}} \sum_{j=1}^d (x_0^j - c^j)^2\right).$$

The transition probability function $p(\vec{x}_0, \mathbf{0}; \vec{x}_T, T)$ follows immediately from the last expression by fixing the endtime $t = T$ with endpoint $\vec{c} = \vec{x}_T$.

For a discussion on the Donsker delta function of general class of stochastic processes with memory see [29].

26
WICK-TYPE LINEAR SDES DRIVEN BY SFBM

In this section we will solve the Langevin equation driven by sfBm and a linear stochastic differential equation (SDE) of Wick-type driven by sfBm by using the S-transform method. First, let us consider the Langevin equation driven by sfBm

$$dX^H(t) = -\lambda X^H(t) dt + \kappa dS^H(t), \quad X^H(0) = x_0, \quad (3)$$

where $\kappa \in \mathbb{R}$ dan $\lambda > 0$. The solution $X^H(t)$ of equation (3) represents the velocity at time t of a free particle which performs a subfractional Brownian-type motion but different from $S^H(t)$. The SDE (3) is interpreted in the integral form by

$$X^H(t) = x_0 - \lambda \int_0^t X^H(s) ds + \kappa S^H(t).$$

Applying the S-transform to $X^H(t)$ we obtain

$$SX^H(t)(\varphi) = x_0 - \lambda \int_0^t SX^H(s)(\varphi) ds + \kappa \left\langle \varphi, \frac{1}{\sqrt{2}} M_-^H \mathbf{1}_{[0,t]}^\circ \right\rangle, \quad (4)$$

for all $\varphi \in \mathcal{S}_1(\mathbb{R})$. Let us denote $z(t) := SX^H(t)(\varphi)$, then the equation (4) is equivalent to

$$z'(t) = -\lambda z(t) + \kappa \frac{d}{dt} \left\langle \varphi, \frac{1}{\sqrt{2}} M_-^H \mathbf{1}_{[0,t]}^\circ \right\rangle. \quad (5)$$

The solution of ODE (5) is given by

$$z(t) = x_0 \exp(-\lambda t) + \frac{\kappa}{\sqrt{2}} \left(\left\langle \varphi, M_-^H \mathbf{1}_{[0,t]}^\circ \right\rangle - \lambda \int_0^t \exp(\lambda(s-t)) \left\langle \varphi, M_-^H \mathbf{1}_{[0,s]}^\circ \right\rangle ds \right).$$

Hence, we have

$$SX^H(t)(\varphi) = x_0 \exp(-\lambda t) + \frac{\kappa}{\sqrt{2}} \left\langle \varphi, M_-^H \mathbf{1}_{[0,t]}^\circ - \lambda \int_0^t \exp(\lambda(s-t)) M_-^H \mathbf{1}_{[0,s]}^\circ ds \right\rangle.$$

Denoting

$$f_H(t) := \frac{1}{\sqrt{2}} \left(M_-^H \mathbf{1}_{[0,t]}^\circ - \lambda \int_0^t \exp(\lambda(s-t)) M_-^H \mathbf{1}_{[0,s]}^\circ ds \right)$$

the S-transform of $X^H(t)$ can be rewritten as

$$SX^H(t)(\varphi) = x_0 \exp(-\lambda t) + \kappa \langle \varphi, f_H(t) \rangle.$$

Taking the inverse of the S-transform yields

$$X^H(t) = x_0 \exp(-\lambda t) + \kappa \langle \cdot, f_H(t) \rangle. \quad (6)$$

Definition 1 The solution $X^H(t)$, $t \geq 0$, given by equation (6) of the Langevin equation (3) is called a subfractional Ornstein-Uhlenbeck process. Its expectation and characteristic function are given by

$$\begin{aligned} \mathbb{E}(X^H(t)) &= x_0 \exp(-\lambda t), \\ \mathbb{E}(\exp(ikX^H(t))) &= \exp\left(ikx_0 \exp(-\lambda t) - \frac{1}{2}k^2 |f_H(t)|_0^2\right). \end{aligned}$$

Next, we will consider linear SDEs of Wick-type of the form

$$dY^H(t) = -\lambda Y^H(t) dt + Y^H(t) \diamond dS^H(t), \quad Y^H(0) = \Phi_0 \in (\mathcal{S})^*. \quad (7)$$

In white noise analysis the connection between Wick product and Skorokhod integration can be expressed as

$$\int_{\mathbb{R}} Y(t) \delta B(t) = \int_{\mathbb{R}} Y(t) \diamond W(t) dt,$$

where the left-hand side denotes the Skorokhod integral of the stochastic process $(Y(t))_{t \geq 0}$ while the right-hand side corresponds to a $(\mathcal{S})^*$ -valued Pettis integral. Here, $(B(t))_{t \geq 0}$ is a standard Brownian motion and $(W(t))_{t \geq 0}$ is the corresponding white noise.

In order to give sense and solve equation (7) we recall the notion of integration of a stochastic distribution process in terms of the S-transform.

Theorem 4 [30] Let (T, \mathcal{F}, m) be a measure space and $\Phi(t) \in (\mathcal{S})^*$ for all $t \in T$. If

1. $S\Phi(\cdot)(\varphi)$ is measurable for any $\varphi \in \mathcal{S}_1(\mathbb{R})$,
2. $S\Phi(\cdot)(\varphi) \in L^1(T)$ for all $\varphi \in \mathcal{S}_1(\mathbb{R})$,
3. for any measurable set $C \subseteq T$ the function $\int_C S\Phi(u)(\cdot) dm(u)$ is the S-transform of some distribution in $(\mathcal{S})^*$,

then there exists a unique distribution $\Psi \in (\mathcal{S})^*$ such that

$$S\Psi(\varphi) = \int_T S\Phi(u)(\varphi) dm(u), \quad \varphi \in \mathcal{S}_1(\mathbb{R}).$$

The distribution Ψ is denoted by $\int_T \Phi(t) dm(t)$ and it is called the Pettis integral of Φ .

Now, equation (7) is interpreted in the integral form by

$$Y^H(t) = \Phi_0 - \lambda \int_0^t Y^H(s) ds + \int_0^t Y^H(s) \diamond W^H(t) dt, \quad (8)$$

where the integrals are to be understood in the Pettis sense as in Theorem 4. Next, we apply the S-transform to equation (8) and use Lemma 2 to obtain

$$SY^H(t)(\varphi) = S\Phi_0(\varphi) - \lambda \int_0^t SY^H(s)(\varphi) ds + \frac{1}{\sqrt{2}} \int_0^t SY^H(s) ((M_+^H \varphi)(s) - (M_+^H \varphi)(-s)) ds.$$

Denote $y(t) := SY^H(t)(\varphi)$, then

$$y'(t) = -\lambda y(t) + \frac{1}{\sqrt{2}} y(t) ((M_+^H \varphi)(t) - (M_+^H \varphi)(-t))$$

and the solution of this ODE is given by

$$y(t) = y(0) \exp(-\lambda t) \exp\left(\frac{1}{\sqrt{2}} ((M_+^H \varphi)(t) - (M_+^H \varphi)(-t))\right).$$

Rewriting the last expression using S-transform we get

$$SY^H(t)(\varphi) = \exp(-\lambda t) S\Phi_0(\varphi) \exp(SW^H(t)(\varphi)).$$

Finally, inverting the S-transform gives

$$Y^H(t) = \exp(-\lambda t) \Phi_0 \diamond \exp^\circ(W^H(t)), \quad t \geq 0. \quad (9)$$

From the previous considerations we obtain the following theorem.

Theorem 5 The solution $Y^H(t)$, $t \geq 0$, given by equation (9) of the linear SDE with Wick-multiplicative subfractional white noise is an element of $(\mathcal{S})^*$. Its generalized expectation is given by

$$\mathbb{E}_\mu(Y^H(t)) = \mathbb{E}_\mu(\Phi_0) \exp(-\lambda t).$$

CONCLUSION

We have discussed sfBm in the framework of white noise theory. The sfBm can be represented as functional of white noise by using some fractional integral/differential operator. We proved that Donsker's delta function of sfBm is a Hida distribution. We constructed the subfractional Ornstein-Uhlenbeck process as the solution of the Langevin equation. Finally, by using S-transform method, we also solved a linear SDE with Wick-multiplicative subfractional white noise.

ACKNOWLEDGMENTS

47

The author would like to thank Prof. Christopher C. Bernido and Prof. M. Victoria Carpio-Bernido for the invitation to the 9th Jagna International Workshop: Stochastic Analysis–Mathematical Methods and Real-World Models and for their hospitality.

REFERENCES

1. A. N. Kolmogorov, *Doklady Acad. Sci. URSS* **26**, 115–118 (1940).
2. B. B. Mandelbrot and J. W. V. Ness, *SIAM Review* **10**, 422–437 (1968).
3. F. Biagini, Y. Hu, B. Oksendal, and T. Zhang, *Stochastic Calculus for Fractional Brownian Motion and Applications* (Springer-Verlag, 2008).
4. Y. S. Mishura, *Stochastic Calculus for Fractional Brownian Motion and Related Processes* (Springer-Verlag, 2008).
5. I. Nourdin, *Selected Aspects of Fractional Brownian Motion* (Springer, 2012).
6. Y. S. Mishura and M. Zili, *Stochastic Analysis of Mixed Fractional Gaussian Processes* (ISTE Press, 2018).
7. G. Samorodnitsky and M. Taqqu, *Stable Non-Gaussian Random Processes* (CRC Press, 1994).
8. T. Bojdecki, L. G. Gorostiza, and A. Talarczyk, *Statist. Probab. Lett.* **69**, 717–739 (2004).
9. T. Bojdecki, L. G. Gorostiza, and A. Talarczyk, *J. Theoret. Probab.* **69**, 405–419 (2004).
10. T. Bojdecki, L. G. Gorostiza, and A. Talarczyk, *Elect. Comm. in Probab.* **12**, 161–172 (2007).
11. C. El-Nouty, *Stochastic Differential Equations and Processes* **7**, 179–196 (2012).
12. I. Mendy, *Annales Mathematiques Blaise Pascal* **17**, 357–374 (2010).
13. C. Tudor, *Stochastics* **79**, 431–448 (2007).
14. C. Tudor, *Statist. Probab. Lett.* **78**, 2201–2209 (2008).
15. C. Tudor, *Ann. Univ. Bucuresti Mathematica* **57**, 199–230 (2008).
16. C. Tudor, *J. Math. Anal. Appl.* **351**, 456–468 (2009).
17. Z. Wang and L. Yan, *Adv. Math. Phys.* **2013**, 827192:1–11 (2013).
18. L. Yan and G. Shen, *Statist. Probab. Lett.* **80**, 296–308 (2010).
19. L. Yan, G. Shen, and K. He, *Comm. Stoch. Anal.* **5**, 135–159 (2011).
20. H. P. Suryawan, *Jurnal Matematika Integratif* **15**, 81–87 (2019).
21. T. Hida, H.-H. Kuo, J. Potthoff, and L. Streit, *White Noise. An Infinite Dimensional Calculus* (Kluwer Academic Publishers, 1993).
22. H.-H. Kuo, *White Noise Distribution Theory* (CRC Press, 1996).
23. N. Obata, *White Noise Calculus and Fock Space* (Springer Verlag, 1996).
24. Y. G. Kondratiev, P. Leukert, and L. Streit, *Acta Appl. Math.* **44**, 269–294 (1996).
25. Y. G. Kondratiev and et al, *J. Funct. Anal.* **141**, 301–318 (1996).
26. H. P. Suryawan, *Int. J. Mod. Phys. Conf. Ser.* **36**, 1560005: 1–11 (2015).
27. C. Bender, *Stoch. Proc. Appl.* **104**, 81–106 (2003).
28. C. Bender, *Bernoulli* **9**, 955–983 (2003).
29. H. P. Suryawan, *J. Math. Fundam. Sci.* **51**, 265–277 (2019).
30. H.-H. Kuo, *Nonlinear Anal. Theory Methods Appl.* **30**, 317–328 (1997).

Turnitin AIP 2020

ORIGINALITY REPORT

23%

SIMILARITY INDEX

15%

INTERNET SOURCES

20%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1

ebin.pub

Internet Source

1%

2

export.arxiv.org

Internet Source

1%

3

nip.upd.edu.ph

Internet Source

1%

4

core.ac.uk

Internet Source

1%

5

Soukaina Douissi, Khalifa Es-Sebaiy, Soufiane Moussaten. "Asymptotics of the Cross-Variation of Young Integrals with Respect to a General Self-Similar Gaussian Process", Acta Mathematica Scientia, 2020

Publication

1%

6

Guangjun Shen. "Necessary and sufficient condition for the smoothness of intersection local time of subfractional Brownian motions", Journal of Inequalities and Applications, 2011

Publication

1%

7	Junfeng Liu, Donglei Tang, Yuquan Cang. "Variations and estimators for self-similarity parameter of sub-fractional Brownian motion via Malliavin calculus", Communications in Statistics - Theory and Methods, 2015 Publication	1 %
8	Karl Patrick S. Casas, Christopher C. Bernido. "From Markovian to non-Markovian stochastic processes: Diffusion coefficients with power law time dependence", AIP Publishing, 2020 Publication	1 %
9	P. J. Hunt, J. E. Kennedy. "Financial Derivatives in Theory and Practice", Wiley, 2004 Publication	1 %
10	N. A. Kachanovsky. "Stochastic integral of Hitsuda–Skorokhod type on the extended Fock space", Ukrainian Mathematical Journal, 06/2009 Publication	1 %
11	Submitted to Universitas Sanata Dharma Student Paper	1 %
12	www.cima.uevora.pt Internet Source	1 %
13	www.karyailmiah.trisakti.ac.id Internet Source	<1 %

14	Submitted to University of Strathclyde Student Paper	<1 %
15	hdl.handle.net Internet Source	<1 %
16	moam.info Internet Source	<1 %
17	arxiv.org Internet Source	<1 %
18	www.menaldi.wayne.edu Internet Source	<1 %
19	www.ripublication.com Internet Source	<1 %
20	123dok.com Internet Source	<1 %
21	B. L. S. Prakasa Rao. "On some maximal and integral inequalities for sub-fractional Brownian motion", Stochastic Analysis and Applications, 2016 Publication	<1 %
22	epdf.tips Internet Source	<1 %
23	eprints.upnyk.ac.id Internet Source	<1 %
24	pure.tue.nl Internet Source	<1 %

25 Xili Zhang, Weilin Xiao. "Arbitrage with fractional Gaussian processes", Physica A: Statistical Mechanics and its Applications, 2017

Publication

<1 %

26 Yong Xu, Rong Guo, Di Liu, Huiqing Zhang, Jinqiao Duan. "Stochastic averaging principle for dynamical systems with fractional Brownian motion", Discrete & Continuous Dynamical Systems - B, 2014

Publication

<1 %

27 journalofinequalitiesandapplications.springeropen.com

Internet Source

<1 %

28 Biagini, Francesca, Holger Fink, and Claudia Klüppelberg. "A fractional credit model with long range dependent default rate", Stochastic Processes and their Applications, 2013.

Publication

<1 %

29 Helge Holden, Bernt Øksendal, Jan Ubøe, Tusheng Zhang. "Chapter 2 Framework", Springer Science and Business Media LLC, 2010

Publication

<1 %

30 Said Ngobi. "A generalization of the Itô formula", International Journal of

<1 %

Mathematics and Mathematical Sciences,
2002

Publication

31

Stochastic Analysis and Applications in
Physics, 1994.

Publication

<1 %

32

publishing.aip.org

Internet Source

<1 %

33

www.lumii.lv

Internet Source

<1 %

34

Alberto Masayoshi F. Ohashi. "Stochastic
Evolution Equations Driven by a Fractional
White Noise", Stochastic Analysis and
Applications, 2006

Publication

<1 %

35

Benth, F.E.. "A White Noise Approach to a
Class of Non-linear Stochastic Heat
Equations", Journal of Functional Analysis,
19970601

Publication

<1 %

36

Constantin Tudor. "On the Wiener integral
with respect to a sub-fractional Brownian
motion on an interval", Journal of
Mathematical Analysis and Applications, 2009

Publication

<1 %

37

Martin Grothaus, Jan Müller, Andreas
Nonnenmacher. "An improved

<1 %

characterisation of regular generalised functions of white noise and an application to singular SPDEs", Stochastics and Partial Differential Equations: Analysis and Computations, 2021

Publication

38

epubs.siam.org

Internet Source

<1 %

39

repositorio.cinvestav.mx

Internet Source

<1 %

40

Constantin Tudor. "Some properties of the sub-fractional Brownian motion", Stochastics, 2011

Publication

<1 %

41

Guangjun Shen, Litan Yan, Chao Chen. "On the convergence to the multiple subfractional Wiener–Itô integral", Journal of the Korean Statistical Society, 2012

Publication

<1 %

42

Joachim Lebovits. "Stochastic Calculus with Respect to Gaussian Processes", Potential Analysis, 2017

Publication

<1 %

43

Litan Yan. "Some properties of the fractional Ornstein–Uhlenbeck process", Journal of Physics A Mathematical and Theoretical, 04/11/2008

Publication

<1 %

44 Nenghui Kuang, Huantian Xie. "Asymptotic behavior of weighted cubic variation of sub-fractional brownian motion", Communications in Statistics - Simulation and Computation, 2014
Publication <1 %

45 "Stochastic Differential Equations and Processes", Springer Science and Business Media LLC, 2012
Publication <1 %

46 Barndorff-Nielsen, Ole E., Fred Espen Benth, Jan Pedersen, and Almut E.D. Veraart. "On stochastic integration for volatility modulated Lévy-driven Volterra processes", Stochastic Processes and their Applications, 2014.
Publication <1 %

47 C. C. Bernido, M. V. Carpio-Bernido. "Preface", International Journal of Modern Physics: Conference Series, 2015
Publication <1 %

48 N. E. Frangos *, S. D. Vrontos, A. N. Yannacopoulos. "Ruin probability at a given time for a model with liabilities of the fractional Brownian motion type: A partial differential equation approach", Scandinavian Actuarial Journal, 2005
Publication <1 %

49 Samuel Daudin. "Optimal Control of Diffusion Processes with Terminal Constraint in Law", *Journal of Optimization Theory and Applications*, 2022

Publication

<1 %

50 Sergio Albeverio. "On the paths Hölder continuity in models of Euclidean quantum field theory", *Stochastic Analysis and Applications*, 1/1/2001

Publication

<1 %

51 Un Cig Ji, Nobuaki Obata. "Unitarity of Generalized Fourier–Gauss Transforms", *Stochastic Analysis and Applications*, 2006

Publication

<1 %

52 sutir.sut.ac.th:8080

Internet Source

<1 %

53 Kinda Abuasbeh, Nazim I. Mahmudov, Muath Awadalla. "Relative Controllability and Ulam–Hyers Stability of the Second-Order Linear Time-Delay Systems", *Mathematics*, 2023

Publication

<1 %

54 Lecture Notes in Physics, 2001.

Publication

<1 %

55 PIOTR MIŁOŚ. "OCCUPATION TIMES OF SUBCRITICAL BRANCHING IMMIGRATION SYSTEMS WITH MARKOV MOTION, CLT AND DEVIATION PRINCIPLES", *Infinite Dimensional*

<1 %

Analysis, Quantum Probability and Related Topics, 2012

Publication

56	ilirias.com Internet Source	<1 %
57	kuscholarworks.ku.edu Internet Source	<1 %
58	repository.kulib.kyoto-u.ac.jp Internet Source	<1 %
59	vmsta.org Internet Source	<1 %
60	www.numdam.org Internet Source	<1 %
61	www.worldscientific.com Internet Source	<1 %
62	Alberto Lanconelli. "Computing conditional expectations of multidimensional diffusion processes", <i>Stochastics</i> , 2006 Publication	<1 %
63	G. N. Gatica. "On the numerical analysis of nonlinear twofold saddle point problems", <i>IMA Journal of Numerical Analysis</i> , 04/01/2003 Publication	<1 %
64	M. Ait Ouahra, S. Moussaten, A. Sghir. "On limit theorems of some extensions of	<1 %

fractional Brownian motion and their additive functionals", Stochastics and Dynamics, 2017

Publication

65

SERGIO ALBEVERIO, MINORU W. YOSHIDA. "HIDA DISTRIBUTION CONSTRUCTION OF NON-GAUSSIAN REFLECTION POSITIVE GENERALIZED RANDOM FIELDS", Infinite Dimensional Analysis, Quantum Probability and Related Topics, 2011

<1 %

Publication

66

Selesi, D.. "Fundamental solutions of singular SPDEs", Chaos, Solitons and Fractals, 201107

<1 %

Publication

67

YURI G. KONDRATIEV, EUGENE W. LYTVYNOV. "OPERATORS OF GAMMA WHITE NOISE CALCULUS", Infinite Dimensional Analysis, Quantum Probability and Related Topics, 2011

<1 %

Publication

68

"Séminaire de Probabilités XLV", Springer Science and Business Media LLC, 2013

<1 %

Publication

69

B. L. S. Prakasa Rao. "Fractional Processes and Their Statistical Inference: An Overview", Journal of the Indian Institute of Science, 2022

<1 %

Publication

70

Mounir Zili, Eya Zougar. "Mixed stochastic heat equation with fractional Laplacian and

<1 %

gradient perturbation", Fractional Calculus and Applied Analysis, 2022

Publication

71

Tudor, C.. "Inner product spaces of integrands associated to subfractional Brownian motion", Statistics and Probability Letters, 20081001

Publication

<1 %

72

Xavier Bekaert, Nicolas Boulanger. "Gauge invariants and Killing tensors in higher-spin gauge theories", Nuclear Physics B, 2005

Publication

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On