

# UV-Vis Spectroscopy to Enable Determination of the Dissolution Behavior of Solid Dispersions Containing Curcumin as Well as Piperine

*by Yustina Sri Hartini*

---

**Submission date:** 15-Nov-2021 08:47PM (UTC+0700)

**Submission ID:** 1703413347

**File name:** of\_Young\_Pharmacist\_Scopus\_Yosi\_et\_al\_Yustina\_second\_auctor.pdf (325.92K)

**Word count:** 5010

**Character count:** 27427

5

# UV-Vis Spectroscopy to Enable Determination of the Dissolution Behavior of Solid Dispersions Containing Curcumin as Well as Piperine

Yosi Bayu Murti<sup>1</sup>, Yustina Sri Hartini<sup>2</sup>, Wouter Leonardus Joseph Hinrichs<sup>3</sup>, Henderik Willem Frijlink<sup>3</sup>, Dewi Setyaningsih<sup>2\*</sup><sup>1</sup>Faculty of Pharmacy, Universitas Gadjah Mada, INDONESIA.<sup>2</sup>Faculty of Pharmacy, Sanata Dharma University, INDONESIA.<sup>3</sup>Department of Pharmaceutical Technology and Biopharmacy, University of Groningen, THE NETHERLANDS.

## ABSTRACT

**Objective:** Curcumin and piperine exhibit poor aqueous solubility and dissolution. Solid dispersion technology is a promising way to improve dissolution of lipophilic compounds. Therefore, to investigate the dissolution behavior, an accurate determination of curcumin and piperine concentrations in a sample containing 0.5 wt-% SLS in sodium phosphate buffer as the dissolution medium was required. The goal of research was to investigate whether UV-Vis spectroscopy can be used to accurately determine curcumin and piperine concentrations in aqueous solutions. **Method:** Stock solutions (1 mg/mL) of curcumin and piperine were prepared in methanol. A series of solutions for calibration were prepared by mixing the stock solutions in various ratios after which they were diluted with dissolution medium (0.5% wt sodium lauryl sulfate in 20 mM phosphate buffer, pH 6.0). Solutions of curcumin (2 µg/mL) and piperine (1 µg/mL) were subjected to overlay scan in a UV-VIS spectrophotometer in 20 min.  $\lambda_{max}$  of 430 and 335.5 nm for curcumin and piperine, respectively. The method was validated according to ICH requirements, such as specificity, linearity, accuracy, precision, limit of detection and limit of quantification. **Results:**

At a concentration range of 0.1 to 5 µg/mL, calibration curves of curcumin and piperine showed linearity with  $R^2 = 0.9980$  and  $0.9982$ , respectively. Precision and accuracy was confirmed by AOAC. Furthermore, LOD of both compounds was 0.23 µg/mL and LOQ of curcumin and piperine was 0.72 and 0.69 µg/mL, respectively. **Conclusion:** UV-Vis spectroscopy can be used to accurately determine curcumin and piperine concentrations in dissolution samples obtained during screening the dissolution behavior of solid dispersions containing these compounds.

**Key words:** Curcumin, Piperine, Dissolution, UV-Vis spectroscopy.

### Correspondence

Dewi Setyaningsih, Faculty of Pharmacy, Sanata Dharma University, INDONESIA.

Phone: +6282135993494

Email: dewi@usd.ac.id

DOI: 10.5530/jyp.2019.11.x

## INTRODUCTION

Curcumin is the principle component of turmeric rhizome and exhibits various pharmacological activities, such as anti-inflammatory, antioxidant, hepatoprotector, antibacterial and anticancer activity.<sup>1-3</sup> Despite having excellent biological activity *in vitro* and *in vivo*, clinical application of this natural compound as a pharmaceutical agent is hampered because of its poor bioavailability after oral administration.<sup>4,5</sup> The poor bioavailability of curcumin can be ascribed to two reasons. First, curcumin exhibits an aqueous solubility of only 11 ng/mL,<sup>6</sup> causing poor dissolution behavior in the gastrointestinal tract, by which only limited amount of dissolved molecules are available for absorption. E.g. in a publication of Wahlström *et al.* it was reported that 75% of the administered dose of curcumin was found in faeces indicating that curcumin was poorly dissolved in the gastrointestinal milieu.<sup>7</sup> Second, once dissolved, curcumin is rapidly metabolized into curcumin glucuronate by UDP-glucuronosyltransferase.<sup>8</sup>

In Indonesian traditional medicine, a combination of turmeric and black pepper extract has been used in therapy.<sup>9</sup> It was believed that additional of black pepper extract, which predominately contains piperine, enhances the efficacy of turmeric extract. Scientific research has meanwhile supported the rational basis of the enhanced efficacy of the combination: co-administration of piperine enhances curcumin's bioavailability.<sup>10</sup> Another study showed that piperine is a UDP-glucuronosyltransferase inhibitor in both the small intestine and liver.<sup>11</sup> Furthermore, piperine

has been reported to increase bioavailability as it also acts as a P-glycoprotein inhibitor in the gastrointestinal tract. Also a synergistic effect of piperine-curcumin has been found. An *in vivo* study in rats revealed that co-administration with piperine improved the anti-inflammatory effect. In addition, piperine alone demonstrated anti-inflammatory, analgesic, antineoplastic, axiolytic activity and stimulator of digestive enzyme.<sup>12</sup> Similar to curcumin, piperine also exhibits a poor aqueous solubility. Therefore, in studying such combination the low aqueous solubility of both of these two compounds needs to be addressed. Solid dispersion technology is an often applied strategy to improve the dissolution behaviour of poorly soluble drugs.<sup>13</sup> Therefore, in order to deal with the low aqueous solubility and to prepare a synergistic combination, which is expected to improve curcumin bioavailability in the pre-clinical study, we prepared microparticle based solid dispersion containing both curcumin and piperine.

To evaluate this strategy, studying the *in vitro* dissolution behavior being the first step in screening formulations. To be able to do so, an accurate and reliable analytical method is needed to determine both curcumin and piperine concentration. Furthermore, since dissolution experiments often involve large number of samples, rapid and economical analytical method is required.

Several analytical methods have been reported to quantify curcumin in the co-mixed curcumin-piperine. As in a methanolic solution, piperine

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

and curcumin exhibit a  $\lambda_{\max}$  of 341 and 420 nm, respectively, it can be expected that both compounds will interfere each other during analysis by spectrophotometry method as the peaks may overlap. Therefore, the reported analytical methods are based on chromatography, i.e. Liquid Chromatography-Mass Spectrometry (LC-MS),<sup>14</sup> High Performance Liquid Chromatography (HPLC),<sup>10,15,16</sup> Ultra-Fast Liquid Chromatography<sup>17</sup> and High Performance Thin Layer Chromatography (HPTLC).<sup>18</sup> Although chromatographic method is suitable for selectively determine both curcumin and piperine concentrations, the reported methods require laborious sample extraction procedures by which it may potentially affect negatively to the samples since curcumin degradation may occur due to inevitably light exposure.<sup>19</sup> However, UV-Vis spectroscopic method has been reported to be suitable to determine concentration of curcumin/piperine/querctin in an Ayurvedic polyherbal formulation.<sup>20</sup> Using a particular calculation involving the absorptivity of the tested compounds, the concentration of the compounds in a co-mixture could be accurately determined.<sup>21</sup> In the current study, we aimed at validating a spectroscopic method for curcumin/piperine quantification from samples collected directly from the dissolution experiment where dissolution medium is present, which differs from the sample reported before.<sup>21</sup> Therefore, the novelty of this study is in developing a spectroscopic based analytical method to estimate curcumin and piperine in dissolution medium using combinatorial equation.

## MATERIALS AND METHODS

### Materials

Curcumin as reference standard was purchased from Nacalai, Japan. *C. longa* standardized extract (97.56%) was supplied by PT Phytochemindo Reksa, Bogor, Indonesia. Piperine was isolated and purified as described before,<sup>22,23</sup> after which it was further characterized based on piperine standard (Sigma Aldrich) using Thin Layer Chromatography method. Sodium dihydrogen phosphate and sodium lauryl sulfate (SLS) were purchased from Merck, Darmstadt, Germany. Mili Q water was prepared in our laboratory. All solvents were of analytical grade and were purchased from Merck, Darmstadt, Germany. Dissolution medium consisting of 0.5 wt-% SLS in 20 mM sodium phosphate buffer of pH 6.0 was prepared in our laboratory.

### Determination of $\lambda_{\max}$

Solutions of curcumin (2  $\mu\text{g/mL}$ ) and piperine (1  $\mu\text{g/mL}$ ) prepared in dissolution medium was subjected to overlay scan in a UV spectrophotometer from 200-600 nm. It was confirmed that  $\lambda_{\max}$  for curcumin and piperine are of 430 and 335.5 nm, respectively. Additionally, as a control, solutions of curcumin and piperine in methanol at the same concentrations were also scanned. The  $\lambda_{\max}$  for curcumin and piperine in methanol was 420 and 341 nm.

### Preparation of calibration curve

Curcumin or piperine stock solutions were prepared in methanol at a concentration of 1.0 mg/mL. Series of solutions containing curcumin and piperine at various concentrations were prepared by spiking the dissolution medium with the stock solutions. The calibration curves were prepared at concentrations of 0.05 - 5  $\mu\text{g/mL}$  for curcumin and piperine, respectively. Curcumin and piperine were simultaneously analyzed. Analyte concentrations were calculated based on combination calculation from the two  $\lambda$  measurements. Linearity of concentration and absorption and significance of the regressions were analyzed using Least Square Analysis method. The experiments were done in triplicate.

### Simultaneous equation for determination of curcumin/piperine

The equations (equation 1, 2) were derived based on the absorption of curcumin and piperine at the each maximum wavelength. Two wave-

lengths were selected (335 nm for curcumin and 430 nm for piperine) to calculate the concentrations based on equations 1 and 2. Furthermore, the absorptivity values E (1%, 1 cm) were determined for the piperine and curcumin at their maximum wavelength.

$$C_c = \frac{(A2.ap1) - (A1.ap2)}{(ac2.ap2) - (ac1.ap2)} \quad \text{equation 1}$$

$$C_p = \frac{(A1.ac2) - (A2.ac1)}{(ac2.ap2) - (ac1.ap2)} \quad \text{equation 2}$$

Cc: curcumin concentration

Cp: piperine concentration

A1: absorbance measured at  $\lambda$  335.5 nm

A2: absorbance measured at  $\lambda$  430.0 nm

Ac1: curcumin absorptivity at  $\lambda$  335.5 nm in absorbance/ppm

Ac2: curcumin absorptivity at  $\lambda$  430.0 nm in absorbance/ppm

Ap1: piperine absorptivity at  $\lambda$  335.5 nm in absorbance/ppm

Ap2: piperine absorptivity at  $\lambda$  430.0 nm in absorbance/ppm

### Method Validation

The spectroscopic method for quantification of curcumin and piperine in the sample solution was validated according Q2R1 ICH guidelines, which are linearity, accuracy and precision, limit of detection (LOD) and limit of quantification (LOQ).<sup>24</sup> Linearity between concentrations and absorbance was assessed by analyzing the series concentration of calibration plot using Least Square Analysis. Correlation coefficient (r) of >0.99 was used to determine the linearity.<sup>25</sup> For accuracy and precision studies, three different concentrations of curcumin and piperine in a mixture solution were prepared. Nine different samples were prepared at concentrations of 1, 2, 4  $\mu\text{g/mL}$  curcumin and 0.5, 1, 2  $\mu\text{g/mL}$  piperine in the dissolution medium. Accuracy was expressed as recovery value while precision was reflected by relative standard deviation (RSD) value. Inter-day precision and accuracy were determined at another 2 consecutive days. RSD were obtained by dividing the standard deviation with the mean value of obtained concentrations  $\times$  100%. Recovery was calculated by ratio of obtained concentration to the accordingly nominal concentration  $\times$  100%. Linearity, accuracy and precision were judged according to AOAC requirements.<sup>25</sup> LOD and LOQ were determined based on standard deviation of response and slope as described previously.<sup>26</sup> LOD and LOQ were defined as  $3.3 \sigma / S$  and  $10 \sigma / S$ , respectively, where  $\sigma$  is the standard error of intercept and S represents the mean of slopes as determined by the Least Square Analysis.

### Application of the validated method

The validated method was applied to quantify curcumin and piperine concentrations after dissolution of a solid dispersion formulation containing both components. The solid dispersion were prepared by spray drying method using polyvinylpyrrolidone K30 (PVP K30) as carrier. The solid dispersion contained 30 wt-% curcuminoid, 1 wt-% piperine. In brief, curcumin, piperine and PVP K30 were dissolved in ethanol then spray dried using Büchi B-290 spray dryer equipped with a B-295 dehumidifier. Atomizing air was set at a flow 600 Ln/h. Inlet temperature and aspirator was set at 100°C and 100%, respectively. These setting resulted in an outlet temperature of 49-55°C.

The obtained spray dried powder were weighed accurately (100.00 mg) and then dissolved in 25.0 mL dissolution medium. A volume of 40  $\mu\text{L}$  was taken and diluted to 5.0 mL to achieve final concentrations of curcumin and piperine of 9.60 and 0.32  $\mu\text{g/mL}$ . The absorbance of the diluted sample was measured at 335.5 and 430 nm for piperine and



curcumin, respectively. The measurement was done in triplicate. The concentration of curcumin and piperine was calculated using equation 1 and 2.

## RESULTS

### Spectral scanning


Spectral scanning of curcumin, piperine and the mixture of curcumin-piperine solution are presented in Figure 1. In the methanolic solutions (Figure 1a),  $\lambda_{\max}$  of single compound of curcumin and piperine appeared to be 420.5 nm and 341 nm, respectively. Slight bathochromic shifts were observed when curcumin-piperine was co-mixed to 421 and 345 nm for curcumin and piperine, respectively. When medium dissolution was used to dissolve curcumin or piperine (Figure 1b),  $\lambda_{\max}$  of curcumin solution appeared to be bathochromic shifted to 430 with respect to  $\lambda_{\max}$  of its methanolic solution (420.5nm). Single piperine solution in medium dissolution appeared to exhibit a hypsochromic shift (335.5 nm) when compared to its methanolic solution (341.5 nm). A slight bathochromic shift was observed in a co-mixed curcumin-piperine in dissolution medium from its single piperine solution in dissolution medium (Figure 1b). From the scanning peaks data (Figure 1b) it is obvious that there was no interference of piperine peak at the  $\lambda_{\max}$  of curcumin. However, when the piperine concentration was determined using  $\lambda_{\max}$  of 335.5 nm, curcumin did interfere significantly. Our data showed that curcumin demonstrates significant absorbance at 0.05 at  $\lambda_{\max}$  of 535.5 nm when the mixture was prepared at ratio curcumin: piperine 1:1 (Figure 1a and 1b). Therefore, to minimize or to avoid the interference of other compound in the determination of piperine concentrations, calculation based on combinatorial equations as presented in equation 2 has to be conducted.

### Method validation


The validation of the spectroscopic method for determination of curcumin-piperine in a dissolution sample followed the procedures as described by ICH.<sup>24</sup>

### Linearity and range


The calibration curve obtained was evaluated by its correlation coefficient (r) and coefficient of determination ( $R^2$ ). To construct calibration curves, 17 different concentrations of co-mixed of curcumin and piperine were prepared in dissolution medium. The concentrations ranged from



**Figure 1:** Spectral of curcumin, piperine and curcumin-piperine mixture prepared in methanol (a) and dissolution medium (b). Maximum wavelength shifts were observed when curcumin or piperine or the mixture of both were dissolved in dissolution medium.



**Figure 2:** Regression analysis of curcumin concentrations (ppm) vs absorbance at wavelength 1 (a) and 2 (b) with  $n = 3$ .



**Figure 3:** Regression analysis of piperine concentrations (ppm) vs absorbance at wavelength 1 (a) and 2 (b) with  $n = 3$ .

**Table 1:** Validation parameters for curcumin and piperine prepared in dissolution medium ( $n= 3$ ).

Parameter	Curcumin	Piperine
$\lambda_{\max}$ (nm)	430.0	335.5
Linear equation	$y = 0.02266x - 0.00049$	$y = 0.076598x - 0.001014$
Slope	0.139807	0.076598
$R^2$	0.9980	0.9982
r	0.9989	0.9990
LOD ( $\mu\text{g/mL}$ )	0.23	0.23
LOQ ( $\mu\text{g/mL}$ )	0.72	0.69

0.005 – 5  $\mu\text{g/mL}$ . The calibration curves of curcumin and piperine are depicted in Figure 2 and Figure 3. Linearity of curcumin or piperine was obtained from 10 different concentrations which ranged from 0.1 to 5.0  $\mu\text{g/mL}$  as indicated by r and  $R^2$  of 0.9989 and 0.9980, respectively, for curcumin and 0.9990 and 0.9982, respectively, for piperine solutions (Table 1).<sup>25</sup> Moreover, regression analysis (Figure 2 and Figure 3) shows the regression was significant ( $p < 0.05$ ) which can be used as prediction model for the determination of curcumin and piperine concentration in the co-mixture solution. The LOD and LOQ were calculated based on the calibration curves of curcumin and piperine. The calculated LOD of curcumin and piperine was 0.23  $\mu\text{g/mL}$  and the calculated LOQ values were 0.69 and 0.72 for piperine and curcumin, respectively.

### Precision and accuracy

Precision was evaluated at three different concentrations prepared in dissolution medium, which were 0.5, 1 and 2  $\mu\text{g/mL}$  for piperine and 1, 2, 4  $\mu\text{g/mL}$  for curcumin. Precision and accuracy data are presented in Table 2 and

Table 3. Intra-day and inter-day precision study for curcumin and piperine of the method shows RSD values between 0.48 – 10.70, which confirms

**Table 2: Intraday and interday precision and accuracy of piperine (n=3).**

Nominal concentration (µg/mL)	Intraday			Interday		
	Found (µg/mL)	Precision (%RSD)	Accuracy (%recovery)	Found (µg/mL)	Precision (% RSD)	Accuracy (% recovery)
0.5	0.54	1.67	107.38	0.51	2.87	102.69
1.0	1.02	10.7	102.00	1.01	8.82	101.00
2.0	2.09	0.48	104.50	2.05	1.73	102.50

**Table 3: Intraday and interday precision and accuracy of curcumin (n=3).**

Nominal concentration (µg/mL)	Intraday			Interday		
	Found (µg/mL)	Precision (%RSD)	Accuracy (%recovery)	Found (µg/mL)	Precision (% RSD)	Accuracy (% recovery)
1.0	1.03	9.64	103.00	0.99	7.29	96.84
2.0	2.09	2.67	104.50	1.98	2.14	94.87
4.0	4.22	5.62	105.50	4.05	3.45	95.40

reliability and stability of the method based on AOAC requirements.<sup>25</sup> Accuracy data as indicated by the recovery values shows recovery of 94-107% which confirms the validity of the method according to the AOAC requirements.<sup>25</sup>

#### Application of the validated method

The validated method was applied to measure the concentration of curcumin and piperine dissolved in the dissolution medium. The sample dilution preparation was conducted as such mimicking the concentration found at the initial time sampling. Considering the dose of 500 mg curcuminoids and 5 mg of piperine in a tablet as conducted in a clinical trial<sup>10</sup> and of at least 5% analyte should be released in 1000 mL of dissolution medium during earlier sampling, it is required to be able to measure concentrations of 25 µg/mL curcumin and 0.25 µg/mL of piperine accurately. Using this validated spectroscopic method, curcumin concentration was found to be  $10.11 \pm 0.09$  µg/mL with RSD value of 0.89% while the piperine concentration was found to be  $0.33 \pm 0.06$  µg/mL with RSD value of 18.18%. Taking the expected concentrations of 25 and 0.25 µg/mL for curcumin and piperine during initial time sampling and the found concentrations of  $10.11 \pm 0.09$  µg/mL (curcumin) and  $0.33 \pm 0.06$  µg/mL (piperine), it can be concluded that this validated spectroscopic method is sensitive enough to simultaneously quantify curcumin and piperine in the dissolution sample. Although we prepared a slightly higher concentration of piperine in the dissolution sample (around 0.32 µg/mL) than the expected concentration (0.25 µg/mL), we were still able to confirm the sensitivity of the method based on the found LOD (0.23 µg/mL). In the sample, the RSD of piperine was quite high, which was 18.18%. The RSD value of piperine was high due to the very low piperine concentration approaching at detection level of the method (0.23 µg/mL).

## DISCUSSION

The aim of this study was to develop and validate a spectroscopic method to determine curcumin and piperine in dissolution samples. As a BCS II drug, dissolution behavior of curcumin and piperine is being the most important predictive parameter for the bioavailability.<sup>27</sup> In screening of a formula candidate, dissolution behavior of a drug is the most useful parameters, which it may end up with large number of dissolution samples and those need quick analysis to allow rapid selection of formula for bio-availability evaluation.

Analytical method validation objective is to demonstrate suitability and reliability of an analytical procedure for its intended purpose.<sup>28</sup> This study validated the spectroscopic method to quantify curcumin and piperine in the dissolution medium. The dissolution medium contained 0.5 wt-% SLS in 20 mM sodium phosphate buffer of pH 6.0. Sodium phosphate buffer at pH 6.0 was used to confirm curcumin stability during dissolution study, since curcumin was found to be stable at pH 6.0.<sup>6</sup> SLS was added in the dissolution medium to attain sink condition.<sup>29</sup> Although chromatographic methods are found in many publications for accurately determine curcumin-piperine, SLS and sodium phosphate buffer in the dissolution media may cause poor peak separation in chromatographic method. Collier *et al.* reported that SLS in the dissolution media interferes the chromatographic separation of levothyroxine as indicated by a dramatic decrease in plate counts after number of samples injection into the HPLC column, and that resulted a high variability in drug content in the dissolution samples.<sup>30</sup> Recently, Setyaningsih *et al.* found that SLS and sodium phosphate buffer cause peak tailing of curcuminoids resulting poor peak separation.<sup>31</sup> The interference caused by SLS and sodium phosphate buffer leads to jeopardize accurate estimation of curcumin in dissolution samples.

Sample preparation is required to obtain satisfactory chromatographic peaks separations. However, extensive sample preparation, which may involve multiple clean up steps followed by extraction procedures, might raise stability concerns of curcumin. A report from Tonnesen and Karlson suggests that when using buffers and ethyl acetate for extraction, polymerization of curcumin can occur at the interface of the two phases.<sup>19</sup>

Although spectroscopic method has been acknowledged as not selective; there is no ability of the method to determine curcumin from curcuminoids mixture, as these curcuminoids appears to be overlapping in UV spectra at the same wavelength,<sup>30</sup> the method are useful to determined curcumin concentration in dissolution study. Despite the selectivity problem, the spectrophotometric method offers simplicity, rapidity and reliability.<sup>32,33</sup> Being the major component in curcuminoids mixture, curcumin concentration in samples can be determined as total curcuminoids which are present in the dosage forms or polyherbal samples.<sup>20</sup> Taken all together, spectrophotometry remains valuable method to determine curcumin in the dissolution samples. Furthermore, using combinatorial equation involving absorptivity molar and absorbance of curcumin and piperine (equation 1 and 2), this developed method are able to reliably determine curcumin in co-existence with piperine



2

This proposed spectroscopic method confirms linearity, accuracy and precision of the method as required by AOAC. The LOD of piperine and curcumin was 0.23 µg/mL and LOQ were 0.69 and 0.72 for piperine and curcumin, respectively. As compared when methanol was used to prepare samples, Aneja *et al* found LOD of 0.71 µg/mL and 0.45 µg/mL and LOQ of 2.1 µg/mL and 1.49 µg/mL for piperine and curcumin, respectively,<sup>21</sup> which indicates that this validated method of curcumin/piperine quantification in dissolution medium is more sensitive compared to the previous report.

## CONCLUSION

The UV-Vis spectroscopy method is accurate, precise and reliable to be used for the quantification of curcumin and piperine in dissolution medium composed of 0.5 wt-% SLS in 20 mM sodium phosphate buffer of pH 6.0 where curcumin and piperine are co-present. Moreover, the LOD shows to be sensitive. Thus validation confirmed that the method is applicable for the determination of curcumin or piperine during dissolution studies.

## ACKNOWLEDGEMENT

This study is fully supported by the Ministry of Research, Technology and Higher Education of Republic Indonesia through the scheme of International Research Collaboration and Scientific Publication year of 2018 (3/E/KPT/2018).

3

## CONFLICT OF INTEREST

There are no conflict of interest in this manuscript.

## ABBEVIATION

**BCS:** Biopharmaceutical classification system; **LOD:** Limit of detection; **LOQ:** Limit of Quantification; **RSD:** Relative standard deviation; **SLS:** Sodium lauryl sulfate; **UV-Vis:** Ultraviolet-visible.

## REFERENCES

- Aggarwal B, Surh YJ, Shishodia S, Menon V, Sudheer A. Antioxidant and anti-inflammatory properties of curcumin The Molecular Targets and Therapeutic Uses of Curcumin in Health and Disease: Springer US. 2007;105-25.
- Kapoor S. Curcumin and Its Emerging Role in Pain Modulation and Pain Management. Korean J Pain. 2012;25(3):202-3.
- Abdollahi E, Momtazi AA, Johnston TP, Sahebkar A. Therapeutic effects of curcumin in inflammatory and immune-mediated diseases: A nature-made jack-of-all-trades?. J Cell Physiol. 2017. <http://dx.doi.org/10.1002/jcp.25778>
- Siviero A, Gallo E, Maggini V, Gori L, Mugelli A, Firenzuoli F, *et al*. Curcumin, a golden spice with a low bioavailability. J Herb Med. 2015;5(2):57-70.
- Hatcher H, Planalp R, Cho J, Torti FM, Torti SV. Curcumin: From ancient medicine to current clinical trials. Cell Mol Life Sci. 2008;65(11):1631-52.
- Wang YJ, Pan MH, Cheng AL, Lin LI, Ho YS, Hsieh CY, *et al*. Stability of curcumin in buffer solutions and characterization of its degradation products. J Pharm Biomed Anal. 1997;15(12):1867-76.
- Wahlström B, Blennow G. A study on the fate of curcumin in the rat. Acta Pharmacol Toxicol. 1978;43(2):86-92.
- Holder GM, Plummer JL, Ryan AJ. The metabolism and excretion of curcumin (1,7-bis-(4-hydroxy-3-methoxyphenyl)-1,6-heptadiene-3,5-dione) in the rat. Xenobiotica. 1978;8(12):761-8.
- Riswan S, Sangat-Roemantyo H. Jamu as traditional medicine in Java, Indonesia. South Pacific Study. 2002;23(1):1-10.
- Shoba G, Joy D, Joseph T, Majeed M, Rajendran R, Srinivas PS. Influence of Piperine on the Pharmacokinetics of Curcumin in Animals and Human Volunteers. Planta Med. 1998;64(4):353-6.

- Volak LP, Ghirmai S, Cashman JR, Court MH. Curcuminoids inhibit multiple human cytochromes P450, UDP-glucuronosyltransferase and sulfotransferase enzymes, whereas piperine is a relatively selective CYP3A4 inhibitor. Drug Metab Dispos. 2008;36(8):1594-605.
- Zhou S, Lim LY, Chowbay B. Herbal modulation of P-glycoprotein. Drug Metab Rev. 2004;36(1):57-104.
- Srinarong P, de Waard H, Frijlink HW, Hinrichs WL. Improved Dissolution Behavior of Lipophilic Drugs by Solid Dispersions: the Production Process as Starting Point for Formulation Considerations. Expert Opin Drug Deliv. 2011;8(9):1121-40.
- Tu YS, Fu JW, Sun DM, Zhang JJ, Yao N, Huang DE, *et al*. Preparation, characterization and evaluation of curcumin with piperine-loaded cubosome nanoparticles. J Microencapsulation. 2014;31(6):551-9.
- Moorthi C, Krishnan K, Manavalan R, Kathiresan K. Preparation and characterization of curcumin-piperine dual drug loaded nanoparticles. Asian Pac J Trop Biomed. 2011;2(11):841-8.
- Grill A, Koniar B, Panyam J. Co-delivery of natural metabolic inhibitors in a self-microemulsifying drug delivery system for improved oral bioavailability of curcumin. Drug Deliv Trans Res. 2014;4(4):344-52.
- Ramaswamy S, Kuppuswamy G, Dwarampudi P, Kadiyala M, Menta L, Kannan E. Development and validation of simultaneous estimation method for curcumin and piperine by RP-UFLC. Pak J Pharm Sci. 2014;27(4):901-6.
- Verma J, Joshi A. Rapid HPTLC method for identification and quantification of curcumin, piperine and thymol in an ayurvedic formulation. JPC J Planar Chromat. 2006;19(11):398-400.
- Tonnesen H, Karlsen J. Studies on curcumin and curcuminoids VI. Kinetics of curcumin degradation in aqueous solution. Z Lebensm Unters Forsch. 1985;180(5):402-4.
- Singh A, Avupati VR. Development and Validation of UV-Spectrophotometric method for the Estimation of Curcumin in Standardised Polyherbal Formulations. J Young Pharm. 2017;9(4):491-5.
- Aneja G, Dave U, Vadodaria K. Simultaneous Estimation of Piperine, Quercetin and Curcumin in A Mixture Using UV-Visible Spectrophotometer and Method Validation. IJTA. 2012;8:14-7.
- Shingate PN, Dongre PP, Kannur DM. New Method Development for Extraction and Isolation of Piperine from Black Pepper. International JPSR. 2013;4(8):3165-70.
- Saha KC, Seal HP, Noor MA. Isolation and characterization of piperine from the fruits of black pepper (*Piper nigrum*). J Bangladesh Agril Univ. 2013;11(1):11-6.
- International Conference on Harmonization (ICH). ICH Harmonised tripartite guideline. Validation of analytical procedures: Text and methodology ed: Geneva. 2005.
- The Association of Official Analytical Chemists (AOAC). Official Methods of Analysis of AOAC International. In: George W. Latimer J, editor. Guidelines for Standard Method Performance Requirements. 19<sup>th</sup> ed. Gaithersburg, Maryland, USA: AOAC International. 2012;9.
- González G, Herrador A. A practical guide to analytical method validation, including measurement uncertainty and accuracy profiles. Trends Analyt Chem. 2007;26(3):227-38.
- Hu L, Shi Y, Li JH, Gao N, Ji J, Niu F, *et al*. Enhancement of Oral Bioavailability of Curcumin by a Novel Solid Dispersion System. AAPS PharmSciTech. 2015;16(6):1327-34.
- Food and Drug Administration (FDA). Analytical Procedures and Methods Validation for Drugs and Biologics-Guidance for Industry. Division of Drug Information Center for Drug Evaluation and Research Food and Drug Administration. 2015;1-13.
- Visser MR, Baert L, Klooster G, Schueller L, Geldof M, Vanwelkenhuysen I, *et al*. Inulin solid dispersion technology to improve the absorption of the BCS Class IV drug TMC240. Eur J Pharm Biopharm. 2009;74(2):233-8.
- Collier JW, Shah RB, Bryant AR, Habib MJ, Khan MA, Faustino PJ. Development and application of a validated HPLC method for the analysis of dissolution samples of levothyroxine sodium drug products. J Pharm Biomed Anal. 2011;54(3):433-8.
- Setyaningsih D, Murti YB, Fudholi A, Hinrichs WLJ, Mudjahid R, Martono S, *et al*. Validated TLC Method for Determination of Curcumin Concentrations in Dissolution Samples Containing *Curcuma longa* Extract. JIFI. 2016;14(2):147-57.
- Sharma K, Agrawal SS, Gupta M. Development and validation of UV spectrophotometric method for the estimation of curcumin in bulk drug and pharmaceutical dosage forms. Int J Drug Dev and Res. 2012;4(2):375-80.
- Mazzarino L, Belletini IC, Minatti E, Lemos-Senna E. Development and validation of a fluorimetric method to determine curcumin in lipid and polymeric nanocapsule suspensions. Braz J Pharm Sci. 2010;46(2):219-26.

**Article History:** Submission Date : xx-xx-xxxx; Revised Date : xx-xx-xxxx; Acceptance Date : xx-xx-xxxx.

**Cite this article:** Murti YB, Hartini YS, Hinrichs WLJ, Frijlink HW, Setyaningsih D. UV-Vis Spectroscopy to Enable Determination of the Dissolution Behavior of Solid Dispersions Containing Curcumin as Well as Piperine. J Young Pharm. 2019;11(1):x-x.

# UV-Vis Spectroscopy to Enable Determination of the Dissolution Behavior of Solid Dispersions Containing Curcumin as Well as Piperine

## ORIGINALITY REPORT

19%

SIMILARITY INDEX

13%

INTERNET SOURCES

16%

PUBLICATIONS

6%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://www.scribd.com">www.scribd.com</a> Internet Source	2%
2	Bhairy Srinivas, Shaikh Ashraf, Nalawade Vivek, Hirlekar Rajashree. "Development and validation of bivariate UV-visible spectroscopic method for simultaneous estimation of curcumin and piperine in their combined nanoparticulate system", Journal of Applied Pharmaceutical Science, 2021 Publication	1%
3	<a href="http://ijpsr.com">ijpsr.com</a> Internet Source	1%
4	<a href="http://fulltxt.org">fulltxt.org</a> Internet Source	1%
5	<a href="http://www.semanticscholar.org">www.semanticscholar.org</a> Internet Source	1%
6	Wisut Wichitnithad. "A simple isocratic HPLC method for the simultaneous determination	1%

of curcuminoids in commercial turmeric extracts", Phytochemical Analysis, 2009

Publication

---

7	Mona Elfiky, Amr M. Beltagi, Osama Abuzalat. "Selective modified stripping voltammetric sensor based on Ce-1,4-benzenedicarboxylic metal-organic frameworks porous nanoparticles for picomolar detection of curcumin", Journal of Electroanalytical Chemistry, 2021 Publication	1 %
8	<a href="http://www.ajpcr.com">www.ajpcr.com</a> Internet Source	1 %
9	Submitted to Taibah University Student Paper	1 %
10	<a href="http://academic.oup.com">academic.oup.com</a> Internet Source	<1 %
11	Mustafa Celebier, Mustafa Sinan Kaynak, Sacide Altinoz, Selma Sahin. "UV Spectrophotometric Method for Determination of the Dissolution Profile of Rivaroxaban", Dissolution Technologies, 2014 Publication	<1 %
12	<a href="http://www.tandfonline.com">www.tandfonline.com</a> Internet Source	<1 %
13	Excipient Applications in Formulation Design and Drug Delivery, 2015.	<1 %



14

Submitted to Liverpool John Moores University

Student Paper

<1 %

---

15

pharmacophorejournal.com

Internet Source

<1 %

---

16

Kim, Jae Kwang, Soo-Yun Park, Sun-Hwa Ha, Si Myung Lee, Sun Hyung Lim, Chang Yeon Yu, Seok-Cheol Suh, and Soon Jong Kweon.

"Stable Isotope Dilution Gas Chromatography–Time-of-Flight Mass Spectrometry Method for Determining Niacin in Rice", Cereal Chemistry, 2011.

Publication

<1 %

---

17

Maryam Karimi, Mohammad Mashreghi, Sara Shokooh Saremi, Mahmoud Reza Jaafari.

"Spectrofluorometric Method Development and Validation for the Determination of Curcumin in Nanoliposomes and Plasma", Journal of Fluorescence, 2020

Publication

<1 %

---

18

Zhenping Liu, Jingzhou Hou, Xianfeng Wang, Changjun Hou, Zhong Ji, Qiang He, Danqun Huo. "A novel fluorescence probe for rapid and sensitive detection of tetracyclines residues based on silicon quantum dots", Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy, 2020

<1 %

19

Lida Liu, Haiyung Cheng, Jamie J. Zhao, J.Douglas Rogers. "Determination of montelukast (MK-0476) and its S-enantiomer in human plasma by stereoselective high-performance liquid chromatography with column-switching", Journal of Pharmaceutical and Biomedical Analysis, 1997

Publication

<1 %

---

20

[doaj.org](http://doaj.org)  
Internet Source

<1 %

---

21

[tvst.arvojournals.org](http://tvst.arvojournals.org)  
Internet Source

<1 %

---

22

Bae - Chan Kim, Eunmi Ban, Jeong - Sook Park, Yun - Kyoung Song, Chong - Kook Kim. "Determination of Simvastatin in Human Plasma by Column - Switching HPLC with UV Detection", Journal of Liquid Chromatography & Related Technologies, 2009

Publication

<1 %

---

23

Natural Products, 2013.  
Publication

<1 %

---

24

[www.nature.com](http://www.nature.com)  
Internet Source

<1 %

---

25

Submitted to Universitas Riau  
Student Paper

<1 %

---

26

"3rd BBBB-Bosphorus International Conference on Pharmaceutical Sciences", European Journal of Pharmaceutical Sciences, 20091026

Publication

<1 %

27

Petropoulou, S.S.E.. "Gas chromatographic-tandem mass spectrometric method for the quantitation of carbofuran, carbaryl and their main metabolites in applicators' urine", Journal of Chromatography A, 20060303

Publication

<1 %

28

[pubmed.ncbi.nlm.nih.gov](http://pubmed.ncbi.nlm.nih.gov)

Internet Source

<1 %

29

Esther Pinilla-Peñalver, M. Laura Soriano, Gema M. Durán, Eulogio J. Llorent-Martínez, Ana M. Contento, Ángel Ríos. "Discrimination between nanocurcumin and free curcumin using graphene quantum dots as a selective fluorescence probe", Microchimica Acta, 2020

Publication

<1 %

30

Prerana Sethi, Virendra K. Dua, S. Mohanty, S. K. Mishra, Rajeev Jain, G. Edwards. "Development and Validation of a Reversed Phase HPLC Method for Simultaneous Determination of Curcumin and Piperine in Human Plasma for Application in Clinical Pharmacological Studies", Journal of Liquid

<1 %

# Chromatography & Related Technologies, 2009

Publication

---

31

[media.proquest.com](http://media.proquest.com)

Internet Source

<1 %

---

32

CYLENI R. A. SOUZA. "STABILITY OF CURCUMINOIB PIGMENTS IN MODEL SYSTEMS", Journal of Food Processing and Preservation, 11/1997

Publication

<1 %

---

33

Durgawati Patel, Puspendra Kumar, Swapnil Sharma, Jaya Dwivedi. "Analytical Methods for Metformin Estimation", Critical Reviews in Analytical Chemistry, 2017

Publication

<1 %

---

34

Jayant Verma, Anil Joshi. "Rapid HPTLC method for identification and quantification of curcumin, piperine and thymol in an ayurvedic formulation", Journal of Planar Chromatography – Modern TLC, 2006

Publication

<1 %

---

35

Vishal Sharad Chaudhari, Roshan M. Borkar, Upadhyayula Suryanarayana Murty, Subham Banerjee. "Analytical Method Development and Validation of Reverse-Phase High-Performance Liquid Chromatography (RP-HPLC) Method for Simultaneous Quantifications of Quercetin and Piperine in

<1 %



# Dual-Drug Loaded Nanostructured Lipid Carriers", Journal of Pharmaceutical and Biomedical Analysis, 2020

Publication

---

36 [pdfs.semanticscholar.org](https://pdfs.semanticscholar.org) <1 %  
Internet Source

---

37 [polimery.umw.edu.pl](https://polimery.umw.edu.pl) <1 %  
Internet Source

---

38 [turkjps.org](https://turkjps.org) <1 %  
Internet Source

---

39 [www.japsonline.com](https://www.japsonline.com) <1 %  
Internet Source

---

40 [www.ncbi.nlm.nih.gov](https://www.ncbi.nlm.nih.gov) <1 %  
Internet Source

---

41 [www.science.gov](https://www.science.gov) <1 %  
Internet Source

---

42 [www.thieme-connect.com](https://www.thieme-connect.com) <1 %  
Internet Source

---

43 Yuan, J.. "Simultaneous determination of six alkaloids and one monoterpene in rat plasma by liquid chromatography-tandem mass spectrometry and pharmacokinetic study after oral administration of a Chinese medicine Wuji Pill", Journal of Chromatography B, 20120501 <1 %  
Publication

---

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography On