

Volume 7 Issue 1 June 2019

pISSN 2302-1616
eISSN 2580-2909

BIOGENESIS

Jurnal Ilmiah Biologi

Chromosome Characterization of Festival Strawberry (*Fragaria x ananassa* D. var. Festival) Result of Polyploidization

Ganes Rizka Aristya, Rezeki Alyza

Contamination of *Escherichia coli* Drinking Water Refills on Drinking Water Depots in Malang City

Wiwik Kusmawati, Lia Rihayu

Genetic Homogeneity of Commerson's Anchovy (*Stolephorus commersonnii*) in Segara Anakan Cilacap Central Java Inferred from PCR-RFLP Markers

Agus Nuryanto, Rani Eva Dewi, Hendro Pramono

Effect of Methyl Jasmonate on Vegetative Growth and Formation of Potato Tuber (*Solanum tuberosum* L. var. Granola)

Triana Rohmawati, Kumala Dewi

Toxicity of Bioactive Compound from Endophytic Fungi Isolated from Red Ginger (*Zingiber officinale* var. rubrum) Utilizing Brine Shrimp Lethality Assay

Angga Prasetyo, Boy Rahardjo Sidhiarta, Yustina Sri Hartini, Exyupransia Mursyanti

Application of Indigenous AMF from ex-coal Mining Soil Combined with Phosphorus Fertilizers to Improved Oil Palm Seedling Growth (*Elaeis guineensis* Jacq.)

Made Deviani Duaja, Elis Kartika, Lizawati Lizawati

The Effect of *Chlorella vulgaris* on Lipid Profile Wistar Strain Rats (*Rattus norvegicus* Berkenhout, 1769) Under Induced Stress

Farah Nadia Karima, Mulyati Sarto

Notes on *Cyclanthera brachystachya* (Ser.) Cog.: Morphology, Ecology, and Its New Distribution

Muhammad Efendi, Ruyayah Ruyayah

Geometric Morphometrics Analysis of Chelae and Carapace of the Freshwater Prawn *Macrobrachium* Bate, 1868

Rich Gemilang Simanjuntak, Rury Eprilurafman

The Potential of Breadfruit Seed and Jackfruit Seed as Alternative Replacement Medium of Potato Dextrose Agar (PDA) with Seedling F0 Mushrooms

Suparti Suparti, Lina Agustina, Putri Agustina, Resti Rahmawati

**DEPARTMENT OF BIOLOGY
FACULTY OF SCIENCE AND TECHNOLOGY
UNIVERSITAS ISLAM NEGERI (UIN) ALAUDDIN
M A K A S S A R**

Volume 7	Issue 1	June 2019	Page 1-72	pISSN 2302-1616 eISSN 2580-2909
-------------	------------	--------------	--------------	------------------------------------



[Home](#) [About](#) [Current](#) [Archives](#) [Announcements](#) [Statistics](#) [Indexing and Accreditation](#) [Sitemap](#) [Contact](#)

[Home](#) / [Archives](#) / Vol. 7 No. 1 (2019)

Vol. 7 No. 1 (2019)



Published: 2019-06-23

Research Articles

Chromosome Characterization of Festival Strawberry (*Fragaria x ananassa* D. var. Festival) Result of Polyploidization

Ganies Riza Aristya, Rezika Alyza

1-8



<https://doi.org/10.24252/bio.v7i1.5250>



Views: 693 / PDF Downloads: 189

Contamination of *Escherichia coli* Drinking Water Refills on Drinking Water Depots in Malang City

Wiwik Kusmawati, Lia Rahayu

9-13



<https://doi.org/10.24252/bio.v7i1.5786>



Views: 593 / PDF Downloads: 136 / PDF Downloads: 0

Genetic Homogeneity of Commerson's Anchovy (*Stolephorus commersonii*) in Segara Anakan Cilacap Central Java Inferred from PCR-RFLP Markers

Agus Nuryanto, Rani Eva Dewi, Hendro Pramono

14-23



<https://doi.org/10.24252/bio.v7i1.6352>



Views: 418 / PDF Downloads: 104

Effect of Methyl Jasmonate on Vegetative Growth and Formation of Potato Tuber (*Solanum tuberosum* L. var. Granola)

Triana Rohmawati, Kumala Dewi

24-29



PDF



<https://doi.org/10.24252/bio.v7i1.5112>



Views: 608 / PDF Downloads: 140

Toxicity of Bioactive Compound from Endophytic Fungi Isolated from Red Ginger (*Zingiber officinale* var. *rubrum*) Utilizing Brine Shrimp Lethality Assay

Angga Prasetyo, Boy Rahardjo Sidharta, Yustina Sri Hartini, Exsyupransia Mursyanti

30-37



PDF



<https://doi.org/10.24252/bio.v7i1.6000>



Views: 910 / PDF Downloads: 182

Application of Indigenous AMF from ex-coal Mining Soil Combined with Phosphorus Fertilizers to Improved Oil Palm Seedling Growth (*Elaeis guineensis* Jacq.)

Made Deviani Duaja, Elis Kartika, Lizawati Lizawati

38-43



PDF



<https://doi.org/10.24252/bio.v7i1.5990>



Views: 480 / PDF Downloads: 104

The Effect of *Chlorella vulgaris* on Lipid Profile Wistar Strain Rats (*Rattus norvegicus* Berkenhout, 1769) Under Induced Stress

Farah Nadia Karima, Mulyati Sarto

44-53



PDF

 <https://doi.org/10.24252/bio.v7i1.7292>

 Views: 380 / PDF Downloads: 115

Notes on *Cyclanthera brachystachya* (Ser.) Cog.: Morphology, Ecology, and Its New Distribution

Muhammad Efendi, Rugayah Rugayah

54-57

 PDF

 <https://doi.org/10.24252/bio.v7i1.6239>

 Views: 498 / PDF Downloads: 113

Geometric Morphometrics Analysis of Chelae and Carapace of the Freshwater Prawn *Macrobrachium Bate*, 1868

Rich Gemilang Simanjuntak, Rury Eprilurahman

58-66

 PDF

 <https://doi.org/10.24252/bio.v7i1.7803>

 Views: 479 / PDF Downloads: 137

The Potential of Breadfruit Seed and Jackfruit Seed as Alternative Replacement Medium of Potato Dextrose Agar (PDA) with Seedling F0 Mushrooms

Suparti Suparti, Lina Agustina, Putri Agustina, Resti Rahmawati

67-72

 PDF

 <https://doi.org/10.24252/bio.v7i1.6102>

 Views: 1542 / PDF Downloads: 456

Editorial Team
Reviewer
Focus and Scope
Submission
Author Guidelines
Author Fees
Publication Ethics
Screening for Plagiarism

Submit Now

TEMPLATE



CITATION ANALYSIS

Scopus[®]
Citation Analysis

WEB OF SCIENCE[™]
Citation Analysis


TOOLS



INDEXED BY:



HOW TO SUBMIT





- All Rights Reserved -

Platform and Workflow by OJS/PKP Version 3.1.2.4
[Rumah Jurnal UINAM](#) | [About this Publishing System](#)

Toxicity of Bioactive Compound from Endophytic Fungi Isolated from Red Ginger (*Zingiber officinale* var. *rubrum*) Utilizing Brine Shrimp Lethality Assay

ANGGA PRASETYO¹, BOY RAHARDJO SIDHARTA², YUSTINA SRI HARTINI³,
EXSYUPRANSIA MURSYANTI¹

¹Faculty of Biotechnology, Universitas Atma Jaya Yogyakarta
Jl. Babarsari 44, Yogyakarta, Indonesia. 55281

Email: saprasetyo5@gmail.com; mursyantiexsyupranksia@gmail.com

²Bioindustry Laboratory, Faculty of Biotechnology, Universitas Atma Jaya Yogyakarta
Jl. Babarsari 44, Yogyakarta, Indonesia. 55281

Email: brsidharta@gmail.com/brsidharta@uajy.ac.id

³Faculty of Pharmacy, Universitas Sanata Dharma
Jl. Paingan, Maguwoharjo, Sleman, Yogyakarta, Indonesia. 55284
Email: justinahartini@usd.ac.id

Received 10 September 2018; Received in revised form 30 March 2019;

Accepted 19 June 2019; Available online 30 June 2019

ABSTRACT

Red ginger (*Zingiber officinale* var. *rubrum*) has been proven to show anticancer activity. Direct use bioactive compound from red ginger has many obstacles such as large amount of red ginger's rhizome needed, limitation of planting area, and very long time of harvesting. Utilization of endophytic fungi from red ginger's rhizome could be an alternative to the problems. The aims of this study were to determine bioactive compound produced by endophytic fungi and toxicity activity based on LC₅₀. Endophytic fungi were isolated from red ginger and were identified macroscopically and microscopically. The bioactive compounds were extracted using ethanol 96%. Flavonoid test was done qualitatively, bioactive compounds were analyzed by Thin Layer Chromatography (TLC), and the toxicity test was done using Brine Shrimp Lethality Assay (BSLA). The present research found two endophytic fungi isolated from red ginger rhizome. Isolate 1 was similar to *Mucor* sp. and isolate 2 was similar to *Trichoderma* sp. Phytochemical test revealed bioactive compound extracted from the isolates were contained flavonoid. TLC analysis did not detect quercetin from the bioactive compound extracted from the isolates. LC₅₀ values of the bioactive compound from the isolates were 2.300 and 1.747 µg/ml, respectively. The toxicological results suggest that both isolates produce non-toxic compound to *Artemia salina*.

Keywords: brine shrimp lethality assay; endophytic fungi; red ginger; toxicity

INTRODUCTION

Red ginger (*Zingiber officinale* var. *rubrum*) contains bioactive compounds that have capabilities as anticancer, such as flavonoid, gingerol, and shogaol (Danciu *et al.*, 2015; Semwal *et al.*, 2015). Usage of the bioactive compounds directly from ginger faces many obstacles, i.e. large biomass needed, land use limitation, and length of harvesting time (Ginting, 2013). One of the alternatives to solve the problems is to utilize endophytic fungi from red ginger.

Endophytic fungi are a group of microorganism that all or parts of their lives are found in the plant tissues (Kumara *et al.*, 2014). Endophytic fungi produce useful bioactive substances such as antimicrobial, fungicide, antivirus, and antioxidant (Lakshmi & Selvi,

2013). Usage of endophytic fungi as sources of bioactive substances is believed to give benefits, i.e. substances produced are homogenous, fast production, ease to be scaled up, and the amount of productions can be multiplied (Kumara *et al.*, 2012).

Brine Shrimp Lethality Assay (BSLA) is the initial screening method for anticancer substances (Kolbeck & Tintjer, 2016). BSLA is done based on acute toxicity of an extract to *Artemia salina* larvae by LC₅₀ value after 24 hours treatment. If LC₅₀ < 1000 µg/ml, the sample extract is considered to have high toxic effect and is hoped to have cytotoxic effect that becomes a prerequisite for an anticancer substance (Kolbeck & Tintjer, 2016; Bucker *et al.*, 2013).

Kaitu *et al.* (2013) reported that endophytic fungi found from red ginger rhizome was able to inhibit *Escherichia coli* and *Streptococcus pyogenes*. Endophytic fungi isolated from red ginger rhizome were also able to inhibit pathogenic fungi *Fusarium oxysporum* (Ginting, 2013). However, the bioactive substances produced by the endophytic fungi and their toxicity have never been reported.

The objectives of the research are to determine bioactive substances produced by endophyte fungi isolated from red ginger qualitatively, toxicity level of the bioactive substances utilizing BSLA method, and LC₅₀ value of the bioactive substances.

MATERIALS AND METHODS

Red ginger rhizomes were purchased from cultivation field at Rajek Lor village, Mlati subdistrict, Sleman district, Yogyakarta Special Province. Other materials used were as follows eggs of *Artemia salina* from Balai Besar Pengembangan Budidaya Air Jepara, Central of Java.

PDA and PDB media preparations. PDA medium was made by diluting 39 g PDA powder in 1 L aquadest and adding it with chloramphenicol antibiotic 0.5 g/L, stirred and boiled to dilute. Afterward it was sterilized in the autoclave for 15 min at 121°C and 1.5 atm (Sigma Aldrich, 2017). PDB medium was made by slicing 300 g potato, then add with 1000 ml aquadest. The medium was boiled and then filtered. As much as 500 ml of the potato extract was put into a bottle and add with 10 g dextrose and NH₄NO₃ 1%. The media were sterilized with autoclave at 121°C for 15 min (Sigma Aldrich, 2017).

Pure culture of endophyte fungi. Red ginger rhizomes were cleansed with running water. The rhizomes were sterilized by soaking into ethanol 70% for 1 minute and then were rinsed three times with sterile water. The rhizomes were soaked again into hypochlorite solution 0.5% for 5 min and were rinsed for six times with sterile water. Red ginger rhizomes were cut into smaller sizes, i.e. 2 x 2 x 2 cm³. The pieces of rhizome were put on the surface of PDA medium and were incubated at 28°C. Observations were done everyday to determine

the growth of endophytic fungi after isolation (Ginting, 2013; Sadrati *et al.*, 2013). Colonies of endophytic fungi growth were purified in new PDA medium and incubated for 5 days. If there were different colonies growth at the PDA medium, thus it will be re-isolated until pure culture was obtained. The pure culture of endophytic fungi was grown on agar slant to be identified (Noverita *et al.*, 2009).

Endophytic fungi identification.

Identification was done macroscopically by observing the morphology, textures, and colors of the colonies. Microscopic identification was done by observing the types of mycelia, spores, and conidia. Microscopic identification was done using lactophenol cotton blue dyes. The identification was done based on Rodriguez *et al.*, (2009); Ginting *et al.* (2013).

Cultivation and production of secondary metabolites. Endophytic fungi were rejuvenated in PDA slant media and were incubated at 28°C for 5 days. Cultivation of the endophytic fungi were done by transferring the isolates into 100 ml PDB media. PDB medium was incubated for 7 days at 30°C into shaker incubator at 120 rpm (Sadrati *et al.*, 2013; Lakshmi & Selvi, 2013; Li *et al.*, 2015). Fermentation medium was separated from the mycelia by filtration and was rinsed with ethanol 96%. Extraction process was done by adding ethanol 96% into the fermentation medium (1:1). The solvent which contains bioactive substances was evaporated using rotary evaporator and put into the oven to get the viscous extract (Nath *et al.*, 2012).

Qualitative flavonoid test. The test was done by adding 5 ml ethanol to 50 mg extract and was boiled for 5 minutes. The solution was added with 5 drops of concentrated HCl and 1 mg powder. Positive results which determine the existence of flavonoid were shown by the appearance of yellow, pink or purple colors (Ali *et al.*, 2018; Sangi *et al.*, 2008).

Bioactive substance analysis. Bioactive substance extracts were weighed as much as 5 mg, diluted into 5 ml methanol, and filtered. Quercetin as the standard substance was weighed as much as 0.01 gram, diluted into 10 ml methanol. Silica gel plate 60 GF₂₅₄ sizes 9 cm x 5 cm was used. Sample substance and

quercetin were spotted at the distance of 1.5 cm from the lower part of the silica plate. The plate was put into the jar that contains mobile phase composed of toluene:ethyl acetate:methanol (5:3:2). The plate was put at the borderline and was observed by UV lamp at the wavelength 254 nm, and the R_f values were then measured (Males *et al.*, 2013; Preethi *et al.*, 2017; Rao & Ahmed, 2013).

Hatching of *Artemia*. Two g of *Artemia* eggs were hatched in the aquarium filled with 500 ml seawater and equipped with aerator. Seawater temperatures were set at 25-30°C and supported with LED lamps. After 48 hours, *Artemia* eggs will be hatched into larvae, the latter were pipetted and ready to be used in the toxicity test (Kolbeck & Tintjer, 2016; Eimanifar, 2014).

Preliminary toxicity test. Preliminary toxicity test was done to determine the critical concentration ranges to be applied at the determinative toxicity test (Bücker *et al.*, 2013). At the preliminary test, every ten larvae of *A. salina* were put into small bottle with extract solutions at different concentrations, namely 0, 10, 100, 1000 µg/ml. The toxicity tests were

done in five replications. After 24 hours, the percentage of death larvae was counted (Naidu *et al.*, 2014; Hiola & Bahri, 2010).

Determinative toxicity test. BSLA was done by putting every ten *A. salina* larvae aged 48 hours into bottle filled with 5 ml extract solution with the concentration determined from preliminary test and was added with 5 ml seawater. The control treatment was seawater with ten *A. salina* larvae without extract. The bottles were incubated for 24 hours with LED lamps. After 24 hours, the numbers of life larvae were counted (Kolbeck & Tintjer, 2016; Naidu *et al.*, 2014). The criterion of life larvae was larvae that still able to move for few seconds during observations (Rajabi *et al.*, 2015).

Data Analysis. Data were analyzed using probit to determine LC_{50} value. Extracts were considered as toxic if the LC_{50} values are lower than 1000 µg/ml (Kolbeck & Tintjer, 2016).

RESULT AND DISCUSSION

Isolation of Endophytic Fungi. There were two isolates of endophytic fungi isolated from red ginger rhizome (Figure 1).




Figure 1. Pure culture of endophytic fungi isolated from red ginger rhizome: A. Isolate 1 with white coloration as cotton; B. Isolate 2 with black green colour

Macroscopic characteristics of the isolate 1 were white color, smooth as cotton, and colony's back color was yellowish white; while the microscopic characters observed were the types of sporangiophore, sporangium, columella, and sporangiospore (Figure 2). Macroscopic and microscopic characteristics showed that isolate 1 was very similar to *Mucor* sp. and according to Rodriguez *et al.* (2009) the species that has white colony color and can be

changed into greyish brown was identified as *Mucor racemosus*, while colony with back color yellowish white was identified as *Mucor hiemalis*, and species with thick walled and branched sporangiospores was identified as *Mucor plumbeus*. The isolate 1 was also similar to *Rhizopus* sp., but differ in that *Rhizopus* sp. has rhizoid, while *Mucor* sp. does not have rhizoid (Purwantisari & Hastuti, 2009; Rodriguez *et al.*, 2009).




Figure 2. A-C. Morphology of isolate 1 by lactophenol cotton blue dyes and observed under light microscope at 450 x; D. Morphology of one of the species of *Mucor* sp., i.e. *Mucor plumbeus* (Purwantisari & Hastuti, 2009); a. Sporangiophore; b. Sporangium; c. columella; d. round shape sporangiospore

Macroscopic characteristics of isolate 2 (Figure 1B) were black green in color, rough surface, and colony's back color were white, while its microscopic characteristics (Figure 3) showed the existence of conidiophore, round shape conidium, and phyalides or cells that formed conidium. The characters were very similar to *Trichoderma* sp. Rodriguez *et al.*, (2009) stated that *Trichoderma* sp. colony has no color at the beginning but will change to greenish white and turned into dull green; has no color for the colony's back. Change of colony color was due to the age of the colony.

Colony of young *Trichoderma* will have white color, but later it will turn into green and finally into black green (Suparno *et al.*, 2016). Macroscopically it has the same color as *Penicillium* sp., i.e. from greyish green (*Penicillium griseofulvum*) to dark green (*Penicillium purpurogenum*). The difference was seen microscopically i.e. *Penicillium* sp. has metulae that is the apical branch of conidiophore that bring phyalides, while at *Trichoderma* sp. metulae does not exist at all (Rodriguez *et al.*, 2009).




Figure 3. A. Morphology of isolate 2 with lactophenol cotton blue dyes observed under light microscope at 450 x; B. Morphology of conidia from isolate 2 with lactophenol cotton blue dyes at 450 x; C. Morphology of *Trichoderma* sp. at 1000 x (Purwantisari & Hastuti, 2009); a. filial or cells that formed conidia; b. conidiophore; c. semicircle conidia shape

Kaitu *et al.* (2013) found an isolate that was similar to *Mucor* sp. from red ginger rhizome. Ginting (2013) isolated *Mucor* sp. and *Trichoderma* sp., but was not from red ginger

rhizome. The existence of endophytic fungi in plant tissues were influenced by climate and geographical position as well as culture medium used; since culture medium gave

similar conditions to natural habitat of endophytic fungi. The mechanism adaptation of endophytic fungi to physiological and microecological conditions of the plant's hosts were well noted (Costa *et al.*, 2012; Fernandes *et al.*, 2015; Noverita *et al.*, 2009).

Qualitative Flavonoid Test. The result of flavonoid qualitative test showed that both isolates were able to produce flavonoid (Table 1). Yellow color appeared due to reduction

reactions between Mg, HCl, and the samples (Danciu *et al.*, 2015). Endophytic fungi were able to produce flavonoid through shikimate pathway that is started with condensation process between phosphoenolpyruvate (PEP) and erythrose-4-phosphate (E-4-P) (Tohge *et al.*, 2013). Shikimate pathway produces chorismate that will become the precursor for phenylalanine and the latter will become the precursor for flavonoid (Tohge *et al.*, 2013).

Table 1. Flavonoid qualitative test

Isolate	Initial Colour	End Colour	Test Result
Isolate 1	White	Yellow	+
Isolate 2	White	Yellow	+

+ = contain flavonoid

Biosynthesis of flavonoid was also done through malonic acid pathway that produced malonyl CoA. Three molecules of malonyl CoA will undergo condensation with one molecule of 4-co

chalcone synthase to produce chalcone. Chalcone was isomerized into naringenin that will undergo several reactions to produce flavanone, flavonol, flavanol, and anthocyanin




Figure 4. TLC results: A. without UV light; B. UV light with wavelength 254 nm; C. UV light with wavelength 365 nm. a. Standard quercetin with R_f value of 0.67; b. Extract of isolate 1; c. Extract of isolate 2

TLC analysis showed R_f value for quercetin standard at 0.67 (Figure 4). Extracts from isolate 1 and 2 were not able to show clear spots at R_f value 0.67. Extracts of bioactive substance from isolate 1 and 2 did not produce quercetin, which was one of the members of flavonoid. Quercetin was not detected from the extracts of both isolates due to very low concentrations. Preethi *et al.*, (2017) reported that one of the disadvantages of TLC i.e. it cannot detect low concentration of substances. Other possibility was that the endophytic fungi did not produce quercetin due to ecological factors such as rain falls, climate, and nutrient

contents in the soil affected secondary metabolites production (Liu *et al.*, 2015).

Figure 4 showed sample substances that were not fully separated by the mobile phase used. Imperfect separation was due to interaction of substances involved, for instance polar substance can form hydrogen bonds; substances bind to immobile phase, and mobile phase did not work optimally. Thus, advance elution needs to be done to utilize mobile phase with different polarity levels (Santosa & Haresmita, 2015).

Toxicity Test. Preliminary toxicity test was done and the result of the death percentage of *A. salina* larvae at concentration 1000 $\mu\text{g/ml}$

was as high as 30%. The concentration was then needed to be increased in order to reach higher percentage value. For determinative toxicity test the concentrations used were varied,

namely 1000, 2000, 3000, 4000, and 5000 $\mu\text{g/ml}$ (Table 2).

Table 2. Toxicity of bioactive substances from isolate 1 and isolate 2

Isolate	Average Death of <i>A. salina</i> larvae (%)					Control
	Concentration ($\mu\text{g/ml}$)					
	1000	2000	3000	4000	5000	
Isolate 1	28	42	52	68	80	4
Isolate 2	38	52	60	70	78	2

LC₅₀ of the bioactive compound from the isolates were 2,300 and 1,747 $\mu\text{g/ml}$, respectively. Bioactive substances from isolate 1 and 2 yielded LC₅₀ value > 1000 $\mu\text{g/ml}$ and categorized as not toxic (Kolbeck & Tintjer, 2016). LC₅₀ value less than 1000 ppm means that the bioactive substance tested has the potential effect as anticancer, but if the value is higher than 1000 ppm means the bioactive substance has no potency as anticancer (Pasilala *et al.*, 2016). Therefore, bioactive substances from isolate 1 and 2 did not show any potency as anticancer.

Flavonoid can kill *A. salina* larvae by disrupting larvae cell walls, inhibiting enzyme activities, disturbing the function of larvae cell membrane, and inhibiting protein synthesis due to the activation of α -subunit of eukaryotic initiation factor 2 (eIF2 α). Thus phosphorylation was occurred at eIF2 α kinase that caused failure of initiation of mRNA translation (Ginting *et al.*, 2013; Roy, 2016). Death of *A. salina* larvae could also cause by flavonoid that gave up its OH⁻ group to bind to cell membrane integral protein of *A. salina* larvae. The OH⁻ group that binds to cell membrane integral protein will inhibit active transport of Na⁺ and K⁺, thus influx of Na⁺ to the cells becomes uncontrolled and caused the cell membrane rupture (Chen *et al.*, 2014).

CONCLUSION

The present research concluded that Isolate 1 and 2 produce bioactive substances such as flavonoid, but quercetin was not detected in the extracts from the two isolates, not toxic based on BSLA, with LC₅₀ value at 2.300 $\mu\text{g/ml}$ (isolate 1), while isolate 2 at 1.747 $\mu\text{g/ml}$.

ACKNOWLEDGMENTS

The researchers are indebted to Faculty of Biotechnology Universitas Atma Jaya Yogyakarta for permitting the research using laboratory facilities. Thank you to Research and Community Service Center Universitas Atma Jaya Yogyakarta for partially funding the research.

REFERENCES

- Ali AMA, El-Nour MEM, Yagi SM. 2018. Total phenolic and flavonoid contents and antioxidant activity of ginger (*Zingiber officinale* Rosc.) rhizome, callus and callus treated with some elicitors. *Journal of Genetic Engineering and Biotechnology*. vol 16(2): 677-682. doi: <https://doi.org/10.1016/j.jgeb.2018.03.003>
- Bücker A, Falcão-Bücker NC, Nunez CV, Pinheiro CCS, Tadei WP. 2013. Evaluation of larvicidal activity and brine shrimp toxicity of rhizome extracts of *Zingiber zerumbet* (L.) Smith. *Revista da Sociedade Brasileira de Medicina Tropical*. vol 46(3): 377-380. doi: <http://dx.doi.org/10.1590/0037-8682-1309-2013>.
- Costa IPMW, Maia LC, Cavalcanti MA. 2012. Diversity of leaf endophytic fungi in mangrove plants of northeast Brazil. *Brazilian Journal of Microbiology*. vol 43(3): 1165-1173. doi: 10.1590/S1517-838220120003000044.
- Danciu C, Vlaia L, Fetea F, Hancianu M, Coricovac DE, Ciurlea SA, Şoica CM, Marincu I, Vlaia V, Dehelean CA, Trandafirescu C. 2015. Evaluation of phenolic profile, antioxidant and

- anticancer potential of two main representants of Zingiberaceae family against B164A5 murine melanoma cells. *Biological Research*. vol 48(1): 1-9. doi: 10.1186/0717-6287-48-1.
- Eimanifar A. 2014. Molecular phylogeography and population genetics of the brine shrimp *Artemia* (Crustacea, Branchiopoda, Anostraca). [Dissertation]. Heidelberg: Faculty of Natural Sciences and Mathematics, University of Heidelberg, Germany
- Fernandes EG, Pereira OL, da Silva CC, Bento CBP, de Queiroz MV. 2015. Diversity of endophytic fungi in *Glycine max*. *Microbiological Research*. vol 181: 84–92. <http://dx.doi.org/10.1016/j.micres.2015.05.010>.
- Ginting RCB. 2013. Keragaman cendawan endofit yang diisolasi dari berbagai spesies tanaman obat potensial. [Tesis]. Bogor: Program Studi Mikrobiologi IPB.
- Ginting RCB, Sukarno N, Widyastuti U, Darusman LK, Kanaya S. 2013. Diversity of endophytic fungi from red ginger (*Zingiber officinale* Rosc.) plant and their inhibitory effect to *Fusarium oxysporum* plant pathogenic fungi. *HAYATI Journal of Biosciences*. vol 20(3): 127-137. doi: 10.4308/hjb.20.3.127.
- Hiola SF, and Bahri A. 2010. Uji toksisitas ekstrak daun ketepeng cina (*Cassia alata* L.) pada keong mas (*Pomacea canaliculata* L.). *Bionature*. vol 11(2): 115-119.
- Kaitu RAM, Sidharta BR, Atmodjo K. 2013. Aktivitas Antibakteri Fungi Endofit Jahe Merah (*Zingiber officinale* var. rubrum) Terhadap *Escherichia coli* dan *Streptococcus pyogenes*. *Jurnal Biologi*. p 1-15.
- Kolbeck MK, and Tintjer TE. 2016. The use of a brine shrimp assay to detect bioactivity in the endophyte infected grass, *Agrostis hyemalis*. *Journal of the Pennsylvania Academy of Science*. vol 90(1): 13-20.
- Kumara PM, Zuehlke S, Priti V, Ramesha BT, Shweta S, Ravikanth G, Vasudeva R, Santhoshkumar TR, Spiteller M, Shaanker RU. 2012. *Fusarium proliferatum*, an endophytic fungus from *Dysoxylum binectariferum* Hook.f, produces rohitukine, a chromane alkaloid possessing anti-cancer activity. *Antonie van Leeuwenhoek*. vol 101: 323–329. doi: 10.1007/s10482-011-9638-2.
- Kumara PM, Shweta S, Vasanthakumari MM, Sachin N, Manjunatha BL, Jadhav SS, Ravikanth G, Ganeshaiyah KN, Shaanker RU. 2014. Endophytes and plant secondary metabolite synthesis: molecular and evolutionary perspective. In: V.C. Verma and A.C. Gange (eds.). *Advances in Endophytic Research*. doi: 10.1007/978-81-322-1575-2_9.
- Lakshmi PJ, and Selvi KV. 2013. Anticancer potentials of secondary metabolites from endophytes of *Barringtonia acutangula* and its molecular characterization. *International Journal of Current Microbiology and Applied Sciences*. vol 2(2): 44-45.
- Li X, Guo Z, Deng Z, Yang J, Zou K. 2015. A new a-pyrone derivative from endophytic fungus *Pestalotiopsis microspora*. *Rec. Nat. Prod*. vol 9(4): 503-508.
- Liu W, Liu J, Yin D, Zhao X. 2015. Influence of ecological factors on the production of active substances in the anticancer plant *Sinopodophyllum hexandrum* (Royle). *PloS One*. vol 10(4): 1-22. <https://doi.org/10.1371/journal.pone.0122981>.
- Males C, Saric D, Bojic M. 2013. Quantitative determination of flavonoids and chlorogenic acid in the leaves of *Arbutus unedo* L. using thin layer chromatography. *Journal of Analytical Methods in Chemistry*. vol 2013: 1-4. doi: <http://dx.doi.org/10.1155/2013/385473>.
- Naidu JR, Ismail R, Sasidharan S. 2014. Acute Oral Toxicity and Brine Shrimp Lethality of Methanol Extract of *Mentha spicata* L (Lamiaceae). *Tropical Journal of Pharmaceutical Research*. vol 13(1): 101-107. doi: <http://dx.doi.org/10.4314/tjpr.v13i1.15>.
- Nath A, Raghunatha P, Joshi SR. 2012. Diversity and biological activities of endophytic fungi of *Embllica officinalis*, an

- ethnomedicinal plant of India. *Mycobiology*. vol 40(1): 8-13. doi: 10.5941/MYCO.2012.40.1.008.
- Noverita, Fitria D, Sinaga E. 2009. Isolasi dan uji aktivitas antibakteri jamur endofit dari daun dan rimpang *Zingiber ottensii* Val. *Jurnal Farmasi Indonesia*. vol 4(4): 171-176.
- Pasilala FB, Daniel, Saleh C. 2016. Uji toksisitas (*Brine Shrimp Lethality Test*) dan aktivitas antioksidan dari daun sintrong (*Crassocephalum crepidioides*) dengan metode *2,2-diphenyl-1-picrylhidrazil* (DPPH). *Jurnal Kimia Mulawarman*. vol 14(1): 13-18.
- Preethi J, Harita B, Rajesh T. 2017. Review on Thin Layer Chromatography. *Journal of Formulation Science and Bioavailability*. vol 1(2017): 107.
- Purwantisari S, and Hastuti RB. 2009. Isolasi dan identifikasi jamur indigenous rhizosfer tanaman kentang dari lahan pertanian kentang organik di Desa Pakis, Magelang. *Bioma*. vol 11(2): 45-53. doi: <https://doi.org/10.14710/bioma.11.2.45-53>.
- Rajabi S, Ramazani A, Hamidi M, Naji T. 2015. *Artemia salina* as a model organism in toxicity assessment of nanoparticles. *DARU Journal of Pharmaceutical Sciences*. vol 23(1): 20. doi: 10.1186/s40199-015-0105-x.
- Rao AS, and Ahmed MF. 2013. Simultaneous estimation of quercetin and rutin in ethanolic extract of *Melia azedarach*. Linn leaves by HPTLC method. *Asian Journal of Biomedical and Pharmaceutical Sciences*. vol 3(21): 56-69.
- Rodriguez RJ, White Jr JR, Arnold AE, Redman RS. 2009. Fungal endophytes: diversity and functional roles. *New Phytologist*. vol 182: 314-330. doi: 10.1111/j.1469-8137.2009.02773.x.
- Roy S. 2016. Polyphenol Content and Differential Expression of Flavonoid Biosynthetic Pathway Genes of *Fragaria* spp with white fruit. [Dissertations]. Lexington: Department of Plant and Soil Sciences. doi: <http://dx.doi.org/10.13023/ETD.2016.033>.
- Sadrati N, Daoud H, Zerroug A, Dahamna S, Bouharati S. 2013. Screening of antimicrobial and antioxidant secondary metabolites from endophytic fungi isolated from wheat (*Triticum durum*). *Journal of Plant Protection Research*. vol 53(2): 128-136.
- Sangi M, Runtuwene MRJ, Simbala HEI, Makang VMA. 2008. Analisis fitokimia tumbuhan obat di kabupaten Minahasa Utara. *Chem. Prog*. vol 1(1): 47-53.
- Santosa D, and Haresmita PP. 2015. Antioxidant activity determination *Garcinia dulcis* (Roxb.) Kurz, *Blumea mollis* (D. Don) Merr, *Siegesbeckia orientalis* L. and *Salvia riparia* H. B. K. which collected from Taman Nasional Gunung Merapi using DPPH (2,2-diphenyl-1-pikril-hidrazil) and thin layer chromatography. *Traditional Medicine Journal*. vol 20(1): 28-36. doi: <https://doi.org/10.22146/tradmedj.7746>.
- Semwal RB, Semwal DK, Combrinck S, Viljoen AM. 2015. Gingerols and shogaols: important nutraceutical principles from ginger. *Phytochemistry*. vol 117: 554-568. doi: 10.1016/j.phytochem.2015.07.012.
- Sigma Aldrich. 2017. Fungal and bacterial detection media. <http://www.sigmaaldrich.com>.
- Suparno, Sukoso, Hakim L, Aidawati N. 2016. *Trichoderma* spp. as agent of biological control in local rice diseases in tidal swamp lands in South Kalimantan, Indonesia. *Journal of Agriculture and Veterinary Sciences*. vol 9(1): 1-6. doi: 10.9790/2380-09120106.
- Tohge T, Watanabe M, Hoefgen R, Fernie AR. 2013. Shikimate and phenylalanine biosynthesis in the green lineage. *Frontiers in Plant Science*. vol 4(62): 1-13. doi: 10.3389/fpls.2013.00062