

Isocratic high-performance liquid chromatography (HPLC) for simultaneous quantification of curcumin and piperine in a microparticle formulation containing *Curcuma longa* and *Piper nigrum*

by Yustina Sri Hartini

Submission date: 15-Nov-2021 08:42PM (UTC+0700)

Submission ID: 1703409148

File name: 7_2021_e06541_Isocratic_high_liquid_Dewi_Setyaningish_et_al.pdf (976.27K)

Word count: 6909

Character count: 35160



Contents lists available at ScienceDirect

Heliyon

journal homepage: www.cell.com/heliyon



Research article

Isocratic high-performance liquid chromatography (HPLC) for simultaneous quantification of curcumin and piperine in a microparticle formulation containing *Curcuma longa* and *Piper nigrum*



Dewi Setyaningsih^{a,*}, Yosua Agung Santoso^a, Yustina Sri Hartini^a, Yosi Bayu Murti^b, Wouter L.J. Hinrichs^c, Christine Patramurti^a

^a Faculty of Pharmacy, Sanata Dharma University, Indonesia

^b Faculty of Pharmacy, Universitas Gadjah Mada, Indonesia

^c Department of Pharmaceutical Technology and Bio-pharmacy, University of Groningen, the Netherlands

ARTICLE INFO

Keywords:

Curcumin
Piperine
Extract
Solid dispersion
Reverse-phase
Validation
Optimization

ABSTRACT

Poor bioavailability has been reported as a major challenge in the development of curcumin as a pharmaceutical agent. However, co-administration of curcumin with piperine has been shown to improve curcumin bioavailability. Therefore, to assure product control quality, an analytical method needs to be developed for the determination of curcumin and piperine content in a dosage form formulation. The objective of this study was to develop a simple isocratic reversed-phase HPLC (RP-HPLC) method to simultaneously quantify curcumin and piperine content in a solid dispersion based microparticle formulation containing *Curcuma longa* and *Piper nigrum* extracts. The method was validated according to the International Council for Harmonization (ICH) guideline. Chromatographic separation of three curcuminoids and piperine could be achieved using acetonitrile-methanol-water of 65:5:35 %, at a flow rate of 1 mL/min and a wavelength of 353 nm for detection. Resolution (Rs) of 3.57 and 1.68 for piperine and curcumin, respectively, a theoretical plate number (n) > 8000 and a tailing factor (T) < 1.5 indicate a satisfactory separation of the compounds. The calibration curve was linear from 1.25–15 µg/mL and 2.5–30 µg/mL for piperine and curcumin, respectively, with the correlation coefficient of >0.999. The intra-day/inter-day accuracy and precision demonstrated a recovery of 99.54–101.50%/99.38–99.89% and 100.78–102.51%/101.15–102.47% with a Relative Standard Deviation (RSD) of 0.53–0.95%/0.13–1.44 % and 0.28–1.62%/0.46–1.14% for piperine/curcumin. The limit of detection (LOD) were 0.27 and 0.42 µg/mL, for piperine and curcumin, which reveals an adequate sensitivity. A solid dispersion based microparticle formulation containing *C. longa* and *P. nigrum* extracts confirmed the validity of the developed method as a recovery of 91.14% and 99.14% for piperine and curcumin, respectively. In conclusion, all the validated parameters confirm the precision, accuracy, and reliability of the method for the simultaneous analysis of curcumin and piperine within a microparticle formulation containing *C. longa* and *P. nigrum* extracts.

1. Introduction

Curcumin, a yellow polyphenolic compound isolated from turmeric (*Curcuma longa*) rhizome, has been claimed as a therapeutic agent for numerous diseases due to its anti-oxidant and anti-inflammatory properties [1, 2, 3]. However, despite extensive *in-vitro* and *in-vivo* studies, a clear therapeutic effect of curcumin in clinical studies has not been found yet [4]. These disappointing results can be ascribed to the poor bioavailability of the compound after oral intake. There are two reasons for the poor absorption of curcumin; 1) It exhibits a very low aqueous

solubility of only 11 ng/mL [4, 5], and 2) It is rapidly metabolized [1]. Therefore, a combination of two strategies addressing poor solubility and rapid metabolism is required.

An often-applied strategy to improve the dissolution behavior of a poorly water-soluble drug is formulating it as a solid dispersion [6]. Furthermore, piperine (Figure 1d), an alkaloid compound of black pepper (*P. nigrum*) extract has been identified as a bio-enhancer of various drugs, by stimulating gut amino acid transporters and inhibiting drug-metabolizing enzymes to improve drug absorption, bioavailability, and bio-efficacy [7]. Both strategies are thought to be useful to be applied

* Corresponding author.

E-mail address: dewi@usd.ac.id (D. Setyaningsih).

<https://doi.org/10.1016/j.heliyon.2021.e06541>

Received 7 August 2020; Received in revised form 14 November 2020; Accepted 12 March 2021

2405-8440/© 2021 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Figure 1. Molecular structure of (a) curcumin; (b) demethoxycurcumin; (c) bis-demethoxycurcumin; (d) piperine.

to curcumin. Hence, we propose a formulation based on solid dispersion technology containing *C. longa* and *P. nigrum* extracts to improve the bioavailability of curcumin. In this study, we will focus on solid dispersion microparticles prepared by spray drying using polyvinylpyrrolidone K30 (PVP K30) as a carrier.

During pre-formulation, the quality control of the product needs to be addressed. Determining compounds in the final product is a part of quality assurance in production processes. Therefore, the amount of curcumin and piperine containing in the microparticle needs to be determined accurately.

Several analytical methods for the quantification of curcumin and piperine have been developed, however, the majority of these methods concern the quantification of curcumin and piperine individually [8, 9, 10, 11]. A few methods have been developed to simultaneously determine curcumin and piperine. A spectroscopic method combining with the application of Vierordt's equation was successfully applied to allow the determination of curcumin and piperine as a mixture of *C. longa* and *P. nigrum* extracts in dissolution medium [12], however, the method is not selective.

HPLC with UV-Vis detection is one of the most common techniques used in the analysis and quality control of several formulations containing curcumin or piperine and in biological samples. Moorthi *et al.* validated an isocratic HPLC method to allow curcumin and piperine quantification in a Eudragit based nanosuspension containing pure curcumin and piperine as a binary mixture [13]. Furthermore, Sethi *et al.* reported a selective and specific HPLC method for simultaneous estimation of curcumin and piperine in human plasma [14]. Also, a sensitive LC/MS method was developed to enable simultaneous quantification of curcumin and piperine for pharmacokinetic evaluation [15]. Although the HPLC methods have been reported to be useful to simultaneously analyze curcumin and piperine in various samples, the developed methods were applied to determine curcumin and piperine as pure compounds in the samples.

To our best knowledge, no HPLC analytical method has been reported for the simultaneous analysis of curcumin and piperine in extracts. It should be emphasized that commercially available curcumin originating from turmeric rhizome is usually composed of a mixture of naturally occurring curcuminoids, namely curcumin, demethoxycurcumin, and bis-demethoxycurcumin (Figure 1a, b, c). This mixture of compounds could easily interfere with chromatographic separation if not properly resolved. Therefore, the study aimed to develop a validated HPLC method to simultaneously quantify curcumin and piperine in solid dispersion-based microparticle formulation containing *C. longa* and *P. nigrum* extracts.

2. Materials and methods

2.1. Materials

Bis-demethoxycurcumin, demethoxycurcumin, curcumin, and piperine reference standards with a purity of >98% were purchased from

Sigma-Aldrich (St. Louis, USA). *C. longa* extract of high curcuminoids content (97.56%) was a kind gift from PT Phytochemindo Reksa, Bogor, Indonesia. *P. nigrum* was extracted and purified as described by Saha *et al.* [16]. Polyvinylpyrrolidone K30 (PVP K30) was kindly given by PT Konimex, Solo, Central Java, Indonesia. Ethanol, HPLC grade acetonitrile, HPLC grade methanol, and analytical grade phosphoric acid were purchased from Merck (Darmstadt, Germany). A PTFE filter with a pore size of 0.22 μ m was obtained from Whatman. Milli-Q water was prepared in the laboratory.

2.2. Preparation of solid dispersion based microparticle formulation and characterization

A PVP-30 based solid dispersion containing 30 w/w % *C. longa* extract and 10 w/w % *P. nigrum* extract was prepared by spray drying. In brief, 1800 mg *C. longa* extract and 600 mg *P. nigrum* extract were dissolved in 400 mL ethanol and 3600 mg of PVP K30 was dissolved in 50 mL ethanol. The two solutions were mixed for approximately 1 min using a magnetic stirrer.

The resulting solution was fed into a BUCHI B-290 micro spray dryer equipped with a B-295 dehumidifier via a two-way channel with a nozzle diameter of 0.7 mm. Drying was conducted at an inlet temperature of 105 °C, aspiration of 100%, feeding rate of 20%, and an atomization pressure of 500 L/h. The outlet temperature was observed at 60–65 °C. Micrographs of the microparticles were taken using a scanning electron microscope (Hitachi TM 3000, Japan).

2.3. Method development

2.3.1. Finding the detection wavelength

Individual methanolic solutions of curcumin (15 μ g/mL) and piperine (5 μ g/mL) were scanned at the range of 315–450 nm using a Vis spectrophotometer (Shimadzu, Japan). The detection wavelength for the simultaneous analysis of curcumin and piperine was selected based on the crossing point of the curcumin and piperine spectra of the spectral overlay.

2.3.2. HPLC instrument and conditions

HPLC analysis was carried out using a Shimadzu LC 2010HT HPLC system (Shimadzu, Kyoto, Japan) equipped with a serial dual plunger pump, an autosampler, and an SPD-20A/20AV series UV-Vis detectors. Solution software was used for peak integration. Chromatographic separation was achieved on a C18 column (250 \times 4.6 mm, Eurospher 27 with 5 μ m) equipped with a pre-column (Knauer, Berlin, Germany). The mobile phase was filtered through a 0.22 μ m filter and degassed by ultrasonication before use. The injection volume was 20 μ L. The column temperature was kept at 33 °C during chromatographic operation.

2.3.3. Selection of mobile phase

The aim was to find a mobile phase composition, which provides an acceptable peak separation of multiple components i.e. curcuminoids

(bis-demethoxycurcumin, demethoxycurcumin, curcumin) and piperine with a short running time of less than 10 min. A sample consisting of a mixture of 10 µg/mL *C. longa* and 5 µg/mL *P. nigrum* extracts in methanolic solution was used to evaluate mobile phases of different compositions.

Peak identification was carried out by running standard solutions at accordingly used mobile phase compositions. The mobile phase compositions were evaluated on the chromatographic separation parameters i.e. tailing factor (T), peak resolution (Rs), and theoretical plate number (N). The acceptance criteria are $T < 2$, $R_s > 1.5$ and $N > 2000$ [17].

2.3.4. Preparation of stock and calibration solutions

Stocks of standard solution of curcumin and piperine at a final concentration of 1 mg/mL were prepared by dissolving an appropriate amount of curcumin and piperine in methanol as an individual solution. All solutions were prepared in light-protected vials to ensure the stability of curcumin and piperine [18].

2.4. Method validation

The developed method was validated as per Q2R1 ICH guidelines [19] including selectivity, system suitability, accuracy, precision, limit of detection (LOD), limit of quantification (LOQ), linearity, and robustness.

2.4.1. System suitability

Six replicate injections of the sample containing a mixture of curcumin and piperine at a concentration of 10 and 5 µg/mL in the methanolic solution were simultaneously analysed. Retention time (Tr), peak area (AUC), Rs, T, and N were determined. Relative standard deviation (RSD) values of these parameters were calculated to evaluate the system suitability of the developed method. The suitability test was accepted when the RSD values of these parameters were less than 2%.

2.4.2. Specificity and selectivity

Specificity refers to the ability of a method to distinguish between the compound(s) of interest and impurity present in the sample matrix [20]. For the specificity study, the chromatogram of blank samples originated from the formulation matrix (no drugs) was compared to the chromatogram of microparticle formulation containing *C. longa* and *P. nigrum* extracts. Selectivity is the ability of the method to clearly show separation between the different peaks. A value of $R_s > 1.5$ is required to confirm the method to be sufficiently selective [20].

2.4.3. Preparation of calibration solutions of curcumin and piperine and linearity

Calibration solutions were prepared using solutions of curcumin and piperine in methanol at different concentrations. Stock solutions of curcumin and piperine were diluted with methanol and mixed to obtain samples with curcumin concentrations of 0.05–45 µg/mL and piperine concentrations of 0.025–20 µg/mL for. Linearity between concentration and peak area was tested using the regression line as provided by the least square analyses. The correlation coefficient (r) was used to judge linearity [25]. The concentration range with an $r > 0.99$ was considered linear. The limit of detection (LOD) and the limit of quantification (LOQ) were determined using the standard deviation approach [20]. To do so, serially diluted concentrations of the calibration solutions were prepared early targeted LOD or LOQ and analyzed using least square analysis. The standard deviation (SD) of the y-intercept and slope were determined with which the LOD and LOQ were calculated based on Eq. (1) and Eq. (2) below [22].

$$LOD = 3.3 \times SD / \text{slope} \quad (1)$$

$$LOQ = 10 \times SD / \text{slope} \quad (2)$$

2.4.4. Accuracy and precision

Accuracy refers to the closeness of the obtained value with that true value of the corresponding sample concentrations, while precision denotes the closeness agreement between independent test results obtained under the developed method. In this study, accuracy and precision were carried out using the assay sample spiking method. The assay sample was the microparticle formulation as described in section 2.2; dissolved in methanol. The methanolic sample solutions at the concentration of 9.9 µg/mL (sample weight/volume) were spiked with curcumin and piperine stock solutions to result in a mixture of final addition concentrations of 2.50; 15.00; 30.00 and 1.25; 7.50; 15.00 µg/mL for curcumin and piperine.

Accuracy was evaluated based on the recovery percentage, and precision was studied using repeatability assay based on the RSD values of three different sample concentrations. Accuracy and precision were conducted on one-day analysis (intra-day) and three consecutive days (inter-day) of freshly prepared samples. The analysis was done in triplicate for each concentration level. The degree of accuracy and precision were judged according to the Association of Official Analytical Chemists requirement [21].

2.4.5. Robustness

The robustness of a developed analytical method refers to its ability to remain unaffected by deliberate changes of the chromatographic conditions which indicates its reliability during normal usage. Peaks retention time and AUC values of curcumin and piperine resulted from the changed parameters of mobile phase composition, flow rates, and detection wavelengths were analyzed for the RSD values. The RSD value of less than 2% is an indication of a robust method.

2.4.6. Assay of curcumin and piperine in the microparticle formulation

A 200 mg of the microparticle formulation was accurately weighed and dissolved in 50 mL of methanol in a volumetric flask. The solution was subjected to ultrasonication for 5 min. The solution was diluted 100 times with methanol to achieve a final concentration of 40 µg/mL and then analyzed by HPLC. The content of curcumin and piperine was calculated based on the calibration curve constructed as described in section 2.4.3.

2.5. Data analysis

A Least Square Analysis supported by Excel Microsoft INC USA was used to analyze the linearity of the calibration curve. Mean and standard deviation was calculated using Excel Software (Microsoft Inc, USA).

3. Results and discussion

3.1. Preparation of solid dispersion based microparticle formulation and characterization

Microparticles of PVP-30 based solid dispersion containing curcuminoids of *C. longa* and piperine of *P. nigrum* extracts could be successfully prepared by spray drying as can be seen in the scanning electron microscopic images (see example in Figure 2).

3.2. Finding the detection wavelength

The spectral scanning of a single methanolic solution of curcumin and piperine is shown in Figure 3. As shown in Figure 3, the methanolic solution of curcumin (15 µg/mL) absorbed wavelength at the maximum value of 425 nm, while piperine dissolved in methanol at the concentration of 5 µg/mL showed a maximum absorption at the wavelength of 342 nm. Both compounds exhibited an overlapping absorption area at around 315–360 nm, where curcumin and piperine can be simultaneously detected.




Figure 2. Scanning electron microscopic image of the spray dried PVP K30 based solid dispersion containing curcuminoids and piperine.




Figure 3. The overlay spectra of curcumin (15 µg/ml) and piperine (5 µg/ml) in methanolic solution.

As can be seen in Figure 3, the two spectra cross each other at 353 nm. Therefore, this wavelength was used for the simultaneous detection of curcumin and piperine in this study.

3.3. Selection of mobile phase

3.3.1. The use of methanol and acetonitrile

Simultaneous analysis of a mixture contains weak acids and weak basics drugs on an RP HPLC is always challenging [23]. The target analytes in this study are curcumin (weak acid) and piperine (weak basic) which are present as co-mixture with other curcuminoids compounds (*bis*-demethoxycurcumin, demethoxycurcumin) as the active components in the microparticle formulation. The combinations methanol-water and acetonitrile-water are the most commonly used solvent mixtures in RP HPLC analysis. However, curcuminoids are poorly resolved in methanol-water mixtures at a volume ratio of 50:50 and 60:40 with calculated polarity index of 7.55 and 7.06. Methanol was therefore replaced by acetonitrile. Although with an acetonitrile-water volume ratio of 60:40 (polarity index of 7.64) a sharp peak was obtained, all three curcuminoids were co-eluted at a retention time of 12.54 min.

3.3.2. The effect of 0.1% phosphoric acid as pH modifier

A study by Espinosa et al. revealed that the retention time of acidic and basic compounds is strongly dependent on the pH of the mobile phase [24]. The addition of acid into the mobile phase can improve peak separation and reduce peak tailing by modification of interaction between the target analyte and the stationary phase of the HPLC column [25, 26].

Therefore, a mobile phase composition as reported by Sethi et al., which was used to simultaneously analyze curcumin and piperine in the plasma sample [14], with slight modifications was evaluated. The mobile phase consisted of acetonitrile-methanol-water-phosphoric acid 0.1% at a volume ratio of 20:30:47.5:2.5 (pH 4.0) and was delivered at 1 mL/min in an isocratic mode. However, under these conditions, the three curcuminoids compounds were co-eluted at a retention time of 11.80 min, whereas no piperine peak appeared within a running time of 15 min, while a maximum running time of 10 min was aimed for. To evaluate the behavior of the weakly basic compound (piperine) at a lower pH, the composition of the mobile phase was adjusted to acetonitrile-methanol-phosphoric acid 0.1% at a volume ratio of 20:30:50 (pH 3.0) and a calculated polarity index of 7.79. However, a peak in the chromatogram that can be assigned to piperine remained absent during a run time of 15 min. Remarkably, when the mobile phase polarity was slightly modified by replacing the methanol with acetonitrile to result in a calculated polarity index of 8.00 of the mobile phase compositions of acetonitrile-water-phosphoric acid 0.1% at a volume of 50:47.5:2.5 (pH 4.0), piperine was eluted already at 0.516 min.

LoBrutto et al. reported the effect of pH on the retention time of small basic compounds, aniline, pyridine on an isocratic HPLC mode. At the pH range of 1.3–8.6 of the mobile phase with a composition of acetonitrile-water at a volume ratio of 10:90, it was reported that as the pH of the mobile phase decreased, the retention time of the targeted compounds increased due to counterion interactions resulted from acidic mobile

Table 1. Mobile phase compositions and the observed peak characteristics.

Code	ACN	MeOH	water	Curcuminoids									Piperine		
				BDMC			DMC			C			Tr	Rs	T
				Tr	Rs	T	Tr	Rs	T	Tr	Rs	T			
1	30	25	45	-	-	-	6.30	4.66	0	6.63	0.76	0	-	-	-
2	40	30	30	8.09	1.55	0	8.25	1.08	0	8.99	1.14	0	10.72	5.34	1.17
3	30	50	20	4.26	5.61	1.11	4.86	0	0	5.35	0	0.91	8.77	3.66	1.11
4	60	10	30	5.03	6.11	1.06	5.38	1.56	1.20	5.78	1.64	1.18	7.84	3.22	1.08
5	65	5	30	4.82	4.74	0.95	5.18	1.48	1.12	5.58	1.68	1.16	7.91	3.57	1.06

Note:

1. ACN = acetonitrile; MeOH = methanol.
2. BDMC = *bis*-demethoxycurcumin; DMC = demethoxycurcumin; C = curcumin.
4. Tr = retention time; Rs = resolution; T = tailing factor.
5. - = not detected.




Figure 4. Chromatographic separation of bis-demethoxycurcumin (BDMC), demethoxycurcumin (DMC), curcumin (C) and piperine (P) after injection of 20 μ L of the sample containing 10 μ g/mL *C. longa* and 5 μ g/mL *P. nigrum* extracts. Separation was carried out using a Knauer C18 column (250 \times 4.6 mm, Eurospher 100 μ m) equipped with a pre-column and an isocratic mobile phase composed of an acetonitrile–anol–water mixture with a volume ratio of 65:5:30 at flow rate of 1 mL/min and a detection wavelength of 353 nm.

phase modifier [27]. Guan and Palmer studied the effect of a chaotropic agent trifluoroacetic acid on the retention of four weakly basic derivatives of triazole with different pKa values [28]. Using mixtures of water and acetonitrile as the mobile phase in the gradient mode, it was found that the lower the degree of protonation, the shorter the retention time. As phosphoric acid is also categorized as a chaotropic agent [11], it is presumed that the absence of piperine during 15 min running in this study using the mobile phase composition of acetonitrile–methanol–water–phosphoric acid 0.1% at a volume ratio of 20:30:47.5:2.5 (pH 4.0) and acetonitrile–methanol–phosphoric acid 0.1% at 20:30:50 (pH 3.0) might be attributed to the effect of chaotropic effect resulting partially protonated piperine, making piperine molecules interact with the stationary phase. However, when methanol was replaced by acetonitrile, a solvent with higher elution strength than methanol at the mobile phase composition of acetonitrile–water–phosphoric acid 0.1% of 50:47.5:2.5 (pH 4.0), the previous interaction of piperine with the stationary phase might be changed, in which piperine might be more solvated in the mobile phase. As a result, piperine was eluted at an early retention time of 0.516 min.

Based on the observation that the exchange of methanol by acetonitrile in the mobile phase and the presence of phosphoric acid had a dramatic effect on the retention time of piperine, various acetonitrile–methanol–water ratios without phosphoric acid were evaluated. In these experiments, phosphoric acid was not included in the mobile phase as acids can substantially reduce the life-time of the column. The effects of the mobile phase composition on peak retention (Tr), peak resolution (Rs), and peak tailing factor (T) are shown in Table 1.

When scrutinizing this table, it can be concluded that the optimal mobile phase composition consisted of an acetonitrile–methanol–water mixture with a volume ratio of 65:5:30, which corresponds to the calculated polarity index of 7.08. Figure 4 shows the resulting baseline chromatographic separation of bis-demethoxycurcumin, demethoxycurcumin, curcumin, and piperine of the methanolic sample containing 10 μ g/mL *C. longa* and 5 μ g/mL *P. nigrum* extracts. Curcumin was eluted at 5.58 min with an Rs of 1.68 and a T of 1.16; piperine was eluted at 7.91

min with an Rs of 3.57 and T of 1.06. The obtained values of Rs, T, N of curcumin and piperine meet the requirements of qualified peak in HPLC ($T < 1.5$, $R_s > 1.5$ and $N > 2000$ [17,29].

3.4. Specificity and selectivity

The method specificity is represented in Figure 5. There are no peaks observed at the retention time of curcumin and piperine in the blank sample (Figure 5a). The retention time for curcumin and piperine in the microparticle formulation sample (Figure 5c) is following the standard solution of curcumin and piperine (Figure 5b). The result indicates the method specificity. The method selectivity is also demonstrated by the baseline separation parameters obtained for curcumin and piperine at which the Rs values for curcumin and piperine are larger than 1.5 (Figure 5c). The curcumin peak was distinguishable from the demethoxycurcumin peak with the Rs of 1.6 and T of 1.1. These results confirm the selectivity of the developed method.

3.5. System suitability test

The optimized HPLC method was subjected to a system suitability test. A sample was injected six times in the system and the T, N, AUC, Tr, and Rs were determined. As shown in Table 2, the RSD values of all these parameters for curcumin as well as piperine were below 2% which indicates all parameters of the proposed HPLC method satisfy the USP and ICH standards. Therefore, the developed HPLC method is concluded to be suitable and effective for the analysis [17].

3.6. Linearity, range and sensitivity

Calibration curves were acquired by plotting peak area as a function of the corresponding concentration (Figure 6). Table 3 shows the regression parameters resulted from the least-square method. As shown in Figure 6 and Table 3, linearity was confirmed at concentration ranges




Figure 5. Chromatographic separation for specificity and selectivity assessment. (a) blank; (b) standard solution containing curcumin (10 μ g/mL) and piperine (5 μ g/mL); (c) solid dispersion based microparticle (9.9 μ g/mL sample) containing *C. longa* and *P. nigrum*. C = curcumin, P = piperine, BDMC = bis-demethoxycurcumin, DMC = demethoxycurcumin.

Table 2. System suitability test of the developed method.

Injection	Tr (min)		AUC		T		Rs		N	
	C	P	C	P	C	P	C	P	C	P
1	5.421	7.660	226134	283307	1.205	1.065	1.624	3.574	8167.5	9561.8
2	5.588	7.914	229048	284215	1.238	1.071	1.649	3.576	8283.6	9356.3
3	5.642	7.970	230929	284790	1.215	1.072	1.680	3.578	8144.9	9243.9
4	5.652	7.994	231654	285454	1.205	1.098	1.690	3.576	8198.7	9409.2
5	5.644	7.974	231274	284831	1.197	1.097	1.688	3.577	8300.8	9474.6
6	5.636	7.977	231993	285380	1.202	1.094	1.690	3.575	8145.6	9353.2
Mean	5.597	7.915	230172	284662	1.210	1.083	1.670	3.576	8206.8	939987
SD	0.089	0.128	2230	803	0.015	0.015	0.027	0.001	69.2	109.8
RSD (%)	1.595	1.614	0.969	0.282	1.221	1.389	1.646	0.040	0.84	1.17

C = Curcumin; P=Piperine; Tr = retention time; Rs = resolution; T = tailing factor; N = theoretical plate number; AUC = Area Under Curve.




Figure 6. Calibration curves of piperine (a) and curcumin (b) standard following the method developed in this study ($n = 3$). The series concentrations of calibration samples were 1.25; 2.50; 5.00; 7.50; 10.00; 12.50; 15.00 $\mu\text{g/mL}$ for piperine and 2.50; 5.00; 10.00; 15.00; 20.00; 25.00; 30.00 $\mu\text{g/mL}$ for curcumin.

of 1.25–15.00 $\mu\text{g/mL}$ and 2.00–30.00 $\mu\text{g/mL}$ for piperine and curcumin, respectively, as the correlation coefficient, r , within these ranges were higher than 0.999. Also, linearity can be further evaluated by calculation of the RSD% of the slope values, which did not exceed 2.0 % [30].

Method sensitivity was tested by determining LOD and LOQ. The calculated LOD of piperine and curcumin was 0.27 $\mu\text{g/mL}$ and 0.42 $\mu\text{g/mL}$, respectively, and the calculated LOQ was 0.91 $\mu\text{g/mL}$ and 1.41 $\mu\text{g/mL}$ for piperine and curcumin, respectively.

3.7. Accuracy and precision

Accuracy and precision studies were conducted according to the ICH recommendations [19] on three different levels of sample concentrations which are low, medium, and high of analyte concentration. The three concentration levels used in this study were 1.25; 7.50; 15.00 and 2.50; 15.00; 30.00 for piperine and curcumin, respectively.

Table 4 summarizes the accuracy and precision data. The results show the intra-day and inter-day recovery of piperine were between 99.54–101.50% and 99.38–99.89%, respectively while the intra-day and inter-day recovery of curcumin were between 100.78–102.51% and 101.15–102.47%, respectively. Repeatability and intermediate precision of the developed method were evaluated by calculating RSD obtained in one day and 3 different days on freshly prepared samples. As compiled in Table 4, it is shown that intra-day and inter-day RSD of piperine were 0.53–0.95% and 0.13–1.44%, respectively; intra-day and inter-day RSD of curcumin were 0.28–1.621% and 0.46–1.14%, respectively.

According to calculated recovery and RSD values, these reported data (Table 4) show that the recovery and RSD values fully satisfy within the generally accepted range of AOAC (90–107% recovery and <5.3% of

RSD) [21], which suggests a high level of accuracy and precision of the developed method.

3.8. Robustness

The robustness parameters of the developed method were obtained by investigating the impact of slight changes in the chromatographic conditions on peak retention time and peak area. Moderate changes in volume ratios of mobile phase composition ($\pm 2\%$), flow rate ($\pm 0.1 \text{ mL/min}$), and detection wavelength ($\pm 2 \text{ nm}$), however, did not significantly affect the retention time and peak area as indicated by the RSD value of <2% (see Table 5).

3.9. Application of the developed method

The developed RP-HPLC method was applied to simultaneously determine the concentration of curcumin and piperine in the microparticle formulation sample containing multicomponent appears in *C. longa* and *P. nigrum* extracts. The developed method was successfully validated to separate multi-compounds in the microparticle formulation i.e. curcumin from its curcuminoids family, and piperine (Table 6). As shown in Table 6, the observed content of curcumin and piperine in the microparticles was $22.42 \pm 0.67 \text{ w/w \%}$ and $9.04 \pm 0.67 \text{ w/w \%}$, respectively. Using reference standards, it was found with the developed HPLC method that *C. longa* extract contained 75.38 w/w % curcumin and *P. nigrum* extract contained 98.97 w/w % piperine. As the claimed percentages of *C. longa* and *P. nigrum* extracts in the microparticles were 30 w/w % and 10 w/w %, respectively, it can be calculated that the theoretical amounts of curcumin and piperine in the microparticles should be 22.61 w/w %

Table 3. Validation parameters for piperine and curcumin (n = 3).

Parameter	Piperine	Curcumin
Linearity range (µg/mL)	1.25–15.00	2.50–30.00
Linear equation	$Y = 140779x + 11549$	$Y = 42144x + 1559.5$
Correlation coefficient of (r)	0.9990	0.9994
RSD of slope	1.017	0.784
Limit of Detection (µg/mL)	0.27	0.42
Limit of Quantification (µg/mL)	0.91	1.41

between 90.0% and 110.0% [31]. Thus, the curcumin and piperine content found in the solid dispersion-based microparticle formulation using the developed method complies with the product specification of the USP monography. The results from the assay of curcumin and piperine confirm the successful implementation of the proposed HPLC method for the simultaneous quantification of curcumin and piperine in microparticles formulation containing multicomponent i.e. curcumin and piperine of *C. longa* and *P. nigrum* extracts. Furthermore, as the running time of HPLC analysis is only 10 min, the proposed method is suitable to

Table 4. Results of intra-day and inter-day accuracy and precision n = 3.

Compound	Final concentration (µg/ml)	Intra-day			Inter-day		
		Recovered quantity (µg/ml)	Recovery (%)	RSD (%)	Recovered quantity (µg/ml)	Recovery (%)	RSD (%)
Curcumin	2.50	2.54 ± 0.04	101.49	1.47	2.61 ± 0.02	102.47	0.79
	15.00	15.12 ± 0.25	100.78	1.62	15.28 ± 0.17	101.88	1.14
	30.00	30.75 ± 0.09	102.51	0.28	31.39 ± 0.14	101.15	0.46
Piperine	1.25	1.27 ± 0.01	101.50	0.95	1.25 ± 0.01	99.79	1.08
	7.50	7.47 ± 0.07	99.54	0.94	7.45 ± 0.01	99.38	0.13
	15.00	15.21 ± 0.08	101.38	0.53	14.98 ± 0.22	99.89	1.44

Table 5. Robustness test of the developed method (n = 3).

Condition	Curcumin				Piperine			
	AUC		Tr (min)		AUC		Tr (min)	
	mean	RSD (%)	mean	RSD (%)	mean	RSD (%)	mean	RSD (%)
Selected mobile phase composition is acetonitrile: methanol: water = 65:5:30 vol-%								
60:10:30 vol-%	21273.6	0.72	5.58	0.10	25209.7	0.308	7.84	0.09
70:5:25 vol-%	22796.7	0.54	5.66	0.10	25358.3	0.500	7.95	0.49
Flow rate of 1 mL/min								
0.9 mL/min	21320.7	0.32	5.49	0.02	25296.3	0.955	7.83	0.05
1.1 mL/min	21255.7	0.83	5.82	0.43	25164.6	0.501	7.99	0.12
Selected detection wavelength of 353 nm								
351 nm	21120.3	1.15	5.61	0.16	325481.6	0.329	7.85	0.63
355 nm	23781.4	0.45	5.63	0.07	252137.6	0.264	7.92	0.17

Tr = retention time; AUC = Area Under Curve.

and 9.90 w/w %, respectively. Using these percentages, it can be calculated that the recovery of curcumin and piperine from the microparticles was 99.14% and 91.31%, respectively, using the developed HPLC method (Table 6).

USP monograph on curcuminoids dosage form requires that the content of the active compound in the dosage form products should be

be applied during routine analysis as part of quality control procedures in the formulation of the microparticles.

4. Conclusion

A suitable, rapid, accurate, and precise RP-HPLC method was developed for the simultaneous determination of curcumin and piperine in a solid dispersion-based microparticle formulation. The developed method offers a linear response across a wide range of analyte concentrations with satisfactory method sensitivity. Furthermore, the proposed RP-HPLC method guarantees the extension of the column lifetime and HPLC system due to the absence of acid in the mobile phase and the use of commonly used organic solvents, i.e. acetonitrile and methanol.

Declarations

Author contribution statement

Dewi Setyaningsih: Conceived and designed the experiments; Analyzed and interpreted the data; Contributed reagents, materials, analysis tools or data; Wrote the paper.

Yosua Agung Santoso: Performed the experiments.

Yustina Sri Hartini: Analyzed and interpreted the data.

Table 6. Curcumin and piperine content in microparticles.

Sample	Amount found in microparticles	
	Curcumin (w/w %)	Piperine (w/w %)
1	21.17	8.41
2	22.90	8.39
3	22.58	8.55
4	22.90	9.98
5	22.82	9.40
6	22.17	9.50
Mean	22.42	9.04
SD	0.67	0.67
RSD (%)	3.05	7.47

Formulation claim: 30 w/w % *C. longa* and 10 w/w % *P. nigrum* extracts.

Yosi Bayu Murti: Contributed reagents, materials, analysis tools or data.

Wouter L. J. Hinrichs: Conceived and designed the experiments; Analyzed and interpreted the data; Wrote the paper.

Christine Patramurti: Conceived and designed the experiments; Performed the experiments.

Funding statement

This work was supported by Ministry of Research and Technology/ National Research and Innovation Agency of Republic Indonesia (contract number of 227/SP2H/LT/DRPM/2019, B/1436.8/L5/RA.00/2019, 029/Penel/LPPM-USD/IV/2019).

Data availability statement

Data included in article/supplementary material/referenced in article.

6

Declaration of interests statement

The authors declare no conflict of interest.

Additional information

No additional information is available for this paper.

References

- [1] S.J. Hewlings, D.S. Kalman, Curcumin: a review of its effects on human health, *Foods* 6 (2017) 92.
- [2] R. Tabrizi, et al., The effects of curcumin-containing supplements on biomarkers of inflammation and oxidative stress: a systematic review and meta-analysis of randomized controlled trials, *Phytother. Res.* 33 (2019) 253–262.
- [3] E.U. Ikonne, V.O. Ikpeazu, E.A. Ugbogu, The potential health benefits of dietary natural plant products in age related eye diseases, *Heliyon* 6 (2020).
- [4] S.C. Gupta, S. Patchva, B.B. Aggarwal, Therapeutic roles of curcumin: lessons learned from clinical trials, *AAPS J.* 15 (2012) 195–218.
- [5] Y.J. Wang, et al., Stability of curcumin in buffer solutions and characterization of its degradation products, *J. Pharmaceut. Biomed. Anal.* 15 (1997) 1867–1876.
- [6] P. Srinarong, et al., Improved dissolution behavior of lipophilic drugs by solid dispersions: the production process as starting point for formulation considerations, *Expet Opin. Drug Deliv.* 8 (2011) 1121–1140.
- [7] D.B. Mhaske, S. S. M. KR, Role of piperine as an effective bioenhancer in drug absorption, *Pharm. Anal. Acta* 7 (2018).
- [8] G. Shoba, et al., Influence of piperine on the pharmacokinetics of curcumin in animals and human volunteers, *Planta. Med.* 64 (1998) 353–356.
- [9] A. Rohman, Analysis of curcuminoids in food and pharmaceutical products, *Int. Food Res. J.* 19 (2012) 19–27.
- [10] H.K. Syed, et al., Stability indicating HPLC-UV method for detection of curcumin in Curcuma longa extract and emulsion formulation, *Food Chem.* 170 (2015) 321–326.
- [11] M.R. Peram, et al., Single robust RP-HPLC analytical method for quantification of curcuminoids in commercial turmeric products, Ayurvedic medicines, and nanovesicular systems, *J. Liq. Chromatogr. Relat. Technol.* 40 (2017) 487–498.
- [12] Y.B. Murti, et al., UV-vis spectroscopy to enable determination of the dissolution behavior of solid dispersions containing curcumin and piperine, *JYP* 11 (2019) 26–30.
- [13] C. Moorthi, et al., Application of validated RP-HPLC-PDA method for the simultaneous estimation of curcumin and piperine in Eudragit E 100 nanoparticles, *J. Pharm. Res.* 7 (2013) 224–229.
- [14] P. Sethi, et al., Development and validation of a reversed phase HPLC method for simultaneous determination of curcumin and piperine in human plasma for application in clinical pharmacological studies, *J. Liq. Chromatogr. Relat. Technol.* 32 (2009) 2961–2974.
- [15] X.-M. Wang, et al., Validated HPLC-MS/MS method for simultaneous determination of curcumin and piperine in human plasma, *TJPR* 11 (2012) 621–629.
- [16] K.C. Saha, H.P. Seal, M.A. Noor, Isolation and characterization of piperine from the fruits of black pepper (*Piper nigrum*), *J. Bangladesh Agric. Univ.* 11 (2013) 6.
- [17] USP, chromatography USP 621, in: National Formulary 37, Pharmacopeial Convention Inc, USA, 2009.
- [18] H.H. Tonnesen, J. Karlsen, G.B. van Henegouwen, Studies on curcumin and curcuminoids. VIII. Photochemical stability of curcumin, *Z. Lebensm. Unters. Forsch.* 183 (2) (1986) 116–122.
- [19] ICH, Validation of analytical procedures: text and methodology, in: International Conference on Harmonisation, in Q2(R1), 2005. Geneva: Geneva.
- [20] G. González, A. Herrador, A practical guide to analytical method validation, including measurement uncertainty and accuracy profiles, *Trends Anal. Chem.* 26 (2007) 227–238.
- [21] AOAC, Official Methods of Analysis of AOAC International, in: J. George W. Latimer (Ed.), Guidelines for Standard Method Performance Requirements, AOAC International, Gaithersburg, Maryland, USA, 2012, p. p9. Appendix F.
- [22] Y.C. Lee, Method validation for HPLC analysis of related substances in pharmaceutical drug products, in: C.C. Chan, et al. (Eds.), Analytical Method Validation and Instrument Performance Verification, John Wiley & Sons, Inc., 2004, pp. 27–49.
- [23] D. Mumane, G.P. Martin, C. Marriott, Validation of a reverse-phase high performance liquid chromatographic method for concurrent assay of a weak base (salmeterol xinafoate) and a pharmacologically active steroid (fluticasone propionate), *J. Pharmaceut. Biomed. Anal.* 40 (2006) 1149–1154.
- [24] S. Espinosa, E. Bosch, M. Rosés, Retention of ionizable compounds on HPLC. 5. pH scales and the retention of acids and bases with acetoneitrile–water mobile phases, *Anal. Chem.* 72 (2000) 5193–5200.
- [25] S. Li, M. Tian, K.H. Row, Effect of mobile phase additives on the resolution of four bioactive compounds by RP-HPLC, *Int. J. Mol. Sci.* 11 (2010) 2229–2240.
- [26] R. McClain, E. Streckfuss, High-throughput purification in support of pharmaceutical discovery, in: P. Knochel (Ed.), Comprehensive Organic Synthesis II, second ed., Elsevier, Amsterdam, 2014, pp. 160–180.
- [27] R. LoBrutto, et al., Effect of the eluent pH and acidic modifiers in high-performance liquid chromatography retention of basic analytes, *J. Chromatogr. A* 913 (2001) 173–187.
- [28] K. Guan, D.C. Palmer, Effects of trifluoroacetic acid concentrations in mobile phases on HPLC retention of zwitterionic and weakly basic triazole derivatives, *J. Liq. Chromatogr. Relat. Technol.* 29 (2006) 415–430.
- [29] L.R. Snyder, J.J. Kirkland, J.L. Glajch, Basics of separation, in: Practical HPLC Method Development, A Wiley-Interscience Publication, John Wiley & Sons, Inc., USA, 1997, pp. 21–58.
- [30] R.A. Shaalan, et al., Validated HPTLC methods for determination of some selected antihypertensive mixtures in their combined dosage forms 52, *Bulletin of Faculty of Pharmacy, Cairo University*, 2014, pp. 225–237.
- [31] USP, Dietary supplements monographs curcuminoids, in: USP Pharmacists' Pharmacopeia, U.S Pharmacopeia, Rockville, 2019, pp. 1–7.

Isocratic high-performance liquid chromatography (HPLC) for simultaneous quantification of curcumin and piperine in a microparticle formulation containing *Curcuma longa* and *Piper nigrum*

ORIGINALITY REPORT

21%

SIMILARITY INDEX

16%

INTERNET SOURCES

16%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|----|
| 1 | www.rug.nl
Internet Source | 2% |
| 2 | C. Moorthi, C. Senthil Kumar, S. Mohan, Kiran Krishnan, K. Kathiresan. "Application of validated RP-HPLC-PDA method for the simultaneous estimation of curcumin and piperine in Eudragit E 100 nanoparticles", <i>Journal of Pharmacy Research</i> , 2013
Publication | 1% |
| 3 | pubmed.ncbi.nlm.nih.gov
Internet Source | 1% |
| 4 | www.hindawi.com
Internet Source | 1% |
| 5 | Rasha A. Shaalan, Tarek S. Belal, Fawzy A. El Yazbi, Sohila M. Elonsy. "Validated HPTLC methods for determination of some selected antihypertensive mixtures in their combined | 1% |

dosage forms", Bulletin of Faculty of Pharmacy, Cairo University, 2014

Publication

6	repository.ubaya.ac.id Internet Source	1 %
7	Submitted to Universitas Merdeka Malang Student Paper	1 %
8	Submitted to Syiah Kuala University Student Paper	1 %
9	Submitted to Higher Education Commission Pakistan Student Paper	1 %
10	Submitted to Jawaharlal Nehru Technological University Student Paper	<1 %
11	bmcchem.biomedcentral.com Internet Source	<1 %
12	www.bioseek.eu Internet Source	<1 %
13	jifi.farmasi.univpancasila.ac.id Internet Source	<1 %
14	mafiadoc.com Internet Source	<1 %
15	Submitted to Universiti Sains Malaysia Student Paper	<1 %

- | | | |
|----|--|------|
| 16 | repositorio.ufsm.br
Internet Source | <1 % |
| 17 | www.pubmedcentral.nih.gov
Internet Source | <1 % |
| 18 | www.scielo.br
Internet Source | <1 % |
| 19 | Epi Erpina, Mohamad Rafi, Latifah Kosim Darusman, Arum Vitasari, Budi Riza Putra, Eti Rohaeti. " Simultaneous quantification of curcuminoids and xanthorrhizol in by high-performance liquid chromatography ", Journal of Liquid Chromatography & Related Technologies, 2017
Publication | <1 % |
| 20 | jast-journal.springeropen.com
Internet Source | <1 % |
| 21 | www.ajol.info
Internet Source | <1 % |
| 22 | Prerana Sethi, Virendra K. Dua, S. Mohanty, S. K. Mishra, Rajeev Jain, G. Edwards. "Development and Validation of a Reversed Phase HPLC Method for Simultaneous Determination of Curcumin and Piperine in Human Plasma for Application in Clinical Pharmacological Studies", Journal of Liquid | <1 % |

Chromatography & Related Technologies, 2009

Publication

23

pharmacophorejournal.com

Internet Source

<1 %

24

engagedscholarship.csuohio.edu

Internet Source

<1 %

25

pubs.rsc.org

Internet Source

<1 %

26

Mireille Hassoun, Maria Malmlof, Otto Scheibelhofer, Abhinav Kumar et al. "Use of PBPK modelling to evaluate the performance of DissolvIt, a biorelevant dissolution assay for orally inhaled drug products", Molecular Pharmaceutics, 2019

Publication

<1 %

27

ijpsr.com

Internet Source

<1 %

28

Hema Chaudhary, Kanchan Kohli, Saima Amin, Saurabh Arora, Vikash Kumar, Sushila Rathee, Permender Rathee. "DEVELOPMENT AND VALIDATION OF RP-HPLC METHOD FOR SIMULTANEOUS ESTIMATION OF DICLOFENAC DIETHYLAMINE AND CURCUMIN IN TRANSDERMAL GELS", Journal of Liquid Chromatography & Related Technologies, 2012

<1 %

29

bjp.sagepub.unboundmedicine.com

Internet Source

<1 %

30

faf.vfu.cz

Internet Source

<1 %

31

sphinxsai.com

Internet Source

<1 %

32

www.e-journals.in

Internet Source

<1 %

33

B. JOE. "Biological Properties of Curcumin-Cellular and Molecular Mechanisms of Action", Critical Reviews in Food Science and Nutrition, 3/1/2004

Publication

<1 %

34

Hasan Basan, Nilgün Günden Göğer, Nusret Ertaş, M.Tevfik Orbey. "Quantitative determination of piroxicam in a new formulation (piroxicam- β -cyclodextrin) by derivative UV spectrophotometric method and HPLC", Journal of Pharmaceutical and Biomedical Analysis, 2001

Publication

<1 %

35

Immaculada Canals, Fadoua Z. Oumada, Martí Rosés, Elisabeth Bosch. "Retention of ionizable compounds on HPLC. 6. pH measurements with the glass electrode in

<1 %

methanol-water mixtures", Journal of Chromatography A, 2001

Publication

36

Teresa Cecchi. "Ion Pairing Chromatography", Critical Reviews in Analytical Chemistry, 2008

Publication

<1 %

37

www.ijppr.humanjournals.com

Internet Source

<1 %

38

www.pharmanest.net

Internet Source

<1 %

39

Bo Hyun Lee, Hyun A. Choi, Mi-Ri Kim, Jungil Hong. "Changes in chemical stability and bioactivities of curcumin by ultraviolet radiation", Food Science and Biotechnology, 2013

Publication

<1 %

40

Malleswara Rao Peram, Sachin R. Patil, Vijay M. Kumbar, Manohar S. Kugaji, Kishore G. Bhat, Prakash V. Diwan, Sunil Jalalpure. "An RP-HPLC Method for Quantitative Analysis of Linagliptin Entrapped in Nanotransfersomes and its Application to Skin Permeation Studies", Current Pharmaceutical Analysis, 2021

Publication

<1 %

41

Marlene Maria Fregonezi-Nery, Marcela Maria Baracat, Rubia Casagrande, Hodnei Takashi

<1 %

Machado et al. "Validação de métodos para determinação de fluoxetina em cápsulas", Química Nova, 2008

Publication

42

Ornchuma Naksuriya, Siriporn Okonogi, Raymond M. Schiffelers, Wim E. Hennink. "Curcumin nanoformulations: A review of pharmaceutical properties and preclinical studies and clinical data related to cancer treatment", Biomaterials, 2014

Publication

<1 %

43

Wisut Wichitnithad. "A simple isocratic HPLC method for the simultaneous determination of curcuminoids in commercial turmeric extracts", Phytochemical Analysis, 2009

Publication

<1 %

44

diginole.lib.fsu.edu

Internet Source

<1 %

45

repositorio.unicamp.br

Internet Source

<1 %

46

www.camag.com

Internet Source

<1 %

47

www.dovepress.com

Internet Source

<1 %

48

Alexey Makarov, Rosario LoBrutto, Yuri Kazakevich. "Liophilic Mobile Phase Additives in Reversed Phase HPLC", Journal of Liquid

<1 %

49

Elliott L. Rodriguez, Chenhua Zhang, Ashley G. Woolfork, Zhao Li et al. "Analysis of curcumin and piperine in biological samples by reversed-phase liquid chromatography with multi-wavelength detection", Journal of Chromatography B, 2021

Publication

<1 %

50

Sonia Espinosa, Elisabeth Bosch, Martí Rosés. "Retention of Ionizable Compounds on HPLC. 5. pH Scales and the Retention of Acids and Bases with Acetonitrile–Water Mobile Phases", Analytical Chemistry, 2000

Publication

<1 %

51

Yi Lu, Ning Tang, Ruyue Lian, Jianping Qi, Wei Wu. "Understanding the relationship between wettability and dissolution of solid dispersion", International Journal of Pharmaceutics, 2014

Publication

<1 %

52

bioone.org

Internet Source

<1 %

53

portal.research.lu.se

Internet Source

<1 %

54

İncilay Süslü, Sacide Altınöz. "Two derivative spectrophotometric determinations of indapamide in pharmaceutical dosage forms", *Journal of Pharmaceutical and Biomedical Analysis*, 2002

Publication

<1 %

55

"Curcumin: Preventive and Therapeutic Properties in Laboratory Studies and Clinical Trials", *Antioxidants & Redox Signaling*, 03/2008

Publication

<1 %

56

1library.net

Internet Source

<1 %

57

Haroon Khalid Syed, Kai Bin Liew, Gabriel Onn Kit Loh, Kok Khiang Peh. "Stability indicating HPLC–UV method for detection of curcumin in *Curcuma longa* extract and emulsion formulation", *Food Chemistry*, 2015

Publication

<1 %

58

Kumar, Sandeep, Viney Lather, and Deepti Pandita. "Stability indicating simplified HPLC method for simultaneous analysis of resveratrol and quercetin in nanoparticles and human plasma", *Food Chemistry*, 2016.

Publication

<1 %

59

Li, Y.G.. "Robustness tests on the United States Pharmacopoeia XXVI HPLC assay for

<1 %

ginsenosides in Asian and American ginseng using an experimental design", *Analytica Chimica Acta*, 20050422

Publication

60

Maximiliano da Silva Sangoi, Daniele Rubert Nogueira, Lucélia Magalhães da Silva, Diogo Paim Leal, Sérgio Luiz Dalmora. "Validation of a Stability Indicating Reversed Phase LC Method for the Determination of Fluticasone Propionate in Pharmaceutical Formulations", *Journal of Liquid Chromatography & Related Technologies*, 2008

Publication

61

McClain, R., and E. Streckfuss. "9.07 High-Throughput Purification in Support of Pharmaceutical Discovery", *Comprehensive Organic Synthesis II*, 2014.

Publication

62

Sang-Ho Lee, Sang-Ho Lee, Jong-Hyuk Lee, Joo-Eun Kim. "Development, Validation, and Application of a High-Performance Liquid Chromatography Method for the Analysis of Dissolution Samples of Telmisartan and Rosuvastatin Calcium Double-Layer Formulation", *Current Pharmaceutical Analysis*, 2021

Publication

63

[Scholar.ufs.ac.za](https://scholar.ufs.ac.za)

Internet Source

<1 %

<1 %

<1 %

<1 %

64

Tomoko Hayashi, Hisao Oka, Yuko Ito, Tomomi Goto et al. "An HPLC Method for the Analysis of Marigold Color in Food Using Lutein as an Indicator", Journal of Liquid Chromatography & Related Technologies, 2007

Publication

<1 %

65

[es.scribd.com](https://www.es.scribd.com)

Internet Source

<1 %

66

kclpure.kcl.ac.uk

Internet Source

<1 %

67

www.japsonline.com

Internet Source

<1 %

68

www.koreascience.kr

Internet Source

<1 %

69

"Analytical chemistry of chemesthetic compounds", Chemesthesis, 2016.

Publication

<1 %

70

Poonam Kushwaha, Babita Shukla, Jyotsana Dwivedi, Sumedha Saxena. "Validated high-performance thin-layer chromatographic analysis of curcumin in the methanolic fraction of *Curcuma longa* L. rhizomes",

<1 %

71

"High-Performance Thin-Layer Chromatography (HPTLC)", Springer Science and Business Media LLC, 2011

Publication

<1 %

72

Maria Manconi, Maria Letizia Manca, Elvira Escribano-Ferrer, Elisabet Martí Coma-Cros et al. "Nanoformulation of curcumin-loaded eudragit-nutriosomes to counteract malaria infection by a dual strategy: Improving antioxidant intestinal activity and systemic efficacy", International Journal of Pharmaceutics, 2019

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On