




# LATIN AMERICAN JOURNAL OF PHYSICS EDUCATION

www.lajpe.org

Volume 15

Number 1

March 2021



A publication sponsored by the Latin American Physics Education Network  
and the Institute of Science Education



---

---

# LATIN AMERICAN JOURNAL OF PHYSICS EDUCATION

---

---

Electronic version of this journal can be downloaded free of charge from the web-resource:

<http://www.lajpe.org>

## Production and technical support

Rubén Sánchez Sánchez  
rbnsnchz@yahoo.com.mx  
Martin Sánchez Huerta  
Halcon.n@live.com.mx

Latin American Journal of Physics  
Education is indexed in:

DOAJ

Dialnet

latindex

EBSCO  
PUBLISHING

INDEX COPERNICUS  
INTERNATIONAL

## EDITORIAL POLICY

Latin American Journal of Physics Education (LAJPE) is a peer-reviewed, electronic international journal for the publication of papers of instructional and cultural aspects of physics. Articles are chosen to support those involved with physics courses from introductory up to postgraduate levels.

Papers may be comprehensive reviews or reports of original investigations that make a definitive contribution to existing knowledge. The content must not have been published or accepted for publication elsewhere, and papers must not be under consideration by another journal.

This journal is published quarterly (March, June, September and December), by the Latin American Physics Education Network (LAPEN) and the Institute of Science Education. Manuscripts should be submitted to [boubarkic@gmail.com](mailto:boubarkic@gmail.com) or [lajpe@lapen.org.mx](mailto:lajpe@lapen.org.mx). Further information is provided in the "Instructions to Authors" on [www.lajpe.org](http://www.lajpe.org)

Direct inquiries on editorial policy and the review process to: C. Boubarki, Editor in Chief, Instituto Pedagógico Latinoamericano y Caribeño (Cuba), calle 154 No. 12906 entre 129 y 133 Reparto Reynold Garcia, Matanzas, Cuba CP 40100. Phone: 53 45 265794

ISSN 1870-9095

Copyright © 2021 Latin American Physics Education Network. ([www.lajpe.org](http://www.lajpe.org))

## INTERNATIONAL ADVISORY COMMITTEE

Ann-Marie Pendrill, Göteborgs University (Sweden)  
Bayram Akarsu, Erciyes University (Turkey)  
Carl Wenning, Illinois State University (USA)  
Diane Grayson, Andromeda Science Education (South Africa)  
David Sokoloff, University of Oregon (USA)  
Dean Zollman, Kansas State University (USA)  
Edward Redish, University of Maryland (USA)  
Freidrich Herrmann, University of Karlsruhe (Germany)  
Hiroshi Kawakatsu, Kagawa University (Japan)  
Jorge Barojas Weber, Universidad Nacional Autónoma de México (México)  
Laurence Viennot, Université Paris 7 (France)  
Marisa Micheli, University of Udine (Italy)  
Marco Antonio Moreira, Universidade Federal do Rio Grande do Sul (Brazil)  
Minella Alarcón, UNESCO (France)  
Orhan Karamustafaoğlu, Amasya University, (Turkey)  
Pratibha Jolly, University of Delhi (India)  
Priscilla Laws, Dickinson College (USA)  
Ton Ellermeijer, (Netherlands)  
Verónica Tricio, University of Burgos (Spain)  
Vivien Talisayon, University of the Philippines (Philippines)

## EDITORIAL BOARD

Deise Miranda, Universidade Federal do Rio de Janeiro (Brasil)  
Eduardo Moltó, Instituto Superior Pedagógico José Varona (Cuba)  
Eduardo Montero, Escuela Superior Politécnica del Litoral (Ecuador)  
Josefina Barrera, Universidade do Estado do Amazonas (Brasil)  
Julio Benegas, Universidad Nacional de San Luis (Argentina)  
Leda Roldán, Universidad de Costa Rica (Costa Rica)  
Celso Ladera, Universidad Simón Bolívar (Venezuela)  
Nelson Arias Ávila, Universidad Distrital, Bogotá (Colombia)  
Octavio Calzadilla, Universidad de la Habana (Cuba)  
Ricardo Buzzo Garrao, Pontificia Universidad Católica de Valparaíso (Chile)  
Zulma Gangoso, Universidad Nacional de Córdoba (Argentina)

## EDITOR-IN-CHIEF

Carlos Boubarki, Instituto Pedagógico Latinoamericano y Caribeño (Cuba)

## ASSOCIATED EDITOR

Josip Slisko, Benemérita Universidad Autónoma de Puebla (México)

# LATIN AMERICAN JOURNAL OF PHYSICS EDUCATION

Volume 15, Number 1, March 2021

## CONTENTS/CONTENIDO

### Papers/Artículos

- Solar radiation and the fraction of the visible sun's disk during an eclipse  
Ign Edi Santosa 1301
- Project method in the education background: a review of recent literature  
Mara Fernanda Parisoto, Mauricio Romani, Emmanuel Zullo Godinho 1302
- Habilidades esenciales para el siglo XXI a través de la educación STEM  
Daniel Gomes da Silva, Maud Rejane de Souza, Josefina Barrera Kalhil 1303
- Learning by projects: theory and practice in  
Mara F. Parisoto, Emmanuel Z. G, Marcos H., Abreu de O., Robert F., Ígor P. 1304
- Curricular configurations of the contemporaneous academic training of medical  
physicists in Rio Grande do Sul- Brazil  
Laura Larré G., Cleidilene Ramos M., Priscila Silva R., Mirko Salomón A. S. 1305
- Eppur se muove", but, we do not understand how  
José Joaquín Lunazzi, Antonio Marcos de Souza 1306
- Euler–Bernoulli beam theory in the kitchen: The fridge-violin  
Luis Darmendrail, Andreas Müller 1307
- Evaluación de un montaje para la práctica de laboratorio efecto Hall en  
semiconductores y modificaciones para elevar su calidad  
Arcelio A. Hernández Ferreira 1308
- Uso alternativo del cronómetro de un teléfono celular en la medición del período de  
oscilación de un péndulo para experimentación en educación a distancia  
Terrazas-Bandala Luisa Piroshka *et al.* 1309
- Abordagem da Lei de Ohm para Estudantes com Deficiência Visual no Ensino  
Superior  
M. Velloso, M. Arana, V. Acioly, A C F Santos 1310
- Development of an Educational Environment by Using Graphical User Interfaces  
Applied to Heat Transfer Problems  
Fran Sérgio Lobato, Fábio de Oliveira Arouca 1311

*continued/continuación*

*continued/continuación*

Computational fluid dynamic simulation of a gas-solid fluidized bed system: A dense phase analysis L.P. Olivo-Arias, L.G. De Araujo, J.B. Rojas-Trigos	1312
Uso de simuladores Phet, para la enseñanza del comportamiento de gases ideales Manuel Sandoval Martínez, Janette Moreno Sandoval, César Mora	1313
Termodinâmica e Revolução Industrial: Uma abordagem por meio da História e da Epistemologia da Ciência Debora Samir Conceição de Souza, Boniek Venceslau da Cruz Silva	1314
The last quitters: nobody's science and physics Nilo Serpa, Richard Cathcart	1315
Investigación en el aula: un trabajo colegiado Adriana Gómez Reyes, Ángel Homero Flores Samaniego	1316
Estudio de dos modelos de aprendizaje semipresencial en educación superior Emilia López-Iñesta, María T. Sanz	1317
Momento de torsión con Aprendizaje Activo para estudiantes de edad adulta en el nivel Superior Carla Kerlegand Bañales, Rubén Sánchez Sánchez, Elvia Rosa Ruiz Ledezma	1318
Selected Experiments with an Arduino Board to Teach Analog-to-Digital Conversion João E. M. Perea Martins	1319
Reseña 3er Coloquio de los Seminarios Repensar del Instituto Politécnico Nacional Rosa Griselda Nava Galve, Guillermina Ávila, Víctor Hugo Luna, Liliana Suárez Téllez, Daniel Sánchez Guzmán	1511
Grupo de estudio como alternativa formativa del Seminario Repensar las Matemáticas William Andrey Suárez Moya, Liceth Katherin Beltrán Perdomo, Domingo Márquez Ortega	1512

# Solar radiation and the fraction of the visible sun's disk during an eclipse



**Ign Edi Santosa**

*Physics Education Dept., Sanata Dharma University,  
Paingan, Maguwohardjo, Sleman, Yogyakarta 55282, Indonesia.*

**E-mail:** edi@usd.ac.id

(Received 12 December 2020, accepted 15 February 2021)

## Abstract

The change of illumination on the 9<sup>th</sup> March 2016 solar eclipse has been monitored using a light sensor a simple instrument that is available in most laboratories. This observation was conducted in Yogyakarta (7<sup>o</sup> 46'S, 110<sup>o</sup> 20'E). The results were normalized to the illumination on the normal day, i.e., on 5<sup>th</sup> March 2017 and on 8<sup>th</sup> March 2018, to obtain the fraction of the uncovered sun's disk. The minimum illumination during eclipse corresponded to the 80.5% obscuration. This simple method showed a good agreement with the available data obtained from Stellarium.

**Keywords:** Eclipse, Solar radiation, Light sensor.

## Resumen

El cambio de iluminación del eclipse solar del 9 de marzo de 2016 se ha monitorizado mediante un sensor de luz, un instrumento sencillo que está disponible en la mayoría de los laboratorios. Esta observación se realizó en Yogyakarta (7<sup>o</sup> 46'S, 110<sup>o</sup> 20'E). Los resultados se normalizaron a la iluminación del día normal, es decir, el 5 de marzo de 2017 y el 8 de marzo de 2018, para obtener la fracción del disco solar descubierto. La iluminación mínima durante el eclipse correspondió al 80,5% de oscurecimiento. Este sencillo método mostró una buena concordancia con los datos disponibles obtenidos de Stellarium.

**Palabras clave:** Eclipse, Radiación solar, Sensor de luz.

## I. INTRODUCTION

Solar radiation has been studied in various fields. These studies are related to the availability of solar energy, the effect of intensity on the season, and temperature distribution. Visible movement of the sun has been modelled. This model can predict daylight duration and be applied for a practical purpose of harvesting the solar energy [1]. The same model is also used to construct global maps with distribution of daily insolation, and to simulate the passage of the seasons on the Earth [2].

Different instruments have been utilized for measuring the solar radiation. Using a homemade pyranometer Zanetti and Zecca showed the light intensity in arbitrary unit for a full day [3]. In the other experiment the pattern of solar radiation has been monitored; thermocouples and an Eppley Precision Spectral Pyranometer were utilized for measuring temperature of the shingles and total solar radiation, respectively. As expected, the temperature follows the pattern of solar radiation [4].

In contrast to the studies above, during the eclipse solar radiation shows a special behaviour. Therefore, the solar eclipse attracts observers from the entire world. Different articles on solar eclipses can be found in the pedagogic journals in physics. Richard and Smart measured a decrease in negative ion concentration in the atmosphere, and

recorded solar magnitude as well as studied shadow bands [5]. Production of shadow bands during an eclipse has also been modelled [6]. For the intermediate astronomy courses a solar eclipse has been predicted. It is assumed that the moon's position is along the circular orbit. This simplification and the data from the almanac, can be utilised to predict the longitude and latitude for the observation location. Furthermore, the width of the eclipse's shadow on the earth and the duration of totality can also be calculated [7]. By observing the moon's shadow during the solar eclipse, Sawicki determined the radius of the moon's orbit, i.e. 376 000 km [8]. Resources for observing solar eclipses can be found in a recent article [9].

A decrease of solar radiation has been monitored during an eclipse [10]. The measurement was performed using a photometer operated in three spectral regions, i.e. infrared, red and blue. This article showed the variation of the ratio of the light intensity during an eclipse day and at the previous day without the theoretical background. In a different way a model of solar radiation during an eclipse was developed. The algorithm has been tested in the global solar UV radiation during the partial solar eclipse in Italy [11]. The change of illuminance during a solar eclipse of 2006 has been measured [12]. Calibrated equipment that was oriented perpendicular to the light was utilized for measuring the illuminance. In this case a model was

presented based on the assumption that the illuminance is proportional to the unobscured area of the sun’s disk. The model is in a good agreement with measured data.

The total solar eclipse of 9 March 2016 over Indonesia was observed from the east to the west regions of the country. This phenomenon could be seen in different parts of Indonesia. In Indonesia the first contact started at 06:19:41 local time. The longest duration of the eclipse was observed in Jayapura, Papua, i.e. 2 hours 55.05 minutes. The maximum duration of totality of the eclipse was 3.325 minutes, which was observed in Maba, North Maluku.

This solar eclipse was also seen partly from Yogyakarta (7° 46’S, 110° 20’E) located on Java Island. This location is not in the total eclipse path. This phenomenon lasted for 2 hours and 14.74 minutes. The maximum eclipse obscuration was 80.5%. It was fortunate that during this time the sky was mostly clear and cloudless; the eclipse could be monitored clearly and completely from the beginning to the end.

This special occasion makes an interesting study. Here the incoming solar radiation depends on the visible sun’s disk. The solar radiation was monitored easily using a light sensor; a simple instrument that is available in most common physics laboratories. Using a simple formula, the fraction of visible sun’s disk can be determined from the measured data. The experiment presented in this paper can be conducted by any students using a simple light sensor during an eclipse. The students will also learn a direct effect of the eclipse.

**II. THEORY**

The direct solar beam radiation intensity on a horizontal sensor surface  $I_b$  is a function of the sun-earth distance, the solar zenith angle, solar constant and the transmittance function [13]. The solar zenith angle depends on the latitude, the declination angle, and the angular displacement of the sun due to rotation of the earth on its axis. The declination angle as well as the sun-earth distance varies due to the earth’s movement around the sun. These factors give the annual and diurnal variation in the incoming solar radiation. For a particular date the sun-earth distance is relatively constant and therefore the incoming solar radiation on a fixed place only shows the diurnal variation and can be expressed as

$$I_b = I_{max} T \cos \theta_z \tag{1}$$

where  $I_{max}$  is the maximum intensity that is determined by the solar constant and the sun-earth distance;  $\theta_z$  is the solar zenith angle;  $T$  is the transmittance function related to the absorption and scattering processes in the atmosphere that depends on the availability of gases in the atmosphere. For the sun’s altitude  $\alpha$  as seen in Fig. 1, Eq. (1) becomes

$$I_b = I_{max} T \cos (90 - \alpha) \tag{2}$$

In this experiment the light sensor was mounted in an inclined holder (Fig. 2). For this tilted sensor the Eq. (2) should be corrected by tilt angle  $\beta$ , and hence [14-15]

$$I_{bt} = I_{max} T \cos (\beta - \alpha) \tag{3}$$

During the eclipse the moon covers part of the sun’s disk. The incoming solar radiation is proportional to the area of the visible bright disk [11, 16]. Therefore, the correction should be given, and its intensity during eclipse day  $I$  becomes.




FIGURE 1. The solar radiation on the horizontal sensor.




FIGURE 2. The solar radiation on the tilt sensor.

Beside this direct radiation there is also a diffuse radiation  $I_d$ . As a result, the total intensity  $I_T$  on the tilt sensor follows

$$I_T = I_{bt} + I_d \tag{4}$$

$$I = A_r I_T \tag{5}$$

where  $A_r$  represents the fraction of the uncovered solar disk. In this case the fraction of the sun’s area occulted by the moon is defined as eclipse obscuration  $E_o$  and hence


$$A_r = 1 - E_o \tag{6}$$

The fraction of visible sun’s disk can be determined from Eq. (5) by measuring the solar radiation during an eclipse day  $I$  and in the normal day  $I_T$ . These measurements should be in the same experiment’s condition, i.e., the date, the

time, the position (orientation and tilt angle) of the sensor and the atmosphere. The requirement of the date was approximately fulfilled by measuring  $I_T$  on the same week of the eclipse day or at least the same week of the year.

### III. EXPERIMENT

The solar radiation intensity has been measured using a light sensor (BTA Vernier), which is a small photodiode (1mm x 1mm). This sensor was placed inside the glass cylinder and was secured near the tip of the cylinder. This sensor is connected to a laptop via an interface *LabPro* product of Vernier [17]. A laptop provides data acquisition and controls the experiment, and for this purpose it is equipped with software *LoggerPro*. The experimental setup is depicted in Fig. 3. The light sensor is mounted on the holder on an incline of  $25^\circ$ . The sensor position is in front of the holder and perpendicular to this incline. The holder is placed in the balcony oriented to the East, directly toward the sun. All measurements utilize this experimental setup condition.




**FIGURE 3.** The experimental setup. The light sensor (1) is mounted on holder (2). The interface Labpro (3) is used to connect the light sensor to a laptop (4).

The eclipse on 9 March 2016 was predicted to occur from 06:20:32 to 08:35 local time, and it reached a maximum obscuration at 7:23:30 local time. Therefore, on this day the solar radiation was collected continuously from 06:00 local time to 10:00 local time. On the day of the eclipse the sky was mostly clear, although it was in the rainy season. The solar radiation was recorded as illumination in unit lux. Fig. 4 shows the result, the shaded area given highlights the period of the eclipse.


At the beginning, the illumination increased gradually due to the diurnal behaviour. As expected, the effect of eclipse reduced the illumination and in overall showed clearly a dip. If we inspected closely, the decrease in the illumination was not instantaneously following the increase of the covered sun's disk. At first the effect was not dominant, an increase of illumination was still observed. After about 25 minutes from the first contact, the illumination started decreasing, reaching minimum at 07:21 local time and then back to increase. The maximum-recorded illumination was observed at 08:55 local time.

To obtain solar radiation on a normal day, with approximately the same experimental condition as during

the eclipse, a series of measurements were also performed from 4<sup>th</sup> March to 14<sup>th</sup> March 2016. Unfortunately, the presence of clouds impacted these measurements. The incoming solar radiation showed dips and peaks e.g., as shown in Fig. 5 for the result of 12<sup>th</sup> March 2016. The heavy clouds in the early morning significantly reduced the solar radiation, subsequently the cloud appeared randomly. Similar or even irregular results were found in this first attempt.



**FIGURE 4.** The illumination observed during eclipse on 9<sup>th</sup> March 2016.



**FIGURE 5.** The illumination observed on 12<sup>th</sup> March 2016.

Due to this situation the second attempt was performed in the following year i.e., 2017. Solar radiation was measured from 4<sup>th</sup> March to 14<sup>th</sup> March 2017. A special attention was given to the measurement on 9<sup>th</sup> March 2017. Similar to the first attempt, the clouds obscured the solar radiation due to the season. The best result was obtained on 5<sup>th</sup> March 2017. On this day the cloud was relatively absent, the result is shown in Fig. 6.

Fig. 6 shows a pattern of solar radiation on a normal day. Although the pattern is not smooth, due to the atmospheric condition, this gives roughly a complete pattern. The illumination increases and reaches its maximum at around 09:00 local time.




FIGURE 6. The illumination observed on 5<sup>th</sup> March 2017.

The last attempt was performed on the third year i.e., from 4<sup>th</sup> March to 14<sup>th</sup> March 2018. Due to the rainy season the presence of clouds hindered the radiation. In this attempt the best result was measured on 8 March 2018, and it is shown in Fig. 7.




FIGURE 7. The illumination observed on 8<sup>th</sup> March 2018.

#### IV. DISCUSSIONS

Following Eq. (5), there are two effects that determined the total incoming solar radiation, the diurnal variation and the percentage of the eclipse. Both of them depend on the time. To show the effects of the eclipse, first we pay close attention to the pattern of solar radiation on a normal day e.g., showed in Fig. 6.

In the beginning the illumination is low, then it increases sharply and reaches its maximum at around 09:00 local time. This pattern is caused by the diurnal variation and the tilt sensor. The low illumination during the first part of the experiment is a diffuse radiation; this is due to the direct solar radiation that was blocked by a building. The following sharp increase of illumination indicates that the sensor was starting to receive the direct solar radiation. An increase of solar altitude will be accompanied by an increase in illumination. Based on Eq. (3), the illumination will increase and reach maximum at  $\alpha=\beta$ . On 5<sup>th</sup> March 2017, the sunrise and the sunset occur at 05:43 and 17:53 local time, respectively. From the day length we can calculate the sun's altitude at 09:00 local time is  $\alpha=24^{\circ}$ . The

sensor holder is tilted ( $\beta=25^{\circ}$ ) causing the sensor surface to be almost perpendicular to the sunrays at 09:00. The rest of the time the oblique sunrays cause less illumination. This similar situation is observed on Fig. 7 for the result of 8<sup>th</sup> March 2018. A difference of absolute illumination found on those figures is caused by the atmospheric condition.

The information of the sun such as position and other related data can be obtained from a free software e.g., Stellarium [18]. Using this software, we can follow the process of eclipse and collect the appropriate data. The collected eclipse obscuration of the 9<sup>th</sup> March 2016 eclipse is used to determine the relative visible sun's size using Eq. (4) and is presented in Fig. 8. In a normal condition the relative sun's size is one. During eclipse the visible sun's size decreases, reaching a minimum and then back to a normal size. The maximum eclipse obscuration observed in Yogyakarta was 80.5%, which gave the minimum relative visible sun's size of 0.195.





FIGURE 8. The relative visible sun's size obtained from the eclipse obscuration on 9<sup>th</sup> March 2016 using Stellarium (location: Yogyakarta).

The visible sun's size can be also determined using Eq. (5). The ratios of the solar radiation during the eclipse (Fig. 4) to the solar radiation on the normal day (Fig. 6 and 7) are shown in Fig. 9 and 10. In Fig. 9 the ratio (red line) is obtained from the illumination during the eclipse on 9<sup>th</sup> March 2016 and on the normal day (5<sup>th</sup> March 2017). In addition, the calculation using the Stellarium (black line) is presented for the comparison. Due to the atmospheric condition on those days, the ratio is not flawless. During the eclipse period (the shaded area) the ratio decreases, reaches a minimum and returns to the previous value. This ratio represents the fraction of the visible solar disk. Its pattern changes the illumination during the eclipse e.g., a clear dip in Fig. 4.


Fig. 10 shows the ratio (blue line) calculated from the illumination during the eclipse on 9<sup>th</sup> March 2016 and on the normal day (8<sup>th</sup> March 2018). It has the same behaviour as Fig. 9. It is clear that the ratio matched to the previous calculation using the eclipse obscuration previously recorded from the Stellarium. The discrepancies in the edge of Fig. 9 and Fig. 10 are caused by the atmospheric conditions and low solar radiation. In this part a small



fluctuation in solar radiation will give a large change in the ratio. Overall, the agreement of the experiment and the calculation is quite good.



**FIGURE 9.** The relative visible sun's size obtained from: (a) the ratio of solar radiation on 9<sup>th</sup> March 2016 to solar radiation on 5<sup>th</sup> March 2017 (red), and (b) the calculation using the Stellarium (black).



**FIGURE 10.** The relative visible sun's size obtained from: (a) the ratio of solar radiation on 9<sup>th</sup> March 2016 to solar radiation on 8<sup>th</sup> March 2018 (blue), and (b) the calculation using the Stellarium (black).

The experiment presented here is quite simple, as it directly shows the effect of the obscuration. The observation place can be chosen in a partially eclipse region, it is not necessary in the total eclipse path. The experiment result is a relative number that can be obtained without a calibrated instrument. Therefore, this experiment can be performed using any light sensor such as a solar cell. The overall constraint of this experiment is the weather conditions. The weather conditions during a normal day should be comparable as those of the eclipse.

## V. CONCLUSIONS

The solar radiation can be monitored using a simple light sensor that is available in most physics laboratories. Using

a simple formula, the visible sun's size can be determined from the solar radiation during an eclipse and a normal day. This experiment can be performed in the partially solar eclipse region.

## REFERENCES

- [1] Khavrus, V. and Shelevytsky, I., *Introduction to solar motion geometry on the basis of a simple model*, Phys. Educ. **45**, 641-53 (2010).
- [2] Khavrus, V. and Shelevytsky, I., *Geometry and the physics of seasons*, Phys. Educ. **47**, 680-92 (2012).
- [3] Zanetti, V. and Zecca, A., *A homemade pyranometer for solar experiments*, Am. J. Phys **51**, 633-6 (1983).
- [4] Russell, D. G. and Bartels, R. A., *The temperature of various surfaces exposed to solar radiation: An experiment*. Phys. Teach. **27**, 179-81 (1989).
- [5] Richards, S. F. and Smart, P., *Liberal Education from a Physics Project*, Phys. Teach. **11**, 463-8 (1973).
- [6] Stanford, Jr., A. L., *On Shadow bands accompanying total solar eclipses* Am. J. Phys **41**, 731-3 (1973).
- [7] Mottmann, J., *Solar eclipse predictions*, Am. J. Phys **48**, 626-8 (1980).
- [8] Sawicki, M., *Using the solar eclipse to estimate earth's distance from the moon*, Phys. Teach. **34**, 232-3 (1966).
- [9] Pasachoff, J. M. and Fraknoi, A., *Resource Letter OSE-1: Observing solar eclipse*, Am. J. Phys. **85**, 485-94 (2017).
- [10] Deliyannis, J., Papatheasoglou, D. and Stathopoulou, T. M., *On the variation of the direct solar radiation during the annular solar eclipse of April 29 1976*, Solar Physics **62**, 401-5 (1979).
- [11] Petkov, B., Tomasi C., Vitale V., Lanconelli C and Mazzola M., *Variation of incoming solar radiation flux during a partial eclipse episode: an improved model simulation*, Arxiv: 1110.2149 (2013).
- [12] Mollmann, K. P. and Vollmer, M., *Measurements and predictions of the illuminance during a solar eclipse*, Eur. J. Phys. **27**, 1299-1314 (2006).
- [13] Psiloglou, B. E. and Kambezidis, H. D., *Performance of the meteorological radiation model during the solar eclipse of 29 March 2006*, Atmos. Chem. Phys. **7**, 6047-6059 (2007).
- [14] Igoe, D. P. and Parisi, A. V., *Solar current output as a function of sun elevation: students as toolmakers*, Phys. Educ. **59**, 657-61 (2015).
- [15] Gueymard, C. A. and Myers, D. R., *Solar Radiation Measurement: Progress in Radiometry for Improved Modelling Modelling Solar Radiation at the Earth's Surface*, Ed. Viorel Badescu (Springer-Verlag, Berlin, 2008), pp. 1-5.
- [16] Hughes, D. W., *Brightness during a solar eclipse*, J. Br. Astron. Assoc. **110**, 4 203-5 (2000).
- [17] Vernier Software & Technology, 13979 SW Millikan Way, Beaverton, OR 97005-2886; <http://www.vernier.com>.
- [18] Stellarium.org.