




Pharmacy Education
*an international
peer review journal*



[Home](#) / [Archives](#) / Vol. 21 No. 2 (2021): IAI Conference 2020

Vol. 21 No. 2 (2021): IAI Conference 2020

We are pleased to confirm the publication of IAI Conference 2020.

Published: 2021-07-28

IAI CONFERENCE: The effect of loss-of-function allele (CYP2C19*3) with Clopidogrel efficacy in coronary heart disease patients

Ike Dhiah Rochmawati, Nur Hidayat, David Pomantow (Author)

p. 178-183



IAI CONFERENCE: The development of Origanum vulgare L. into nanoparticles in dosage forms

Lutfi Chabib, Arman Suryani, Muhammad Iqbal Pangestu, Adnan Muhammad Uno J Hidayat, A.M. Bagas Trianloka (Author)

p. 205-209



IAI CONFERENCE: The effect of ethanol extract from Portulaca oleracea on inhibiting total cholesterol on animal subjects

Afifah Bambang Sutjiatmo, Fahmy Ahsanul Haq, Sulaeman Al Jati, Suci Nar Vikasari (Author)

p. 22-26



IAI CONFERENCE: Analysis of white pepper essential oil components using gas chromatography-mass spectroscopy

Purwaniati, Gilang Eka Permana, Indro Pamudjo (Author)

p. 230-234

 PDF

IAI CONFERENCE: The evident gap between actual and perceived facilities supporting value-added pharmacy services

Andi Hermansyah, Catur Dian Setiawan, Firqin Fuad Riansyah (Author)

p. 48-51

 PDF

IAI CONFERENCE: Antiseptic gel formulated from ethanol extract of Citronella grass (*Cymbopogon nardus*) using CMC-Na, arabic gums, and gelatin as gelling agents

Reynelda Juliani Sagala, Lorensia Yolanda, Michael (Author)

p. 258-263

 PDF

IAI CONFERENCE: Dissolution profile of Curcumin from solid dispersion prepared at a high drug load of Curcumin using Poloxamer 407 as the carrier

Dewi Setyaningsih, Yustina Sri Hartini, Christine Patramurti, Sastira Putri, Yosi Bayu Murti (Author)

p. 77-80

 PDF

IAI CONFERENCE: The therapeutic outcomes and adverse drug reactions study of Clozapine on Schizophrenia inpatients in the Grhasia psychiatric hospital Yogyakarta, Indonesia

p. 287-295

 PDF

IAI CONFERENCE: The potential of banana fruit Ranggap (*Musa paradisiaca* var. *Troglodytarum*) as an excipient alternative to oral tablet dosage form

Dolih Gozalil, Iyan Sopyan, Resmi Mustarichi, Wahyu Priyo Legowo (Author)

p. 98-107



IAI CONFERENCE: Pandemic 2020: Economic pressure and evaluation of a primary health care innovation programme for type 2 diabetes mellitus treatment

Fransiskus Samuel Renaldi, Rani Sauriasari, Woro Riyadina (Author)

p. 132-142



IAI CONFERENCE: The influence of extracts and fractions from matoa leaves (Pometia pinnata) on angiotensin I levels

Ika Purwidyaningrum, Jason Merari Peranginangin, Iyem Sahira (Author)

p. 168-171



IAI CONFERENCE: IR spectroscopy coupled with chemometrics used as a simple and rapid method to determine the caffeine content of tea products

Lestyo Wulandari, Diana Hanifiyah Sutipno, Dwi Koko Pratoko (Author)

p. 195-200



IAI CONFERENCE: The impact of pharmacist shortage on the inventory management of medicines at primary healthcare centres in East Java, Indonesia

Abdul Rahem, Umi Athiyah, Catur Dian Setiawan, Andi Hermansyah (Author)

p. 8-14



IAI CONFERENCE: Activity test of fruit and pomegranate seeds (*Punica granatum* L) as a hepatoprotector against white male Wistar rats

p. 220-224



IAI CONFERENCE: The remuneration of the community pharmacist in the developing world: the case in Indonesia

Andi Hermansyah, Anila Impian Sukorini, Abdul Rahem (Author)

p. 36-41



IAI CONFERENCE: Satisfaction of drug information services implementation in Air Itam, Selindung, Girimaya, Gerunggang and Melintang health centre Pangkalpinang City

Rachmawati Felani Djuria (Author)

p. 246-250



IAI CONFERENCE: The effect of stress level on the therapeutic outcomes of type 2 diabetes mellitus at the regional public hospital of West Nusa Tenggara province

Baiq Leny Nopitasari, Baiq Nurbaety, Made Krisna Adi Jaya (Author)

p. 67-70



IAI CONFERENCE: Role of pharmacist in providing drug information and education for patients with chronic diseases during Transition of Care

Umi Athiyah, Abdul Rahem, Catur Dian Setiawan, Andi Hermansyah (Author)

p. 275-280



IAI CONFERENCE: Formulation and physical evaluation of facial cream preparations from Ceremai fruit juice (*Phyllanthus acidus* (L.) Skeels)

Danang Indriatmoko, Nani Suryani, Tarso Rudiana, Mila Kurniah (Author)

p. 87-92

**IAI CONFERENCE: Drug interactions in patients with hypertension at Persahabatan hospital in 2015**

Zainul Islam, Atika Vitasari, Muhammad Arif Ridwan (Author)

p. 309-314

**IAI CONFERENCE: Formulation and evaluation of Kirinyuh Leaf effervescent granules (*Chromolaena Odorata*. L) as an antioxidant**

Firman Gustaman, Keni Idacahyati, Winda Trisna Wulandari (Author)

p. 123-125

**IAI CONFERENCE: Self-medication and self-treatment with short-term antibiotics in Asian countries: A literature review**

p. 152-162

**IAI CONFERENCE: In vivo activity of *Phaseolus vulgaris* as an anti-hypercholesterolemic**

Keni Idacahyati, Yedy Purwandi Sukmawan, Nopi Yanti, Winda Trisna Wulandari, Firman Gustaman, Indra (Author)

p. 184-188

**IAI CONFERENCE: Therapeutic potential of *Cymbopogon schoenanthus* (L.) developed into nanoparticle technology**

Lutfi Chabib, Adnan Muhammad Uno J Hidayat, A.M. Bagas Trianloka, Muhammad Iqbal Pangestu, Arman Suryani, Yulianto (Author)

p. 210-214



IAI CONFERENCE: In vitro antimalarial activity assay of Ashitaba Leaf ethanolic extract (Angelica keiskei)

Alvi Kusuma Wardani, Abdul Rahman Wahid, Miftahul Jannah (Author)

p. 27-30



IAI CONFERENCE: Formulation and physical properties of lotion Kalakai root ethanol extract (Stenochlaena palustris Bedd)

Rabiatul Adawiyah, Riska Apriliyanti, Agustinawati Umaternate (Author)

p. 235-240



IAI CONFERENCE: Description of medication adherence in hypertensive respondents at Mandalika Mataram elderly social centre

Anna Pradiningsih, Dzun Haryadi Ittiqo, Neti Puput Arianti (Author)

p. 52-55



IAI CONFERENCE: Mother's knowledge and practices towards self-medication of fever among children under five years in Muncar Banyuwangi, Indonesia

Sinta Rachmawati, Khusnul Khotimah, Ika Norcahyanti (Author)

p. 264-268



IAI CONFERENCE: Hypoglycemic activity test on smooth pigweed (Ammaranthus Hybridus L) leaf water extract on male Wistar rats

Asti Yunia Rindarwati (Author)

p. 56-60



IAI CONFERENCE: Enhancing solubility and antibacterial activity using multi-component crystals of trimethoprim and malic acid

p. 296-304



IAI CONFERENCE: Correlation between the antioxidant capacity of plasma and blood glucose level

Eva Nurinda, Emelda, Nurul Kusumawardani (Author)

p. 108-115



IAI CONFERENCE: Acute toxicity study of the ethanolic extract of Eleutherine bulbosa Urb in Wistar rats

Helmina Wati, Rahmi Muthia, Kartini, Finna Setiawan (Author)

p. 143-147



IAI CONFERENCE: The effect of biofilm formation on the outcome therapy of diabetic foot infections (DFIs) patients in the outpatient clinic and inpatient ward of Dr Sardjito General Hospital Yogyakarta

Ika Puspitasari, Titik Nuryastuti, Rizka Humardewayanti Asdie, Hemi Sinorita, Nusaibah Umaroh⁴, Wahyu Tri Hapsari (Author)

p. 172-177



IAI CONFERENCE: Acute toxicity test of 96% ethanol extract of Syzygium myrtifolium leaves in white mice (Mus musculus)

Lusi Indriani, E. Mulyati Effendi, Kevin Christofer Fadillah (Author)

p. 201-204



IAI CONFERENCE: Adverse drug reactions associated with successful treatment of multidrug-resistant tuberculosis patients in Cempaka Putih Islamic Hospital Central Jakarta

Adin Hakim Kurniawan, Harpolia Cartika, Siti Aisyah (Author)

p. 15-21



IAI CONFERENCE: Comparing the quality of life of neuropathic patients treated with gabapentin and pregabalin at the neuropathic poly of the NTB provincial hospital in 2019

Nurul Qiyaam, Baiq Leny Nopitasari, Haerul Muhajiji (Author)

p. 225-229



IAI CONFERENCE: Exploring pharmacist experience and acceptance for debunking health misinformation in the social media

p. 42-47



IAI CONFERENCE: Medicine management in districts and primary health care centres (PHC) in the national health insurance (JKN) programme

Raharni, Rini Sasanti, Yuyun Yuniar (Author)

p. 251-257



IAI CONFERENCE: A pharmacoeconomic study: cost-utility analysis of modern wound dressings vs conventional wound dressings in patients with diabetic foot ulcer

Cyntiya Rahmawati, Baiq Leny Nopitasari (Author)

p. 71-76



IAI CONFERENCE: Proximate analysis on animal feed granules composed of raw material from fish innards wastes

Wahyuning Setyani, Christine Patramurti, Agatha Budi Susiana Lestari, Raysha Mcseer, Day Stella Maris Gewab, Maria Felix Zita Ina Bulu, Maria Regina Lusiana Kya (Author)

p. 281-286

**IAI CONFERENCE: The evaluation of compounding prescription and its availability of a licensed product for children at a private hospital in Yogyakarta, Indonesia**

p. 93-97

**IAI CONFERENCE: The potential effect of the green coffee extract on reducing atherogenic index in hyperlipidemic rats**

Fransiska Maria Christianty, Fifteen Aprila Fajrin, Andrean Roni (Author)

p. 126-131

**IAI CONFERENCE: Antibiotic use on paediatric inpatients in a public hospital in Bangil, Indonesia**

Ika Norcahyanti, Malikatur Rosyidah, Abdul Kadir Jaelani, Antonius N.W Pratama (Author)

p. 163-167

**IAI CONFERENCE: Analysis of pharmacists' knowledge and attitude in the pharmaceutical industry of halal certification and their readiness to produce halal medicine**

Abdul Rahem, Mustofa Helmi Effendi, Hayyun Durrotul Faridah (Author)

p. 1-7

**IAI CONFERENCE: Microencapsulation of Jeringau Rhizome essential oils (Acorus calamus L.) using β -Cyclodextrin**

Ledianasari, Deby Tristiyanti, Elva Maulydha Tanjung, Lovelyta Barani (Author)

p. 189-194

 PDF

IAI CONFERENCE: Effectiveness of combination of Moringa Leaf Extract (*Moringa oleifera* Lamk.) and Papaya Seed Extract (*Carica papaya* L.) in reducing blood sugar levels of diabetic rats

Novi Ayuwardani, Yetti Hariningsih (Author)

p. 215-219

 PDF

IAI CONFERENCE: Identification of herbal products used by families in the campus of Darussalam Gontor University

Amal Fadholah, Solikah Ana Istikomah, Cania Sofyan Islamanda, Evi Rohana Ma'rufi Jannah (Author)

p. 31-35

 PDF

IAI CONFERENCE: Effectiveness of public service advertisements on the use of antibiotics in Pangkalpinang

p. 241-245

 PDF

IAI CONFERENCE: Correlation between the level of knowledge of drug managers and drug management in several primary health centres in Malang regency

Ayuk Lawuningtyas Hariadini, Nur Ishmah, Hananditia Rachma Pramestutie (Author)

p. 61-66

 PDF

IAI CONFERENCE: The comparison of the actual cost to case-mix of type 2 diabetes mellitus inpatient in Pandan Arang Boyolali hospital

Sri Bintang Sahara Mahaputra Kusuma Negara, Devi Ristian Octavia, Primanitha Ria Utami (Author)

p. 269-274



IAI CONFERENCE: In silico screening of mint leaves compound (*Mentha piperita* L.) as a potential inhibitor of SARS-CoV-2

p. 81-86



IAI CONFERENCE: Sambiloto (*Andrographis paniculata* Nees.) leaf extract activity as an α -Amylase enzyme inhibitor

Yustina Sri Hartini, Dewi Setyaningsih, Maria Josephine Vivian Chang, Maria Cyrilla Iglesia Adi Nugrahanti (Author)

p. 305-308



IAI CONFERENCE: Fluconazole-tartaric acid co-crystal formation and its mechanical properties

Fikri Alatas, Nia Suwartiningsih, Hestiary Ratih, Titta Hartyana Sutarna (Author)

p. 116-122



IAI CONFERENCE: Evaluation of antidiarrheal effect of combination of Salam Leaves (*Syzygiumpolyanthum*) and Jackfruit Leaves (*Artocarpus heterophyllus* Lam.) infusum in rats induced by castor oil

Husnul Khuluq, Evi Marlina (Author)

p. 148-151



Open Journal Systems

PKP Publishing Services

Part of the
PKP Publishing Services Network



ISSN: 1477-2701

IAI CONFERENCE

RESEARCH ARTICLE

Dissolution profile of Curcumin from solid dispersion prepared at a high drug load of Curcumin using Poloxamer 407 as the carrier

Dewi Setyaningsih¹, Yustina Sri Hartini¹, Christine Patramurti¹, Sastira Putri¹, Yosi Bayu Murti²

¹Faculty of Pharmacy, Sanata Dharma University, Yogyakarta, Indonesia

²Faculty of Pharmacy, Universitas Gadjah Mada, Yogyakarta, Indonesia

Keywords

BCS II
Bioavailability
Curcuma longa
Poloxamers
Rotary evaporator

Correspondence

Dewi Setyaningsih
Faculty of Pharmacy
Sanata Dharma University
Yogyakarta
Indonesia
dewi@usd.ac.id

Abstract

Introduction: Curcumin, a BCS II drug, suffers from poor bioavailability. Increasing curcumin dissolution is a way to increase its bioavailability. Solid dispersion formulation can be used to improve curcumin dissolution. However, the successful curcumin solid dispersion is limited to a relatively low drug load (< 20%). **Aim:** This study aimed to investigate the dissolution behaviour of curcumin at a higher drug load (27.9%, 42.3%, and 56.6%) using a surfactant carrier of poloxamer 407. **Methods:** The solvent evaporation method was employed to prepare high drug load solid dispersion of curcumin. A physical mixture of the corresponding solid dispersion formulation was prepared as a control. Drug load, dissolution behaviour in 180 minutes, and dissolution efficiency (DE180) were determined. **Results:** The results showed that incorporating curcumin into a poloxamer 407 solid dispersion significantly improves the dissolution rate of curcumin. In the solid dispersion formula, the dissolution behaviour of curcumin was found to be carrier-dependent.

Introduction

Curcuminoid, a mixture compound of curcumin, demethoxycurcumin, and bisdemethoxycurcumin, is widely known as a functional food and is obtained from the isolation of turmeric (*Curcuma longa*) rhizome and other species of Zingiberaceae. Extensive investigations revealed the beneficial effects of curcumin to cure several diseases with its main activities as antioxidant and anti-inflammation (Basnet & Skalko-Basnet, 2011; Abdollahi *et al.*, 2017). Although extensive *in vitro* and *in vivo* studies highlighted the potential activities of curcumin, a clear therapeutic benefit of curcumin in clinical studies remained questionable due to the poor bioavailability of curcumin after oral administration (Gupta, Patchva & Aggarwal, 2012). The poor bioavailability of curcumin has been attributed to the low aqueous solubility and dissolution; the solubility of

curcumin in water was reported to be 11 ng/mL (Wang *et al.*, 1997).

An increased dissolution has been proposed to tackle the bioavailability problem of curcumin. Solid dispersion is one of the potential strategies to improve curcumin dissolution. Several factors enhance the dissolution rate of drugs, including increased surface area by reducing particle size, improvement of wetting, reduced agglomeration, and increased saturated concentration resulting from the amorphous form (Craig, 2002). Different types of carriers and methods are used to prepare solid dispersions (Leuner & Dressman, 2000). One of the carriers used in solid dispersion preparation is the poloxamer, a non-ionic triblock copolymer of polyoxyethylene-polyoxypropylene-polyoxyethylene, which has surfactant properties (Bodratti & Alexandridis, 2018). The dissolution enhancement rate of poloxamer

solid dispersion is achieved by improving the wettability and the ability of intermolecular interaction of poloxamer-drug to form a molecular dispersion (Ali, Williams & Rawlinson, 2010). This work aimed to investigate the dissolution behaviour of curcumin at a higher drug load (above 20% to less than 60%), using a surfactant carrier of poloxamer 407.

Material and Method

Material

Curcuma longa standardised extract (84.67% curcumin as detected by spectrophotometric method) was given by PT Phytochemindo Reksa, Bogor, Indonesia. Curcumin standard compound was purchased from Sigma. The poloxamer 407 was given by PT Konimex, Solo, Indonesia. Citric acid, methanol, ethanol, sodium dihydrogen phosphate dihydrate, and sodium lauryl sulfate (SLS) were purchased from Merck. Capsule shells of the size of 00 were purchased from Kapsulindo Nusantara, Indonesia. MiliQ water was supplied by our laboratory.

Method

1. Preparation of the solid dispersion formulation

The poloxamer 407 based solid dispersion (SD-PC) was prepared at 27.9%, 42.3%, and 56.6-wt % of the *C. longa* extract by a solvent evaporation method using a vacuum rotary evaporator. The formulas were coded as SD-PC 33, SD-PC 50, and SD-PC 67 assigned to each extract -wt. % of 27.9%, 42.3%, and 56.6-wt %, respectively in the solid dispersion formulations. Briefly, an accurate weight of *C. longa* extract and poloxamer 407 according to each code of formula was dissolved in ethanol at concentrations of 5 mg/mL and 45 mg/mL. Both solutions were stirred for 30 minutes to form a clear yellow solution. The ethanol was evaporated using a BUCHI vacuum rotary evaporator at 80°C. The obtained viscous solution was then subsequently dried using a vacuum oven. The dried product was pulverized in a mortar and sieved using a 60-mesh size. Physical mixture formulations (PM-PC) were prepared at the drug load as that prepared in the SD-PC formulations. The extract and poloxamer 407 were gently mixed in a mortar using a spatula and sieved using a 60-mesh size. For the dissolution study, the SD powder was prepared into an approximately 500 mg capsule of size 00. The drug load of curcumin in the SD formulations was quantified using the calibration curve prepared from the serial concentrations of curcumin reference standard in methanolic solution.

The linearity between the concentration and the absorption was confirmed with $y=0.1349x + 0.0035$ and a correlation coefficient of 0.9972.

2. Dissolution

The dissolution of the SD-PC or PM-PC prepared in capsules was evaluated using a SOTAX AT7 dissolution tester according to the USP type II dissolution method. For the dissolution study of lipophilic compounds like curcumin, 0.5% surfactant of SLS was added into 20 mM sodium phosphate buffer of pH 6.0. This pH value was chosen because the curcumin solution is highly stable at pH 6.0 (Wang *et al.*, 1997). The dissolution was carried out in 900 mL media with a paddle speed of 100 rpm and a temperature of $37 \pm 0.5^\circ\text{C}$. The samples of 2 mL were taken at predetermined time intervals of 0, 10, 15, 30, 45, 60, 90, 120, and 180 minutes. The withdrawn volume was replaced with the same volume of a fresh dissolution medium, maintained at 37° to keep the volume and sink condition constant. Curcumin in the dissolution sample was determined using the validated spectroscopic method at 430 nm against the blank sample of the dissolution medium. Curcumin concentrations were quantified based on the calibration curve at which the result shows linearity with a linear equation of $y=0.1279x + 0.009$ and a relation coefficient of 0.997.

3. Data analysis

The dissolution profile was determined by calculating the Dissolution Efficiency (DE) at 180 min, as in equations 1 and 2. The area under the curve of dissolved curcumin was calculated by the trapezoidal method, as shown in the following equations:

$$DE_t = \left(\frac{\int_{t_1}^{t_2} y \cdot dt}{y_{100} \cdot t} \right) \times 100\% \quad \text{Equation 1} \quad \text{Equation 2} \quad AUC = \sum_{i=1}^{i=n} \frac{(t_i - t_{i-1})(y_{i-1} + y_i)}{2}$$

Equation 1

Equation 2

DE_t : Dissolution efficiency at time (t)

y : Area under the curve of dissolved drug at time t

$y_{100.t}$: Rectangle area where 100% of drug dissolved at time t

Results

Table I shows the amount of curcumin content in the solid dispersion and physical mixture formulations. The % assay values, calculated based on the recovery at which the obtained curcumin contents were divided by the theoretical values and multiplied by 100%, were

between 94.33-99.48% for SD-PC and PM-PC formulations. The curcumin content for all SD and PM formulations was calculated for the relative standard

deviation (RSD) values. The RSD values for all SD and PM formulations are between 0.12%-4.50%.

Table I: Curcumin content was observed in the solid dispersion (SD) and physical mixture

% weight of <i>C. longa</i> extract	Code	SD-PC			Code	PM-PC		
		% found	% assay	RSD		% found	% assay	RSD
27.9	SD-PC 33	27.59 ± 0.38	99.24	1.37	PM-PC 33	27.42 ± 0.32	99.48	1.16
42.3	SD-PC 50	40.87 ± 0.67	96.54	1.63	PM-PC 50	41.22 ± 0.71	96.38	1.72
56.6	SD-PC 67	56.29 ± 0.07	99.38	0.12	PM-PC 67	53.52 ± 2.41	94.33	4.50

Note: Data were presented as mean and SD of 3 assays. % assay was calculated by dividing the observed amount of curcumin by the theoretical amount of curcumin present in the formulation at the accordingly wt % of *C. longa* extract. The theoretical amount of curcumin was determined based on the curcumin content in the standardized *C. longa* extract.

The *in vitro* dissolution profile of the different solid dispersion formulations using poloxamer 407 and the respective physical mixture in 20 mM sodium phosphate buffer of pH 6.0 are shown in Figure 1 (a, b, and c). At the end of 180 minutes, 26.62%, 18.38%, 12.22%, 41.18%, 23.41%, and 21.92% were released from PM-PC 33, PM-PC 50, PM PC 67, SD-PC 33, SD-PC 50, and SD PC 67, respectively. The dissolution profile of SD and PM formulation show stagnant dissolution rate after 30 minutes (SD/PM-PC 33), 45 minutes (SD/PC-50), and 60 (SD/PC-PC 67).

DE₁₈₀ was used to compare the dissolution profiles resulting from all SD and PC formulations and presented in Figure 1d, showing that all SD formulations increased the dissolution rate of curcumin compared to the physical mixture formulation as monitored in 180 minutes. It also revealed that the solid dispersion formulation increased the dissolution rate of curcumin as compared to the corresponding physical mixture formulation.




Figure 1: Dissolution behaviour of curcumin from SD and PM formulations

Discussion

As shown in Table I, the RSD values for SD and PM formulations are all below 5%. The data indicate that the solid dispersion and physical mixtures formulations are homogenous; independent assays of three samples taken from any dispersion showed an even drug distribution within each system. All formulas showed that % values between the founded curcumin content and the theoretical curcumin concentration were close, as reflected by a high % assay values of between 90 and 100%. Moreover, curcumin content in the SD-PC is quite close to the PM-PC, indicating that the curcumin remains preserved during the preparation processes of the solid dispersion formulations.

Surfactants can improve the hydrophobicity of poorly water-soluble drugs and increase the dissolution rate by increasing wettability. Solid dispersion formulation of *C. longa* extract using poloxamer 407 as carrier improves the dissolution profile of curcumin compared to the physical mixture formulation (Figure 1a, b, c) due to the improvement of wetting with the addition of poloxamer 407. A significant increase in dissolution rate was observed from the SD-PC 33 and SD-PC 67 compared to the corresponding physical mixture formulation (PM-PC 33 and PM-PC 67). The dissolution rate declined as higher extract concentration in the solid dispersion formulations was used (SD-PC 67). The release of curcumin during 180 minutes was about 40%; however, at higher extract concentration (SD-PC 67), the dissolution decreased to 20%. A lower dissolution rate at a higher drug load was also reported by other publications, suggesting that the dissolution mechanism is controlled by the carrier.

Dissolution efficiency (DE) is a dissolution parameter used to compare characteristics of the dissolution profile between formulas. The effect of increasing extract content in the solid dispersion formulation markedly reduced the dissolution rate of curcumin, as indicated by the DE values (Figure 1d). Lowering the dissolution rate by increasing the extracted content was observed in the solid dispersion formulation and demonstrated by the physical mixture formulation.

Conclusion

Poloxamer 407 is an effective carrier to increase the dissolution rate of curcumin during a 180-minute study in solid dispersion formulations containing *C. longa* standardized extract. An increased extracted content in the solid dispersion formulation results in reduced dissolution of curcumin. The highest dissolution rate of curcumin is found in the extracted

content of 27.9 -wt% of curcumin of the SD-PC 33 formulation.

Acknowledgements

The authors would like to thank Ministry of Research, Technology, and Higher Education of Republic Indonesia for providing the grant in 2018 under the scheme of International Research Collaboration.

References

- Abdollahi, E., Momtazi, A. A., Johnston, T. P., & Sahebkar, A. (2017). Therapeutic effects of curcumin in inflammatory and immune-mediated diseases: A nature-made jack-of-all-trades? *Journal of Cellular Physiology*, <https://doi.org/10.1002/jcp.25778>
- Ali, W., Williams, A. C., & Rawlinson, C. F. (2010). Stoichiometrically governed molecular interactions in drug: Poloxamer solid dispersions. *International Journal of Pharmaceutics*, **391**(1-2), 162-168. <https://doi.org/10.1016/j.ijpharm.2010.03.014>
- Basnet, P., & Skalko-Basnet, N. (2011). Curcumin: an anti-inflammatory molecule from a curry spice on the path to cancer treatment. *Molecules*, **16**(6), 4567-4598
- Bodratti, A. M., & Alexandridis, P. (2018). Formulation of Poloxamers for Drug Delivery. *Journal of Functional Biomaterials*, **9**(1), 11
- Craig, D. Q. (2002). The mechanisms of drug release from solid dispersions in water-soluble polymers. *International Journal of Pharmaceutics*, **231**(2), 131-144. [https://doi.org/10.1016/s0378-5173\(01\)00891-2](https://doi.org/10.1016/s0378-5173(01)00891-2)
- Gupta, S. C., Patchva, S., & Aggarwal, B. B. (2012). Therapeutic Roles of Curcumin: Lessons Learned from Clinical Trials. *The AAPS Journal*, **15**(1), 195-218
- Leuner, C., & Dressman, J. (2000). Improving drug solubility for oral delivery using solid dispersions. *European Journal of Pharmaceutics and Biopharmaceutics*, **50**(1), 47-60
- Wang, Y. J., Pan, M. H., Cheng, A. L., Lin, L. I., Ho, Y. S., Hsieh, C. Y., & Lin, J. K. (1997). Stability of curcumin in buffer solutions and characterization of its degradation products. *Journal of Pharmaceutical and Biomedical Analysis*, **15**(12), 1867-1876