

International Journal of Applied Sciences and Smart Technologies

Volume 04, Issue 02, December 2022

Sugarcane Production Modeling Using Machine Learning in Western Maharashtra
Chhaya Narvekar, Madhuri Rao

A White Noise Approach to Occupation Times of Brownian Motion
Herry Pribawanto Suryawan

Dijkstra Algorithm Implementation in Determining the Shortest Route of Industrial Gas Distribution in PT Tira Austenite Tbk Cikarang with Python Programming Language
Bernadetha Nathalia Fitra Soverenty, Cyrenia Novella Krisnamurti

An Enhanced Multi-Level Authentication Electronic Voting System
Ayodeji O.J Ibitoye, Halleluyah O. Aworinde, Esther T. Adekunle

**Appropriate Technology-Based Project-Based Learning:
3D Printing Utilization for Learning Media Class**
Monica Cahyaning Ratri

Employee Presence Using Body Temperature Detection and Face Recognition
Arif Ainur Rafiq, Erna Alimudin, Della Puspa Rani

**The Experiment of Wind Electric Water Pumping for Salt Farmers
in Remote Area of Demak-Indonesia**
S. Dio Zevalukito, Y.B. Lukiyanto, F. Risky Prayogo

Response Surface Modelling of the Mechanical Properties of Oil Palm Empty Fruit Bunch Fibre Reinforced Polyester Composites
Chinwe Evangeline Kamma

Comparative Study of Master Oil (MO) and Lophira Lanceolata (Ochnaceae) Oil (LLO) Lubricants in Sewing Machines
J. B. Yerima, G. O. Mokoro, D. William, S. D. Najoji

Design of Someone's Character Identification Based on Handwriting Patterns Using Support Vector Machine
Rosalia Arum Kumalasanti

Optimization of Vacuum Forming Parameter Settings to Minimize Burning Defect on Strawberry Packaging Products Using the Taguchi Method
Th. Adi Nugroho, Adhi Setya Hutama, Perwita Kurniawan

The Effect of the Number of Cooling Pads on the Output Air Condition and Effectiveness of Air Cooler
Wibowo Kusbandono

Pre-service Science Teachers' Competence and Confidence in Scientific Inquiry
Rohandi

p-ISSN 2655-8564 & e-ISSN 2685-9432

CONTENTS

| | |
|---|---------|
| CONTENTS | i |
| EDITORIAL BOARD | Iii |
| PREFACE | iv |
| Sugarcane Production Modeling Using Machine Learning in Western Maharashtra <i>Chhaya Narvekar and Dr.Madhuri Rao</i> | 123–130 |
| A White Noise Approach to Occupation Times of Brownian Motion <i>Herry Pribawanto Suryawan</i> | 131–140 |
| Dijkstra Algorithm Implementation in Determining the Shortest Route of Industrial Gas Distribution in PT Tira Austenite Tbk Cikarang with Python Programming Language <i>Bernadetha Nathalia Fitra Soverenty, Cyrenia Novella Krisnamurti</i> | 141–148 |
| An Enhanced Multi-Level Authentication Electronic Voting System <i>Ayodeji O.J Ibitoye, Halleluyah O. Aworinde, Esther T. Adekunle</i> | 149–158 |
| Appropriate Technology-Based Project-Based Learning: 3D Printing Utilization for Learning Media Class <i>Monica Cahyaning Ratri</i> | 159–172 |
| Employee Presence Using Body Temperature Detection and Face Recognition <i>Arif Ainur Rafiq, Erna Alimudin, and Della Puspa Rani</i> | 173–184 |
| The Experiment of Wind Electric Water Pumping for Salt Farmers in Remote Area of Demak-Indonesia <i>S. Dio Zevalukito, YB. Lukiyanto and F. Risky Prayogo</i> | 185–194 |
| Response Surface Modelling of the Mechanical Properties of Oil Palm Empty Fruit Bunch Fibre Reinforced Polyester Composites <i>Chinwe Evangeline Kamma</i> | 195–220 |
| Comparative Study of Master Oil (MO) and Lophira Lanceolata (Ochnaceae) Oil (LLO) Lubricants in Sewing Machines <i>J. B Yerima, G. O Mokoro, D. William, and S. D Najoji</i> | 221-232 |
| Design of Someone's Character Identification Based on Handwriting Patterns Using Support Vector Machine <i>R. A. Kumalasanti</i> | 233-240 |
| Optimization of Vacuum Forming Parameter Settings to Minimize Burning Defect on Strawberry Packaging Products Using the Taguchi Method <i>Th. Adi Nugroho, Adhi Setya Hutama and Perwita Kurniawan</i> | 241-254 |

| | |
|---|---------|
| The Effect of the Number of Cooling Pads on the Output Air Condition and Effectiveness of Air Cooler <i>Wibowo Kusbandono</i> | 255-266 |
| Pre-service Science Teachers' Competence and Confidence in Scientific Inquiry <i>Rohandi</i> | 267-280 |
| AUTHOR GUIDELINES | 281-282 |

EDITORIAL BOARD

Editor in Chief

Dr. I Made Wicaksana Ekaputra (*Sanata Dharma University, Yogyakarta, Indonesia*)
Email: made@usd.ac.id

Associate Editor

Dr. Pham Nhu Viet Ha (*Vietnam Atomic Energy Institute, Hanoi, Vietnam*)
Dr. Hendra Gunawan Harno (*Gyeongsang National University, Jinju, The Republic of Korea*)
Dr. Iswanjono (*Sanata Dharma University, Yogyakarta, Indonesia*)
Dr. Mukesh Jewariya (*National Physical Laboratory, New Delhi, India*)
Dr. Mongkolsery Lin (*Institute of Technology of Cambodia, Phnom Penh, Cambodia*)
Dr. Yohanes Baptista Lukiyanto (*Sanata Dharma University, Yogyakarta, Indonesia*)
Dr. Apichate Maneewong (*Thailand Institute of Nuclear Technology, Bangkok, Thailand*)
Prof. Dr. Sudi Mungkasi (*Sanata Dharma University, Yogyakarta, Indonesia*)
Dr. Pranowo (*Universitas Atma Jaya Yogyakarta, Yogyakarta, Indonesia*)
Dr. Monica Cahyaning Ratri (*Sanata Dharma University, Yogyakarta, Indonesia*)
Dr. Mahardhika Pratama (*Nanyang Technological University, Singapore*)
Dr. Augustinus Bayu Primawan (*Sanata Dharma University, Yogyakarta, Indonesia*)
Prof. Dr. Leo Hari Wiryanto (*Bandung Institute of Technology, Bandung, Indonesia*)

Editorial Proofreader

Ir. Ignatius Aris Dwiatmoko, M.Sc. (*Sanata Dharma University, Yogyakarta, Indonesia*)
P. H. Prima Rosa, S.Si., M.Sc. (*Sanata Dharma University, Yogyakarta, Indonesia*)

Editorial Assistant

Eduardus Hardika Sandy Atmaja, M.Cs. (*Sanata Dharma University, Yogyakarta, Indonesia*)
Rosalia Arum Kumalasanti, M.T. (*Sanata Dharma University, Yogyakarta, Indonesia*)
Vitalis Ayu, M.Cs. (*Sanata Dharma University, Yogyakarta, Indonesia*)

Administration

Catharina Maria Sri Wijayanti, S.Pd. (*Sanata Dharma University, Yogyakarta, Indonesia*)

Contact us

International Journal of Applied Sciences and Smart Technologies
Faculty of Science and Technology
Sanata Dharma University
Kampus III Paingan, Maguwoharjo, Depok, Sleman
Yogyakarta, 55282
Phone : +62 274883037 ext. 523110, 52320
Fax : +62 272886529
Email : editorial.ijasst@usd.ac.id
Website : <http://e-journal.usd.ac.id/index.php/IJASST>

International Journal of Applied Sciences and Smart Technologies

Volume 4, Issue 2, page iii

p-ISSN 2655-8564, e-ISSN 2685-9432

IJASST is an open-access peer-reviewed journal that mediates the dissemination of research and studies conducted by academicians, researchers, and practitioners in science, engineering, and technology.

PREFACE

Dear readers, we are delighted to serve you Volume 4, Issue 2 of *International Journal of Applied Sciences and Smart Technologies* (IJASST), which is managed and published by the Faculty of Science and Technology, Sanata Dharma University. IJASST is an open-access peer-reviewed journal that mediates the dissemination of research and studies conducted by academicians, researchers, and practitioners in science, engineering, and technology. Its scope also includes basic sciences which relate to technology, such as applied mathematics, physics, and chemistry.

In this edition, we have thirteen papers authored by researchers from Indonesia, India, Turkey, Nigeria, and Palestine. Submitted papers are reviewed fairly using the open journal system (OJS) of IJASST. After the review process, accepted papers of the journal are publicly available for free at the website of IJASST.

For future issues, we are looking forward to your contributions to IJASST.

Dr. I Made Wicaksana Ekaputra
Editor in Chief
IJASST

A White Noise Approach to Occupation Times of Brownian Motion

Herry Pribawanto Suryawan

Department of Mathematics, Faculty of Science and Technology

Sanata Dharma University, Yogyakarta, Indonesia

Corresponding Author: herrypribs@usd.ac.id

(Received 02-11-2022; Revised 17-11-2022; Accepted 18-11-2022)

Abstract

Occupation times of a stochastic process models the amount of time the process spends inside a spatial interval during a certain finite time horizon. It appears in the fiber lay-down process in nonwoven production industry. The occupation time can be interpreted as the mass of fiber material deposited inside some region. From application point of view, it is important to know the average mass per unit area of the final fleece. In this paper we use white noise theory to prove the existence of the occupation times of one-dimensional Brownian motion and provide an expression for the expected value of the occupation times.

Keywords: occupation times, Brownian motion, white noise analysis

1 Introduction

Technical textiles have attracted great attention to diverse branches of industry over the last decades due to their comparatively cheap manufacturing. By overlapping thousands of individual slender fibers, random fiber webs emerge yielding nonwoven materials that find applications e.g. in textile, building and hygiene industry as integral components of baby diapers, closing textiles, filters and medical devices, to name but a few. They are

produced in melt-spinning operations: hundreds of individual endless fibers are obtained by the continuous extrusion of a molten polymer through narrow nozzles that are densely and equidistantly placed in a row at a spinning beam. The viscous or viscoelastic fibers are stretched and spun until they solidify due to cooling air streams. Before the elastic fibers lay down on a moving conveyor belt to form a web, they become entangled and form loops due to the highly turbulent air flows. The homogeneity and load capacity of the fiber web are the most important textile properties for quality assesment of industrial nonwoven fabrics. The optimization and control of the fleece quality require modeling and simulation of fiber dynamics and lay-down. Available data to judge the quality, at least on the industrial scale, are usually the mass per unit area of the fleece.

Since the mathematical treatment of the whole process at a stroke is not possible due to its complexity, a hierarchy of models that adequately describe partial aspects of the process chain has been developed in research during the last years. A stochastic model for the fiber deposition in the nonwoven production was proposed and analyzed in [4, 5, 7, 10]. The model is based on stochastic differential equations describing the resulting position of the fiber on the belt under the influence of turbulent air flows. In [1] parameter estimation of the Ornstein-Uhlenbeck process from available mass per unit area data, the occupation time in mathematical terms, was done.

Definition 1.1. Let $X = (X_t)_{t \in [0, T]}$, $T > 0$, be a stochastic process and consider an interval $[a, b] \subset \mathbb{R}$. The occupation time $M_{T, [a, b]}(X)$ is defined as

$$M_{T, [a, b]}(X) := \int_0^T \mathbf{1}_{[a, b]}(X_t) dt = \int_0^T \int_a^b \delta_0(X_t - x) dx dt.$$

Here, $\mathbf{1}_{[a, b]}$ denoted the indicator function of the interval $[a, b]$ and δ_0 is the Dirac-delta distribution.

Formally, occupation times models the time the stochastic process spends inside the spatial interval $[a, b]$ during the time interval $[0, T]$. In terms of our physical model for the nonwoven production, the occupation time can be interpreted as the mass of fiber material deposited inside the interval $[a, b]$, i.e. the mass per unit area of the final fleece.

Motivated by the above mentioned problem, in this paper we study the occupation time of one-dimensional Brownian motion. In particular, we show that occupation times of one-dimensional Brownian motion is a white noise distribution in the sense of Hida.

Although it is possible to study the problem by classical probabilistic method, we use a white noise approach to generalize the concept also to higher dimensions in later research. Moreover, in future work an extension to more general process (e.g. with fractional Gaussian noise) is planned. In the next section we provide necessary background on the white noise theory. The main result together with its proof are given afterward.

2 White Noise Analysis

In this section we give background on the white noise theory used throughout this paper. For a more comprehensive discussions including various applications of white noise theory we refer to [8, 9, 12, 13] and references therein. We start with the Gelfand triple

$$\mathcal{S}(\mathbb{R}) \hookrightarrow L^2(\mathbb{R}) \hookrightarrow \mathcal{S}'(\mathbb{R}),$$

where $\mathcal{S}(\mathbb{R})$ is the space of real-valued Schwartz test function, $\mathcal{S}'(\mathbb{R})$ is the space of real-valued tempered distributions, and $L^2(\mathbb{R})$ is the real Hilbert space of all real-valued Lebesgue square-integrable functions. Next, we construct a probability space $(\mathcal{S}'(\mathbb{R}), \mathcal{C}, \mu)$ where \mathcal{C} is the Borel σ -algebra generated by cylinder sets on $\mathcal{S}'(\mathbb{R})$ and the unique probability measure μ is established through the Bochner-Minlos theorem by fixing the characteristic function

$$C(f) := \int_{\mathcal{S}'(\mathbb{R})} \exp(i\langle \omega, f \rangle) d\mu(\omega) = \exp\left(-\frac{1}{2}\|f\|_0^2\right)$$

for all $f \in \mathcal{S}(\mathbb{R})$. Here $\|\cdot\|_0$ denotes the usual norm in the $L^2(\mathbb{R})$, and $\langle \cdot, \cdot \rangle$ denotes the dual pairing between $\mathcal{S}'(\mathbb{R})$ and $\mathcal{S}(\mathbb{R})$. The dual pairing is considered as the bilinear extension of the inner product on $L^2(\mathbb{R})$, i.e.

$$\langle g, f \rangle = \int_{\mathbb{R}} g(x)f(x) dx,$$

for all $g \in L^2(\mathbb{R})$ and $f \in \mathcal{S}(\mathbb{R})$. This probability space is known as the real-valued white noise space since it contains the sample paths of the one-dimensional Gaussian white noise. In this setting a one-dimensional Brownian motion can be represented by a continuous modification of the stochastic process $B = (B_t)_{t \geq 0}$ with

$$B(t) := \langle \cdot, \mathbf{1}_{[0,t]} \rangle,$$

where $\mathbf{1}_A$ denotes the indicator function of a set $A \subset \mathbb{R}$.

In the sequel we will use the Gel'fand triple

$$(\mathcal{S}) \hookrightarrow L^2(\mu) := L^2(\mathcal{S}'(\mathbb{R}), \mathcal{C}, \mu) \hookrightarrow (\mathcal{S})^*$$

where (\mathcal{S}) is the space of white noise test functions obtained by taking the intersection of a family of Hilbert subspaces of $L^2(\mu)$. The space of white noise distributions $(\mathcal{S})^*$ is defined as the topological dual space of (\mathcal{S}) . Elements of (\mathcal{S}) and $(\mathcal{S})^*$ are known as *Hida test functions* and *Hida distributions*, respectively. Within this framework white noise can be considered as the time derivative of Brownian motion with respect to the topology of $(\mathcal{S})^*$. An important tool in white noise analysis is the S-transform which can be considered as the Laplace transform with respect to the infinite dimensional Gaussian measure. The S-transform of $\Phi \in (\mathcal{S})^*$ is defined as

$$(S\Phi)(\varphi) := \langle \langle \Phi, : \exp(\langle \cdot, \varphi \rangle) : \rangle \rangle, \quad \varphi \in \mathcal{S}(\mathbb{R}),$$

where

$$: \exp(\langle \cdot, \varphi \rangle) := C(\varphi) \exp(\langle \cdot, \varphi \rangle),$$

is the so-called Wick exponential and $\langle \langle \cdot, \cdot \rangle \rangle$ denotes the dual pairing between $(\mathcal{S})^*$ and (\mathcal{S}) . We define this dual pairing as the bilinear extension of the sesquilinear inner product on $L^2(\mu)$. The S-transform provides a convenient way to identify a Hida distribution $\Phi \in (\mathcal{S})^*$, in particular, when it is hard to find the explicit form for the Wiener-Itô chaos decomposition of Φ .

Theorem 2.1. [12]

1. The S-transform is injective, i.e. if $S\Phi(\varphi) = S\Psi(\varphi)$ for all $\varphi \in \mathcal{S}(\mathbb{R})$, then $\Phi = \Psi$.
2. If a stochastic distribution process $X : I \rightarrow (\mathcal{S})^*$ is differentiable, then $S \frac{d}{dt} X(t)(\varphi) = \frac{d}{dt} SX(t)(\varphi)$ for all $\varphi \in \mathcal{S}(\mathbb{R})$.

In the following we state a sufficient condition on the Bochner integrability of a family of Hida distributions which depend on an additional parameter.

Theorem 2.2. [11] Let $(\Omega, \mathcal{A}, \nu)$ be a measure space and $\lambda \mapsto \Phi_\lambda$ be a mapping from Ω to $(\mathcal{S})^*$. If

- (1) the mapping $\lambda \mapsto S(\Phi_\lambda)(\varphi)$ is measurable for all $\varphi \in \mathcal{S}(\mathbb{R})$, and
- (2) there exist $C_1(\lambda) \in L^1(\Omega, \mathcal{A}, \nu)$, $C_2(\lambda) \in L^\infty(\Omega, \mathcal{A}, \nu)$ and a continuous seminorm $\|\cdot\|$ on $\mathcal{S}(\mathbb{R})$ such that for all $z \in \mathbb{C}$, $\varphi \in \mathcal{S}(\mathbb{R})$

$$|S(\Phi_\lambda)(z\varphi)| \leq C_1(\lambda) \exp(C_2(\lambda)|z|^2 \|\varphi\|^2),$$

then Φ_λ is Bochner integrable with respect to some Hilbertian norm which topologizing $(\mathcal{S})^*$. Hence $\int_\Omega \Phi_\lambda d\nu(\lambda) \in (\mathcal{S})^*$, and furthermore

$$S\left(\int_\Omega \Phi_\lambda d\nu(\lambda)\right) = \int_\Omega S(\Phi_\lambda) d\nu(\lambda).$$

Let $0 < T < \infty$ and $B = (B_t)_{t \in [0, T]}$ be a one-dimensional standard Brownian motion. The corresponding Donsker's delta distribution is given by

$$\delta_0(B_t - x) = \delta_0(\langle \cdot, \mathbf{1}_{[0, t]} \rangle - x) = \frac{1}{2\pi} \int_{\mathbb{R}} \exp(i\lambda(\langle \cdot, \mathbf{1}_{[0, t]} \rangle - x)) d\lambda.$$

It has been proved that $\delta_0(B_t - x) \in (\mathcal{S})^*$. Furthermore, its S-transform is given by

$$S\delta_0(B_t - x)(\varphi) = \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{1}{2t}(\langle \varphi, \mathbf{1}_{[0, t]} \rangle - x)^2\right),$$

for any $\varphi \in \mathcal{S}(\mathbb{R})$. For details and proofs see e.g. [8, 12]. The Donsker delta distribution is an important research object in the Gaussian analysis. For example, it can be used to study local times, self-intersection local times, stochastic current and Feynman integrals, see e.g. [2, 3, 6, 14, 15, 18]. The derivatives of Donsker's delta distribution has been also studied in [16]. In [17] Donsker's delta distribution is analyzed in the context of stochastic processes with memory.

3 Main Result

Now we are ready to prove the main finding of the paper.

Theorem 3.1. *Let $B = (B_t)_{t \in [0, T]}$ be a one-dimensional standard Brownian motion and consider an interval $[a, b] \subset \mathbb{R}$. The occupation time*

$$M_{T, [a, b]}(B) := \int_0^T \int_a^b \delta_0(B_t - x) dx dt$$

is a Hida distribution.

Proof: It is apparent, at least formally, that occupation times can be obtained by integrating Donsker’s delta distribution with respect to the product measure on $[a, b] \times [0, T]$. In this regard we will use Kondratiev-Streit integration theorem (Theorem 2.2) to prove the statement. Observe that for any $\varphi \in \mathcal{S}(\mathbb{R})$

$$S\delta_0(B_t - x)(\varphi) = \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{1}{2t} \left(\int_0^t \varphi(s) ds - x\right)^2\right)$$

is a measurable function with respect to the product measure on $[0, T] \times [a, b]$. Now for any $z \in \mathbb{C}$ and $\varphi \in \mathcal{S}(\mathbb{R})$ we have

$$\begin{aligned} & |S\delta_0(B_t - x)(z\varphi)| \\ &= \left| \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{1}{2t} \left(\int_0^t z\varphi(s) ds - x\right)^2\right) \right| \\ &= \frac{1}{\sqrt{2\pi t}} \left| \exp\left(-\frac{1}{2t} (\langle z\varphi, \mathbf{1}_{[0,t]} \rangle - x)^2\right) \right| \\ &= \frac{1}{\sqrt{2\pi t}} \left| \exp\left(-\frac{1}{2t} (\langle z\varphi, \mathbf{1}_{[0,t]} \rangle^2 - 2x \langle z\varphi, \mathbf{1}_{[0,t]} \rangle + x^2)\right) \right| \\ &= \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{x^2}{2t}\right) \left| \exp\left(-\frac{1}{2t} \langle z\varphi, \mathbf{1}_{[0,t]} \rangle^2\right) \exp\left(\frac{x}{t} \langle z\varphi, \mathbf{1}_{[0,t]} \rangle\right) \right| \\ &\leq \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{x^2}{2t}\right) \exp\left(\frac{1}{2t} |\langle z\varphi, \mathbf{1}_{[0,t]} \rangle|^2\right) \exp\left(\left|\frac{x}{t} \langle z\varphi, \mathbf{1}_{[0,t]} \rangle\right|\right) \\ &= \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{x^2}{2t}\right) \exp\left(\frac{1}{2t} |z|^2 |\langle \varphi, \mathbf{1}_{[0,t]} \rangle|^2\right) \exp\left(\frac{|x|}{\sqrt{t}} \frac{|z|}{\sqrt{t}} |\langle \varphi, \mathbf{1}_{[0,t]} \rangle|\right) \\ &\leq \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{x^2}{2t}\right) \exp\left(\frac{1}{2t} |z|^2 |\varphi|^2 |\mathbf{1}_{[0,t]}|^2\right) \exp\left(\frac{x^2}{2t}\right) \exp\left(\frac{1}{2t} |z|^2 |\varphi|^2 |\mathbf{1}_{[0,t]}|^2\right) \\ &= \frac{1}{\sqrt{2\pi t}} \exp(|z|^2 |\varphi|^2). \end{aligned}$$

The first factor

$$C_1(t, x) = \frac{1}{\sqrt{2\pi t}}$$

is an integrable function on $[0, T] \times [a, b]$ while the second factor

$$C_2(t, x) = \exp(|z|^2 |\varphi|^2)$$

is a bounded function of t and x . Theorem 2.2 gives the desired result. ■

Corollary 3.2. *The S-transform of the occupation times of Brownian motion is given by*

$$SM_{T,[a,b]}(B)(\varphi) = \int_0^T \frac{1}{\sqrt{2\pi t}} \int_a^b \exp\left(-\frac{1}{2t} \left(\int_0^t \varphi(s) ds - x\right)^2\right) dx dt,$$

for any $\varphi \in \mathcal{S}(\mathbb{R})$.

Proof: Since $M_{T,[a,b]}(B) \in (\mathcal{S})^*$ then according to Theorem 2.2 for every $\varphi \in \mathcal{S}(\mathbb{R})$ it holds

$$\begin{aligned} SM_{T,[a,b]}(B)(\varphi) &= S \int_0^T \int_a^b \delta_0(B_t - x) dx dt(\varphi) \\ &= \int_0^T \int_a^b S \delta_0(B_t - x)(\varphi) dx dt \\ &= \int_0^T \int_a^b \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{1}{2t} \left(\int_0^t \varphi(s) ds - x\right)^2\right) dx dt \\ &= \int_0^T \frac{1}{\sqrt{2\pi t}} \int_a^b \exp\left(-\frac{1}{2t} \left(\int_0^t \varphi(s) ds - x\right)^2\right) dx dt. \blacksquare \end{aligned}$$

Corollary 3.3. *The expected value of the occupation times of Brownian motion is given by*

$$\mathbb{E}_\mu(M_{T,[a,b]}(B)) = \int_0^T \frac{1}{\sqrt{2\pi t}} \int_a^b \exp\left(-\frac{x^2}{2t}\right) dx dt.$$

Proof: The expected value with respect to the white noise measure of the occupation times of Brownian motion is obtained by taking the S-transform and evaluating the value at 0:

$$\begin{aligned} \mathbb{E}_\mu(M_{T,[a,b]}(B)) &= SM_{T,[a,b]}(B)(0) \\ &= \int_0^T \frac{1}{\sqrt{2\pi t}} \int_a^b \exp\left(-\frac{1}{2t} \left(\int_0^t 0 ds - x\right)^2\right) dx dt \\ &= \int_0^T \frac{1}{\sqrt{2\pi t}} \int_a^b \exp\left(-\frac{x^2}{2t}\right) dx dt. \blacksquare \end{aligned}$$

4 Conclusions

We give a mathematically rigorous meaning to the occupation times of a standard Brownian motion as a Hida distribution. An expression for the expected value for the occupation times is also obtained. For the application point of view it is desirable to have a more explicit form for the expected value. This will be done in the future work. We would like also to mention that our present result is limited to one-dimensional setting. For further research we plan to generalize the result to higher spatial dimensions.

Acknowledgements

The author would like to thank anonymous reviewers for their valuable comments and suggestions. Financial support from the Institute for Research and Community Service (LPPM) Universitas Sanata Dharma through research grant No. 017/Penel./LPPM-USD/IV/2022 is gratefully acknowledged.

References

- [1] W. Bock et al, "Parameter estimation from occupation times—a white noise approach," *Communications on Stochastic Analysis*, **26**(3), 29-40, 2014.
- [2] W. Bock, J. L. da Silva, and H. P. Suryawan, "Local times for multifractional Brownian motion in higher dimensions: A white noise approach," *Infinite Dimensional Analysis, Quantum Probability and Related Topics*, **19**(4), id. 1650026, 16 pp, 2016.
- [3] W. Bock, J. L. da Silva, and H. P. Suryawan, "Self-intersection local times for multifractional Brownian motion in higher dimensions: A white noise approach," *Infinite Dimensional Analysis, Quantum Probability and Related Topics*, **23**(1), id. 2050007, 18 pp, 2020.
- [4] T. Götz et al, "A stochastic model and associated Fokker-Planck equation for the fiber lay-down process in nonwoven production processes," *SIAM Journal of Applied Mathematics*, **67**(6), 1704-1717, 2007.
- [5] M. Grothaus et al, "Application of a three-dimensional fiber lay-down model to nonwoven production processes," *Journal of Mathematics in Industry*, **4**(4), 1-19, 2014.
- [6] M. Grothaus, F. Riemann, and H. P. Suryawan, "A White Noise approach to the Feynman integrand for electrons in random media," *Journal of Mathematical Physics*, **55**(1), id. 013507, 16 pp, 2014.
- [7] M. Herty et al, "A smooth model for fiber lay-down processes and its diffusion approximations," *Kinetic and Related Models*, **2**(3), 489-502, 2009.
- [8] T. Hida et al, "White Noise. An Infinite Dimensional Calculus," *Kluwer Academic Publisher, Dordrecht*, 1993.

- [9] Z.Y. Huang and J. Yan, "Introduction to Infinite Dimensional Stochastic Analysis," *Kluwer Academic Publisher, Dordrecht*, 2000.
- [10] A. Klar, J. Maringer, and R. Wegener, "A 3D model for fiber lay-down nonwoven production processes," *Mathematical Models and Methods in Applied Sciences*, **22**(9), 1-18, 2012.
- [11] Y. Kondratiev et al, "Generalized functionals in Gaussian spaces: The characterization theorem revisited," *Journal of Functional Analysis*, **141** article number 0130, 301-318, 1996.
- [12] H.H. Kuo, "White Noise Distribution Theory," *CRC Press, Boca Raton*, 1996.
- [13] N. Obata, "White Noise Calculus and Fock Space," *Springer Verlag, Berlin*, 1994.
- [14] H.P. Suryawan, "A white noise approach to the self intersection local times of a Gaussian process," *Journal of Indonesian Mathematical Society*, **20**(2), 111-124, 2014.
- [15] H.P. Suryawan, "Weighted local times of a sub-fractional Brownian motion as Hida distributions," *Jurnal Matematika Integratif*, **15**(2), 81-87, 2019.
- [16] H.P. Suryawan, "Derivative of the Donsker delta functionals," *Mathematica Bohemica*, **144**(2), 161-176, 2019.
- [17] H.P. Suryawan, "Donsker's delta functional of stochastic processes with memory," *Journal of Mathematical and Fundamental Sciences*, **51**(3), 265-277, 2019.
- [18] H.P. Suryawan, "Pendekatan analisis derau putih untuk arus stokastik dari gerak Brown subfraksional," *Limits: Journal of Mathematics and Its Applications*, **19**(1), 15-25, 2022.

This page intentionally left blank