



ISSN 0975-7058

International Journal
of
Applied Pharmaceutics

Editorial Board

Editor-in-Chief

[Dr. Gaurav Kant Saraogi](#)

SriAurobindo Institute of Pharmacy, Indore-Ujjain State Highway, Indore, Madhya Pradesh, India

Email: editor@ijaponline.org, gauravsaraogi13@gmail.com

Associate Editors

[Dr. Genta Ida](#)

Department of Drug Sciences, University of Pavia, Italy

Email: ida.genta@unipv.it

Assistant Editors

[Dr. Awesh Kumar Yadav](#)

National Institute of Pharmaceutical Education and Research Raebareilly, Uttar Pradesh, India

Email: aweshyadav@gmail.com

[Dr. Arvind Gulbake](#)

National Institute of Pharmaceutical Education and Research: Changsari, Assam, India

Email: arvind.gulbake@gmail.com

Editorial Members

[Dr. Kailash C. Petkar](#)

Scientist 'C', Government of India, DSIR, Min. of Science and Technology, New Delhi, India

Email: petkar.kailash@gmail.com

[Dr. Alankar Shrivastava](#)

Amity Institute of Pharmacy, Amity University, Raipur, Chhattisgarh, India

Email: alankarshrivastava@gmail.com

[Dr. Tarang Nema](#)

International Flavours and Fragrances, Singapore

[Dr. Carlotta Marianecchi](#)

Dipartimento di Chimica e Tecnologie del Farmaco, Sapienza Università di Roma, Rome, Italy

Email: carlotta.marianecchi@uniroma1.it

[Dr. Manoj Nahar](#)

Sun Pharmaceutical Industries Limited, Vadodara, Gujarat, India

Email: mnjnahar@yahoo.co.in

[Dr. Tarek Abdelnaby Ahmed](#)

Department of Pharmaceutics and Industrial Pharmacy, Faculty of Pharmacy, KAU, Jeddah, KSA

Email: dr_tarek_nour@yahoo.com

[Dr. Elizabeth Igne Ferreira](#)

Faculty of Pharmaceutical Sciences, University of Sao Paulo, Brazil

Email: elizabeth.igne@gmail.com

[Dr. Surya Prakasarao Kovvasu](#)

Western University of Health Sciences, Pomona, California, USA

Email: skovvasu@westernu.edu

[Dr. N. Kanagathara](#)

Saveetha School of Engineering, Saveetha University, Chennai, India

Email: kanagathara23275@gmail.com

[Dr. Mohammed Elmowafy Gomaa Aburaia](#)

Department of Pharmaceutics, College of Pharmacy, Jouf University, Saudi Arabia

Email: melmowafy@ju.edu.sa

[Dr. Liang Chen](#)

Wenzhou Medical University, Wenzhou, P. R. China

Email: cheng_zhuan0101@wuxibiotics.com

[Dr. Franca Castiglione](#)

Department "G. Natta", Politecnico di Milano, Italy

Email: franca.castiglione@polimi.it

[Dr. Iman Emam Omar Gomaa](#)

Faculty of Pharmacy, University for Modern Sciences and Arts (MSA) Cairo - Egypt

Email: igomaa@msa.eun.eg

[Dr. Basant Amarji](#)

UIPS, Punjab University, Chandigarh, Punjab, India

Email: basantamarji@gmail.com

[Dr. Rabab Kamel](#)

Pharmaceutical Technology Department, National Research Centre, Egypt

Email: drbabak@hotmail.com

[Dr. Satish Shilpi](#)

Ravishankar College of Pharmacy, Bhopal, MP, India

Email: shilpisatish@gmail.com

[Dr. Umeyor Chukwuebuka Emmanuel](#)

Faculty of Pharmaceutical Sciences, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria

Email: ec.umeyor@unizik.edu.ng

[Dr. Yosra S.R. Elnaggar](#)

Faculty of Pharmacy and Drug Manufacturing, Pharos University, Alexandria, Egypt

Email: yosra_pharm@yahoo.com

[Dr. Sumeet Kapoor](#)

IIT New Delhi, India

Email: s.kapooriitd@gmail.com

[Dr. Arif Nur Muhammad Ansori](#)

Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia

Email: ansori.anm@gmail.com

[Dr. Dinesh Nyavanandi](#)

Research Scientist, Formulation Development Cerevel Therapeutics, MA 02141, USA

Email: ndinesh624@gmail.com



Online ISSN: 0975-7058



[CONTINUED IN 2024 \[Q2\]](#)

How we claim? [Click Scopus indexing IJAP 2024](#) to learn and understand

1.4

2023
CiteScore

62nd percentile

Powered by 



Plagiarism Check



Embase

It's Embase, Not Expanded Embase, [Learn in 1 Min](#)

MOST READ

[FORMULATION AND EVALUATION OF TRANSDERMAL PATCHES OF PANTOPRAZOLE SODIUM](#)

📄 1035

[OCULAR DRUG DELIVERY SYSTEM: CHALLENGES AND APPROACHES](#)

📄 674

[OVERVIEW ON FLOATING DRUG DELIVERY SYSTEM](#)

📄 638

[DERMATOLOGIC GELS SPREADABILITY MEASURING METHODS COMPARATIVE STUDY](#)

📄 450

[FORMULATION AND EVALUATION OF EFFERVESCENT GRANULES OF IBUPROFEN](#)

📄 417

[Our Journals](#) || [Open Access Policy](#) || [Publication & Peer Review Policy](#) || [Publication Ethics](#)

The publication is licensed under a [Creative Commons License \(CC BY\)](#). [View Legal Code](#)

Copyright © 2024 All Rights Reserved, [Innovare Academic Sciences](#) | Powered By [CyberDairy](#)



Vol 16, Issue 5 (Sep-Oct), 2024

PUBLISHED: 07-09-2024

REVIEW ARTICLE(S)

NAVIGATING NITROSAMINES: ORIGIN, DETECTING, ANALYSING AND REGULATING IMPURITIES IN PHARMACEUTICALS

TEJAS SANJAY SONAWANE, TANVI PAINGINKAR, RUCHIR BAWADIA, SG VASANTHARAJU, MLIDDUKRISHNA BADAMANE SATHYANARAYANA, GUNDAWAR RAVI

1-11

[VIEW ABSTRACT](#)

[PDF](#)

[DOWNLOAD PDF](#)

10.22159/ijap.2024v16i5.51376

BEYOND THE HORIZON: RECENT ADVANCES IN HOT MELT EXTRUSION TECHNIQUES AND TECHNOLOGIES

MAGED MOHAMMED ABDO MOHSEN, AMIT B. PATIL, MAGED ALKANAD, DARSHAN PATIL

12-21

[VIEW ABSTRACT](#)

[PDF](#)

[DOWNLOAD PDF](#)

10.22159/ijap.2024v16i5.51425

METALLOPROTEIN PARAMETERS IN MOLECULAR DYNAMICS SIMULATION FOR AMBER, CHARMM, GROMACS, AND NAMD: A SYSTEMATIC REVIEW

PURNAWAN PONTANA PUTRA, NAJMIATUL FITRIA, AIYI ASNAWI, AKMAL DJAMAAN

22-30

[VIEW ABSTRACT](#)

[PDF](#)

[DOWNLOAD PDF](#)

10.22159/ijap.2024v16i5.51513

STATINS: A NEW THERAPEUTIC APPROACH FOR THE TREATMENT OF NEUROPATHIC PAIN

INDU MELKANI, BIMLESH KUMAR, NARENDRA KUMAR PANDEY, DILEEP SINGH BAGHEL, SAURABH SINGH

22-30

[VIEW ABSTRACT](#)

[PDF](#)

[DOWNLOAD PDF](#)

10.22159/ijap.2024v16i5.50456

DIAGNOSTIC AND THERAPEUTIC ROLE OF MESOPOROUS SILICA NANOPARTICLES IN COMBATING CANCER

NUPUR KATARIYA, ARVIND SINGH FARSWAN, NIDHI NAINWAL, GANESH KUMAR

31-37

[VIEW ABSTRACT](#)

[PDF](#)

[DOWNLOAD PDF](#)

10.22159/ijap.2024v16i5.51647

CURRENT PERSPECTIVES ON USING NANOPARTICLES FOR DIABETES MANAGEMENT

NITESH KUMAR YADAV, RUPA MAZUMDER, ANJINA RANI, ARVIND KUMAR

38-45

[VIEW ABSTRACT](#)

[PDF](#)

[DOWNLOAD PDF](#)

10.22159/ijap.2024v16i5.51084

THERAPEUTIC IMPACT OF NANOMEDICINE FOR THE TREATMENT OF NEUROPATHIC PAIN: PRINCIPLE, PROSPECTIVE AND FUTURE

INDU MELKANI, BIMLESH KUMAR, NARENDRA KUMAR PANDEY, SAURABH SINGH, DILEEP SINGH BAGHEL, KAWATALA SUDHAKAR
46-58

[VIEW ABSTRACT](#)

[PDF](#)

[DOWNLOAD PDF](#)

[doi: 10.22159/ijap.2024v16i5.50457](#)

LEVERAGING THE SUCCESS OF MRNA LIPID NANOPARTICLE VACCINE IN COVID-19 TREATMENT: A NARRATIVE REVIEW ON ITS POTENTIAL APPLICATION IN MALARIA TREATMENT

FRANKLYN NONSO IHEAGINAM, ENIOLA DEBORAH ADEDOYIN
59-66

[VIEW ABSTRACT](#)

[PDF](#)

[DOWNLOAD PDF](#)

[doi: 10.22159/ijap.2024v16i5.50885](#)

AQUASOMES UNVEILED: TRANSFORMING DRUG DELIVERY WITH CUTTING-EDGE THERAPEUTIC CARRIERS AND RECENT BREAKTHROUGHS

VIJAYARAGHAWAN KRISHNAN, VIVEKANANDAN ELANGO, SAM HARRISON SAM JENKINSON, BALAGEE MUTHUKUMAR, SANGAY RAJA DHANRAJ, DHANDAPANI NAGASAMY VENKATESH
67-76

[VIEW ABSTRACT](#)

[PDF](#)

[DOWNLOAD PDF](#)

[doi: 10.22159/ijap.2024v16i5.50936](#)

EMERGING FRONTIERS: ADVANCEMENTS IN BIO-NONMATERIAL'S AND NON-INVASIVE STRATEGIES FOR COMBATING CANCER THROUGH PHOTO THERMAL THERAPY

YOKESH S., TEEJESWARI R., JALANIY V., LOKESHWAR R.
86-98

[VIEW ABSTRACT](#)

[PDF](#)

[DOWNLOAD PDF](#)

[doi: 10.22159/ijap.2024v16i5.51113](#)

ORIGINAL ARTICLE(S)

DEVELOPMENT AND OPTIMIZATION OF THE WOUND HEALING ELECTROSPUN POLYURETHANE/COLLAGEN/PHYTOCERAMIDES NANOFIBERS USING THE BOX-BEHNKEN EXPERIMENTAL DESIGN) QUALITY BY DESIGN)

TASSNEIM M. EWEDAH, MOHAMED EL-NABARAWI, MAHMOUD H. TEAIMA, SAMMAR FATHY ELHABAL, KAMEL R. SHOUEIR, ABDALLAH M. HAMDY, AHMED ABDALLA
99-100

[VIEW ABSTRACT](#)

[PDF](#)

[DOWNLOAD PDF](#)

[doi: 10.22159/ijap.2024v16i5.51510](#)

UNLOCKING THE POTENTIAL: ENHANCING SOLUBILITY AND BIOAVAILABILITY OF ACYCLOVIR THROUGH SOLID DISPERSION FORMULATIONS

RUBA MALKAWI, SULEIMAN AL-OLIMAT, JUMANA TAWALBEH
111-118

[VIEW ABSTRACT](#)

[PDF](#)

[DOWNLOAD PDF](#)

[doi: 10.22159/ijap.2024v16i5.51313](#)

FORMULATION AND EVALUATION OF ETODOLAC AND TRIAMCINOLONE ACETONIDE LOADED NANO LIPID CARRIER GELS FOR THE THERAPEUTIC MANAGEMENT OF OSTEOARTHRITIS PAIN THROUGH TOPICAL ADMINISTRATION: A COMPARATIVE STUDY

SOUVIK CHAKRABORTY, N. VISHAL GUPTA, VIKAS JAIN, BALAMURALIDHARA V.
119-132

[VIEW ABSTRACT](#)

[PDF](#)

[DOWNLOAD PDF](#)

[doi: 10.22159/ijap.2024v16i5.51326](#)

COMBINING LIQUID CHROMATOGRAPHY WITH TANDEM MASS SPECTROMETRY (LC-MS/MS) TECHNIQUE, SPECTROPHOTOMETERVERIFICATION AND AN IN SILICO STUDY OF 96% ETHANOL EXTRACT OF SPIRULINA PLATENSIS

TIWUK SUSANTININGSIH, FADILAH FADILAH, ANI RETNO PRIJANTI, NOVI SILVIA HARDIANY
133-138

[VIEW ABSTRACT](#)[PDF](#)[DOWNLOAD PDF](#)[doi](#) 10.22159/ijap.2024v16i5.51339

ANALYTICAL QUANTIFICATION OF BOSUTINIB IN NANOCARRIER USING UV AND HPLC: METHOD DEVELOPMENT AND VALIDATION

RISHABH AGADE, UJIBAN HUSSAIN, SAGAR TRIVEDI, VEENA BELGANWAR

139-145

[VIEW ABSTRACT](#)[PDF](#)[DOWNLOAD PDF](#)[doi](#) 10.22159/ijap.2024v16i5.51394

DEVELOPMENT OF NANOSPONGES-BASED TOPICAL FORMULATION FOR THE EFFECTIVE DELIVERY OF SELECTED ANTIFUNGAL DRUG

RUDRĀJU ANUSHA, MOTHILAL M.

146-156

[VIEW ABSTRACT](#)[PDF](#)[DOWNLOAD PDF](#)[doi](#) 10.22159/ijap.2024v16i5.51466

INTEGRATIVE QSAR ANALYSIS OF OXADIAZOLE DERIVATIVES: RESOLVING MOLECULAR DETERMINANTS FOR ANTI-TUBERCULAR ACTIVITY AND RATIONAL DRUG DESIGN

NEHA H. SUWARNA, VIKSHITH RAJ, SHIHABUDHEEN HAREES, JESSY ELIZABETH MATHEW, LALIT KUMAR, RUCHI VERMA

157-165

[VIEW ABSTRACT](#)[PDF](#)[DOWNLOAD PDF](#)[doi](#) 10.22159/ijap.2024v16i5.51468

FORMULATION AND EVALUATION OF LICORICE OIL-BASED EMULGEL FOR THE TREATMENT OF PSORIASIS

PRIYANKA JUREL, SHIV BAHADUR, MEENAKSHI BAIJPAI

166-175

[VIEW ABSTRACT](#)[PDF](#)[DOWNLOAD PDF](#)[doi](#) 10.22159/ijap.2024v16i5.51471

INVESTIGATING MULTITARGET POTENTIAL OF MUCUNA PRURIENS AGAINST PARKINSON'S DISEASE: INSIGHTS FROM MOLECULAR DOCKING, MMGBSA, PHARMACOPHORE MODELLING, MD SIMULATIONS AND ADMET ANALYSIS

ZAKIYA FATHIMA C., JAINEY P. JAMES, MAHENDRA GOWDRU SRINIWASA, SINDHU T. J, MARIYAM JOUHARA B. M., B. C.

REWANASIDDAPPA, SUDEEP D. GHATE

176-193

[VIEW ABSTRACT](#)[PDF](#)[DOWNLOAD PDF](#)[doi](#) 10.22159/ijap.2024v16i5.51474

THE DEVELOPMENT OF METHOD FOR ACTIVATING PHARMACEUTICAL SUBSTANCES WITH SUBSEQUENT IN SITU STUDY OF MODIFIED POWDER PROPERTIES

ELENA USPENSKAYA, EKATERINA S. KLZMINA, HOANG T. N. QUYNH, ALEKSEY A. TIMOFEEV, TATIANA V. MAXIMOVA

194-203

[VIEW ABSTRACT](#)[PDF](#)[DOWNLOAD PDF](#)[doi](#) 10.22159/ijap.2024v16i5.51481

FORMULATION AND IN VIVO EVALUATION OF NANOEMULGEL-CONTAINING COCOA POD HUSK (THEOBROMA CACAO L.) EXTRACT AS TOPICAL ORAL PREPARATION

YANI CORVIANINDYA RAHAYU, ERNIE MADURATNA SETIAWATIE, RETNO PUDJI RAHAYU, BANLIN KUSUMAWARDANI, NINIK MAS ULFA

204-210

[VIEW ABSTRACT](#)[PDF](#)[DOWNLOAD PDF](#)[doi](#) 10.22159/ijap.2024v16i5.51294

MOLECULAR TARGETS AS POTENTIAL PI3KA INHIBITORS AGAINST AGGRESSIVE METASTATIC DUCTAL AND LOBULAR CARCINOMA

ARPITH MATHEW, SUBHAM DAS, LATE ALEX JOSEPH, SUMIT RAOSAHEB BIRANGAL, JANE MATHEW

211-219

[VIEW ABSTRACT](#)[PDF](#)[DOWNLOAD PDF](#)[doi](#) 10.22159/ijap.2024v16i5.51514

AUTHENTICATION OF GRAPE SEED FACE OIL USING FTIR SPECTROSCOPY COMBINED WITH CHEMOMETRICS TECHNIQUES

ANDI PRAYOGA, ANJAR WINDARSIH, WJURI APRIYANA, FLORENTINUS DIKA OCTA RISWANTO, ENADE PERDANA ISTYASTONO
220-224

[VIEW ABSTRACT](#)[PDF](#)[DOWNLOAD PDF](#)[doi](#) 10.22159/ijap.2024v16i5.51525

HESPERETIN DERIVATIVES AS PPAR γ AGONIST: A PHARMACOPHORE APPROACH

RAMANATHAN MUTHIAH, VIJAYALAKSHMI CHINNIAH, MAIDA ENGELS. S. E.
225-233

[VIEW ABSTRACT](#)[PDF](#)[DOWNLOAD PDF](#)[doi](#) 10.22159/ijap.2024v16i5.51538

QUANTITATIVE DETERMINATION OF METHYL-4-CHLOROBUTYRATE, A POTENTIAL GENOTOXIC IMPURITY, CONTENT IN MOXIFLOXACIN HCL BY GC-EI-MS

K. APARNA, K. VIJAYA RACHEL, K. M. V. NARAYANA RAO
234-241

[VIEW ABSTRACT](#)[PDF](#)[DOWNLOAD PDF](#)[doi](#) 10.22159/ijap.2024v16i5.51551

DEVELOPMENT OF GUGGULSTERONE-LOADED PHYTOSOMES: A QUALITY BY DESIGN-BASED CHARACTERIZATION AND OPTIMIZATION STUDIES

JAMAL BASHA DUDEKULA, JEBASTIN KOILPILLAI, DAMODHARAN NARAYANASAMY
242-251

[VIEW ABSTRACT](#)[PDF](#)[DOWNLOAD PDF](#)[doi](#) 10.22159/ijap.2024v16i5.51559

SYNTHESIS, CHARACTERIZATION AND IN VITRO EVALUATION OF VATERITE MICROPARTICLES

DEENA JOSE, D. KUMUDHA
252-257

[VIEW ABSTRACT](#)[PDF](#)[DOWNLOAD PDF](#)[doi](#) 10.22159/ijap.2024v16i5.51569

DESIGNING, DEVELOPMENT AND EVALUATION OF GASTRORETENTIVE FLOATING HBS SYSTEM OF METFORMIN: IN VITRO IN VIVO STUDIES

RADHESHYAM SAMANTA, GAURAV TIWARI, NAVEEN GUPTA, DHARMENDRA SINGH RAJPUT
258-265

[VIEW ABSTRACT](#)[PDF](#)[DOWNLOAD PDF](#)[doi](#) 10.22159/ijap.2024v16i5.51674

FORMULATION, CHARACTERIZATION, AND TYROSINASE INHIBITORY ASSAYS OF NIACINAMIDE-LOADED NANOPARTICLE GEL AS A SKIN WHITENING AGENT

RISAAHDYANI, NOR LATIFAH, HAYATUS SA'ADAH, ERLINA FATMASARI, IRFAN ZAMZANI
266-274

[VIEW ABSTRACT](#)[PDF](#)[DOWNLOAD PDF](#)[doi](#) 10.22159/ijap.2024v16i5.51750

VALIDATED REVERSED-PHASE HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY METHOD FOR THE ESTIMATION OF TETRABENAZINE IN SELF-NANO EMULSIFYING DRUG DELIVERY SYSTEMS

SHASHI, NARENDRA KUMAR PANDEY, BIMLESH KUMAR, SACHIN KUMAR SINGH, DILEEP SINGH BAGHEL, KALWATALA SUDHAKAR, SAURABH SINGH
275-282

[VIEW ABSTRACT](#)[PDF](#)[DOWNLOAD PDF](#)[doi](#) 10.22159/ijap.2024v16i5.50489

NANOCREAM FOUNDATION FORMULATION LOADED WITH ETHYL ACETATE FRACTION FROM MELASTOMA MALABATHRICUM L.

LIZA PRATIWI, RAFIKA SARI, PRATIWI APRIDAMAYANTI

283-290

 VIEW ABSTRACT

 PDF

 DOWNLOAD PDF

 10.22159/ijap.2024v16i5.50580

MOLECULAR DOCKING OF BIFLAVONOIDS FROM GENUS ARAUCARIA AS DENV NS5 RNA-DEPENDENT RNA POLYMERASE INHIBITOR USING YASARA AND PLANTS PROGRAMS

LAKSMI AMBARSARI, NAJMA AULIA NUR, SYIFA SILFANI RODOTUL ZANAH, KURNIAWANTI, HANHAN DIAN HAR, SITI WARNASIH, DYAH UTAMI CAHYANING RAHAYU, PURNANTININGSIH SUGITA

291-299

 VIEW ABSTRACT

 PDF

 DOWNLOAD PDF

 10.22159/ijap.2024v16i5.50833

BIO-ANALYTICAL METHOD DEVELOPMENT AND VALIDATION OF FINASTERIDE, TADALAFIL, AND ITS APPLICATION TO PHARMACOKINETIC STUDIES IN RAT PLASMA BY USING LC-MS/MS

YESUPADAMU RAYINUTHALA, DAVID RAJU MEDEPALLI, A. LAKSHMANARAO

300-308

 VIEW ABSTRACT

 PDF

 DOWNLOAD PDF

 10.22159/ijap.2024v16i5.50858

COMPUTATIONAL ASSESSMENT OF UNDARIA PINNATIFIDA AND MORINGA OLEIFERA COMPOUNDS AS ANTI-OBESITY AGENTS

C. SAI KALYANI YOGINI, CHITTA SURESH KUMAR, C. M. ANURADHA, C. H. M. KUMARI CHITTURI

309-326

 VIEW ABSTRACT

 PDF

 DOWNLOAD PDF

 10.22159/ijap.2024v16i5.50867

INVESTIGATING THE ROLE OF NANOPARTICLE-BASED CURCUMIN IMPLANTS IN PREVENTION OF POST-LAPAROTOMY PERITONEAL ADHESION: AN IN VIVO STUDY

MOHAMMAD AMIN KABOLI, DHIRYA ALTEMEY, MOOSA JAVDANI, HOSSEIN AMINI KHOEI, PARISA MEHREGANZADEH, FATEMEH DRISS, MEHRDAD KARIMI, PEGAH KHOSRAVIAN

327-332

 VIEW ABSTRACT

 PDF

 DOWNLOAD PDF

 10.22159/ijap.2024v16i5.50976

DRUG EXCIPIENT ADSORPTION TECHNIQUE FOR THE PREPARATION OF LOW-DOSE TABLETING

NIKHITHA K. SHANMUKHAN, RAMYA G., ARUN RADHAKRISHNAN

333-339

 VIEW ABSTRACT

 PDF

 DOWNLOAD PDF

 10.22159/ijap.2024v16i5.50991

MOLECULAR DOCKING STUDY AS THERAPEUTIC APPROACH FOR TARGETING CHOLECYSTOKININ IN PANCREATIC CANCER

OKTAVIAN ARYA PUTRA, TESIA AISYAH RAHMANIA, EDITHA RENESTEEN

340-349

 VIEW ABSTRACT

 PDF

 DOWNLOAD PDF

 10.22159/ijap.2024v16i5.51027

DEVELOPMENT AND VALIDATION OF A SPECIFIC AND UNIQUE DUAL POLARITY ESI-LC-MS/MS METHOD FOR SIMULTANEOUS DETERMINATION OF SEMAGLUTIDE AND DAPAGLIFLOZIN IN HUMAN PLASMA

JAGAPATHI RAJU WATSAWYI, NALANDA BABY REVU

350-358

 VIEW ABSTRACT

 PDF

 DOWNLOAD PDF

 10.22159/ijap.2024v16i5.51095

FABRICATION, OPTIMIZATION AND IN VITRO CYTOTOXICITY EVALUATION OF DASATINIB MONOHYDRATE-LOADED NANOPARTICLES

AJAY SAROHA, RAWINDER VERMA, VINEET MITTAL, DEEPAK KAUSHIK

359-367

 [VIEW ABSTRACT](#)

 [PDF](#)

 [DOWNLOAD PDF](#)

 [10.22159/ijap.2024v16i5.51125](https://doi.org/10.22159/ijap.2024v16i5.51125)

THE BINARY AND TERNARY AMORPHOUS SYSTEMS OF CANDESARTAN CILEXETIL PREPARATION TO IMPROVE ITS SOLUBILITY

FIKRI ALATAS, HESTIARY RATIH, TITTA HARTYANA SUTARNA, MUHAMAD LUTHFI FAUZI

368-373

 [VIEW ABSTRACT](#)

 [PDF](#)

 [DOWNLOAD PDF](#)

 [10.22159/ijap.2024v16i5.51141](https://doi.org/10.22159/ijap.2024v16i5.51141)

PULMONARY DRUG DELIVERY SYSTEMS OF POSACONAZOLE: DEVELOPMENT AND OPTIMIZATION THROUGH QUALITY BY DESIGN

INDIRAMUZIB YALLAMALLI, SREEVIDYA PUVVALA

374-382

 [VIEW ABSTRACT](#)

 [PDF](#)

 [DOWNLOAD PDF](#)

 [10.22159/ijap.2024v16i5.51163](https://doi.org/10.22159/ijap.2024v16i5.51163)

SMART CHITOSAN-BASED MICROBEAD FORMULATION FOR COLON-TARGETED DELIVERY OF LACTOFERRIN

LUQMAN OLADOYE, AZAD SADRADDIN, SHWANABRAIM

383-388

 [VIEW ABSTRACT](#)

 [PDF](#)

 [DOWNLOAD PDF](#)

 [10.22159/ijap.2024v16i5.51185](https://doi.org/10.22159/ijap.2024v16i5.51185)

INHIBITION ACTIVITY OF LIQUID SMOKE COCOS NUCIFERA L. ON DPP-IV AND AGE-RAGE IN SILICO AND IN VITRO: ANTIDIABETIC AND ANTI-INFLAMMATORY ACTIVITY

MASFRIA, HENNY LUCIDA, YUSIATIFAH, HAFID SYAHPUTRA, HANNA MURTI SIHOMBING

389-396

 [VIEW ABSTRACT](#)

 [PDF](#)

 [DOWNLOAD PDF](#)

 [10.22159/ijap.2024v16i5.51231](https://doi.org/10.22159/ijap.2024v16i5.51231)

BRINZOLAMIDE-LOADED ETHO-LECIPEX FOR EFFECTIVE OCULAR MANAGEMENT OF GLAUCOMA: D-OPTIMAL DESIGN OPTIMIZATION AND IN VIVO EVALUATION

SARA NAGEEB EL-HELALY, HAYDER A. HAMMOODI, MINA I. TADROS, NERMEEN A. ELKASABGY

397-406

 [VIEW ABSTRACT](#)

 [PDF](#)

 [DOWNLOAD PDF](#)

 [10.22159/ijap.2024v16i5.51259](https://doi.org/10.22159/ijap.2024v16i5.51259)

ZAP-AIRTMICRONIZER: IMPACT ON NEBULIZED SALBUTAMOL SULPHATE AND IPRATROPIUM BROMIDE PARTICLE SIZE

MUHAMMAD IZZUDDIN ZAMERY, SHARIZA SAHUDIN, SHAHARIZUAN ABDUL RAHMAN, ALBERT KOW

407-417

 [VIEW ABSTRACT](#)

 [PDF](#)

 [DOWNLOAD PDF](#)

 [10.22159/ijap.2024v16i5.51277](https://doi.org/10.22159/ijap.2024v16i5.51277)

EVALUATION OF POPULATION PHARMACOKINETICS OF ORAL DIGOXIN IN VENOUS PLASMA

SIRAJUDEEN MAHABOOB, ARUN K. P., S. D. RAJENDRAN, GNK GANESH

418-424

 [VIEW ABSTRACT](#)

 [PDF](#)

 [DOWNLOAD PDF](#)

 [10.22159/ijap.2024v16i5.51291](https://doi.org/10.22159/ijap.2024v16i5.51291)



Online ISSN: 0975-7058



CONTINUED IN 2024 [Q2]

How we claim? [Click Scopus indexing IJAP 2024](#) to learn and understand

1.4 2023
CiteScore

62nd percentile

Powered by **Scopus**



Plagiarism Check



Embase

It's Embase, Not Expanded Embase, [Learn in 1 Min](#)

MOST READ

FORMULATION AND EVALUATION OF TRANSDERMAL PATCHES OF PANTOPRAZOLE SODIUM

📄 1066

OCULAR DRUG DELIVERY SYSTEM: CHALLENGES AND APPROACHES

📄 680

OVERVIEW ON FLOATING DRUG DELIVERY SYSTEM

📄 662

DERMATOLOGIC GELS SPREADABILITY MEASURING METHODS COMPARATIVE STUDY

📄 465

FORMULATION AND EVALUATION OF EFFERVESCENT GRANULES OF IBUPROFEN

📄 422

[Our Journals](#) || [Open Access Policy](#) || [Publication & Peer Review Policy](#) || [Publication Ethics](#)

The publication is licensed under a [Creative Commons License \(CC BY\)](#). [View Legal Code](#)

Copyright © 2024 All Rights Reserved, [Innovare Academic Sciences](#) | Powered By [CyberDairy](#)

AUTHENTICATION OF GRAPE SEED FACE OIL USING FTIR SPECTROSCOPY COMBINED WITH CHEMOMETRICS TECHNIQUES**ANDI PRAYOGA¹, ANJAR WINDARSIH², WURI APRIYANA², FLORENTINUS DIKA OCTA RISWANTO^{1,3}, ENADE PERDANA ISTYASTONO^{1,3}**¹Magister of Pharmacy Study Program, Faculty of Pharmacy, Universitas Sanata Dharma, Yogyakarta-55282, Indonesia, ²Research Center for Food Technology and Processing, National Research and Innovation Agency, Jl. Jogja Wonosari km 31.5, Gunungkidul, Yogyakarta, Indonesia, ³Research Group of Computer-Aided Drug Design and Discovery of Bioactive Natural Products, Faculty of Pharmacy, Sanata Dharma University, Yogyakarta-55282, Indonesia*Corresponding author: Florentinus Dika Octa Riswanto; Email: dikaocta@usd.ac.id

Received: 21 May 2024, Revised and Accepted: 04 Jul 2024

ABSTRACT**Objective:** This research aims to authenticate grape seed oil products using FTIR spectroscopy combined with chemometric methods.**Methods:** In the initial stage, exploratory data analysis was carried out by applying the main components with the Principal Component Analysis (PCA) model. Second derivative spectra resulting from preprocessing of the original spectra are used to create multivariate Principal Component Regression (PCR) and Partial Least Squares (PLS) calibration models. The second derivative spectra of Grape Seed Oil (GO), Olive Oil (OO), and the binary mixture GO+OO are utilized to generate a sparse partial least squares-discriminant analysis (SPLS-DA) model.**Results:** The PCA model was successfully obtained with visualization that depicted a total of 93.8% in the first and second dimensions. Multivariate calibration produced the best model in PLS with second derivative spectra for both GO and OO. PLS model for GO resulted the value of R_{cv}^2 , R_{cv}^2 , R_{cv}^2 , RMSEC, RMSECV, and RMSEP of 0.990, 0.992, 0.982, 0.700, 1.557, and 2.331, respectively. The SPLS-DA model was successfully built and discriminated with AUC-ROC values of 1.000, 1.000, and 0.994 for GO, OO, and GO+OO, respectively.**Conclusion:** Authentication of grape seed face oil can be undertaken using FTIR spectroscopic methods and chemometric techniques, which can produce high sensitivity and specificity values.**Keywords:** Grape seed oil, Face oil, Adulteration, FTIR spectroscopy, Chemometrics© 2024 The Authors. Published by Inovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<https://creativecommons.org/licenses/by/4.0/>) DOI: <https://doi.org/10.22159/ijap.2024v16i5.51525> Journal homepage: <https://inovareacademic.in/journals/index.php/ijap>**INTRODUCTION**

The high demand for cosmetic products for both men and women provides opportunities for the cosmetic industry in Indonesia [1-3]. Modern cosmetics are reported to contain harmful chemicals and have ecological impacts on the environment [4]. With growing awareness, consumers tend to prefer natural products [5-8]. One of the most needed cosmetic products for both teenagers and adults is skincare, especially facial skincare [9]. One type of facial skincare is face oil, aimed at moisturizing and smoothing. One widely used type of face oil is grape seed oil.

Grape seed oil contains antioxidants that are beneficial for skin care. The antioxidants contained in it are vitamin E and Oligomeric Proanthocyanidins (OPC) [10]. In the previous research, grape seeds were reported to contain 8-20% oil in the dry state [11]. Grape seed oil has a high vitamin E content, ranging from 1 to 53 mg per 100 g of oil, which is higher than olive oil [10]. Even though grape seed oil has greater economic value than other oils, they appear to share many similar physical characteristics. Currently, research on grape seed oil to detect whether the grape seed extract is adulterated or not is still lacking. Therefore, it is very significant to develop a simple and fast method for the detection of grape seed oil adulteration [12].

In previous related research, analysis of sunflower oil in a ternary mixture with grape seed oil and candlenut oil in a mixed system using FTIR spectroscopy and chemometrics was successfully carried out [13]. However, research that applies the FTIR spectroscopy method combined with chemometric techniques on grape seed face oil has never been carried out. This research aims to authenticate grape seed oil products using FTIR spectroscopy combined with chemometric methods. The undertaken chemometric techniques include Principal Components Analysis (PCA), Partial Least Squares (PLS), and Partial Least Squares Discriminant Analysis (PLS-DA).

MATERIALS AND METHODS**Materials**

Grape Seed Oil (GO) and Olive Oil (OO) are obtained from distributors in Indonesia, with a Certificate of Analysis to verify and prove their authenticity.

Tools and software

In this research, a complete instrument was used with Thermo Nicolet IS10 Thermo Scientific FT-IR spectrophotometer, Sacorex® Micropipette with sizes of 20-200 μ l and 100-1000 μ l and a Globolab measuring cup, the results of the FTIR spectrum data were exported to Excel 2019 software (Microsoft Inc, USA) and saved as a .csv file before being analyzed using statistical software R Version 4.3.2.

Methods

Methods used in this study were developed according to the previous study from Riswanto et al (2023) [14] with several modifications. In this research, chemometrics techniques were implemented to authenticate grape seed face oil from olive oil as adulterant.

Calibration and validation sample preparation

Calibration and validation solution test samples containing grape seed oil and olive oil were made from a mixture of the two ingredients with a concentration range of 0-100% (v/v) to obtain 15 variations of the concentration of the calibration solution and 30 variations of the concentration of the validation solution.

Sample preparation for discrimination

The solution for PLS-DA analysis is available in three different samples, namely grape seed oil, olive oil, and a binary mixture that contains grape seed oil and olive oil with a range of concentration 0-100% (v/v). Eight pure grape seed oil solutions, eight pure olive oil

solutions, and forty-three solutions containing grape seed oil and olive oil were obtained and labelled to create a discrimination analysis model.

FTIR spectra acquisition

FTIR spectra acquisition was carried out using an FTIR spectrophotometer in which samples of pure grape seed oil, pure olive oil, and the binary mixed of grape seed oil were placed on the ATR crystal and then measured in the mid-infrared area (4000-600 cm^{-1}). FTIR spectra used absorbance mode on the resolution of 8 cm^{-1} with a number of scans of 32. Air (blank) spectra measurements were applied before each sample measurement as the background spectra. Each sample was measured using three replications. The cleaning of ATR crystals was done using ethanol after each sample measurement.

Exploratory data analysis using the PCA method

The FTIR spectra obtained from the data acquisition stage were then evaluated [15]. The dominant peaks of the analysis, namely grape seed oil, olive oil, and a mixture of samples containing grape seed oil and olive oil, were used to construct the PCA model. Scree plots, variable plots and individual plots are displayed to visualize the constructed model.

Multivariate calibration analysis

The PLS multivariate calibration model produced predictive models for the two oils. The performance of the multivariate calibration model was examined by evaluating statistical parameters, including coefficients of determination for calibration (R_{cal}^2), cross-validation (R_{cv}^2), validation (R_{val}^2), root mean square error of calibration (RMSEC), root mean square error of cross-validation (RMSECV) and root mean square error of prediction (RMSEP). The cross-validation process, as an internal validation, applies the leave-one-out technique. The selected multivariate calibration model for each oil was determined by evaluating R_{cal}^2 , R_{cv}^2 , R_{val}^2 , RMSEC, RMSECV, and RMSEP [16].

Construction of the PLS-DA model

The PLS-DA model utilized the FTIR spectrum of grape seed oil, olive oil, and a mixture of samples containing grape seed oil and olive oil. Background prediction and 3D individual plots visualized the

discrimination model. Evaluation of model performance applied Area Under Curve-Receiver Operating Characteristics (AUC-ROC). Optimization of this model was further carried out by selecting output variables until the final result was obtained considering the error rate in classification and feature selection.

RESULTS

This research develops a rapid detection technique for counterfeiting grape seed face oil using the Fourier Transform Infrared (FTIR) spectroscopy method combined with chemometric techniques. The FTIR spectroscopy method combined with chemometrics was chosen in this research because this analysis requires minimal material preparation, is easy, and fast [17, 18].

The infrared spectrum characteristics of samples in the range of 4000-600 cm^{-1} area produce 12 dominant peaks for grape seed oil, olive oil and mixtures, namely at 2970, 2947, 2922, 2912, 1743, 1653, 1462, 1376, 1238, 1160, 1099, and 718 cm^{-1} (fig. 1).

Exploratory data analysis using PCA

Principal Component Analysis (PCA) is a data processing technique to construct linear multivariate models from complex data sets [19]. PCA facilitates visualization of data groupings, initial evaluation of similarities between groups or classes, and finding factors or reasons behind observed patterns through correlation by considering chemical or chemical-physics properties [20]. The processing of data from an analytical instrument using an FTIR spectrophotometer with a range of 4000-600 cm^{-1} with a resolution of 8 cm^{-1} produces the absorbance of twelve main peaks; this data then produces a PCA model (fig. 2).

Multivariate calibration analysis

In this research, a multivariate calibration is developed using PCR and PLS techniques; these two techniques are used to predict GO and OO content and to obtain the best prediction model in various spectral modes (table 1). Absorption data at each wave number in the range of 4000-600 cm^{-1} is further processed through a preprocessing stage in order to produce the original spectrum, first derivative, second derivative, standard normal variate (SNV), and savitzky golay (SG). Plot of grape seed oil and olive oil predictions resulting from the best calibration model were presented in fig. 3.




Fig. 1: FTIR spectra of GO (Grape seed Oil), OO (Olive Oil), and GO+OO (Grape seed Oil+Olive Oil)




Fig. 2: Results of principal component analysis of the original spectra with 12 selected wavenumbers in the form of scree plot (a), variable plot (b), and individual plot (c)

Table 1: PCR and PLS models to predict the content of mixtures containing grape seed oil and olive oil

| Oils | Multivariate calibration | Type of spectra | Number of components | R_{cal}^2 | RMSEC | R_{cv}^2 | RMSECV | R_{val}^2 | RMSEP |
|----------------|--------------------------|--------------------------|----------------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Grape seed oil | PCR | Original | 8 | 0.995 | 1.175 | 0.990 | 1.762 | 0.984 | 2.169 |
| | | First derivative | 5 | 0.995 | 1.197 | 0.993 | 1.410 | 0.982 | 2.311 |
| | | Second derivative | 7 | 0.995 | 1.178 | 0.992 | 1.559 | 0.976 | 2.660 |
| | | SNV | 2 | 0.993 | 1.431 | 0.991 | 1.657 | 0.966 | 3.191 |
| | PLS | SG | 8 | 0.995 | 1.180 | 0.989 | 1.771 | 0.984 | 2.159 |
| | | Original | 4 | 0.994 | 1.317 | 0.989 | 1.785 | 0.977 | 2.603 |
| | | First derivative | 3 | 0.996 | 1.088 | 0.994 | 1.383 | 0.982 | 2.342 |
| | | Second derivative | 5 | 0.998 | 0.700 | 0.992 | 1.557 | 0.982 | 2.331 |
| Olive oil | PCR | SNV | 2 | 0.994 | 1.397 | 0.991 | 1.644 | 0.967 | 3.127 |
| | | SG | 4 | 0.994 | 1.320 | 0.989 | 1.788 | 0.978 | 2.591 |
| | | Original | 8 | 0.995 | 1.175 | 0.990 | 1.762 | 0.984 | 2.169 |
| | | First derivative | 5 | 0.995 | 1.197 | 0.993 | 1.410 | 0.982 | 2.311 |
| | PLS | Second derivative | 7 | 0.995 | 1.178 | 0.992 | 1.559 | 0.976 | 2.660 |
| | | SNV | 2 | 0.993 | 1.431 | 0.991 | 1.657 | 0.966 | 3.191 |
| | | SG | 8 | 0.995 | 1.180 | 0.989 | 1.771 | 0.984 | 2.159 |
| | | Original | 4 | 0.994 | 1.317 | 0.989 | 1.785 | 0.977 | 2.603 |
| PLS | First derivative | 3 | 0.996 | 1.088 | 0.994 | 1.383 | 0.982 | 2.341 | |
| | Second derivative | 5 | 0.998 | 0.700 | 0.992 | 1.557 | 0.982 | 2.331 | |
| | SNV | 2 | 0.994 | 1.397 | 0.991 | 1.644 | 0.967 | 3.127 | |
| | SG | 4 | 0.994 | 1.320 | 0.989 | 1.788 | 0.978 | 2.591 | |

Note: Selected model of calibration for each material were marked with bold. PCR: Principal Component Regression; PLS: Partial Least Squares; SNV: Standard Normal Variate; SG: Savitzky-Golay smoothing with polynomial order of 3 and window width of 11 points.




Fig. 3: Plot of grape seed oil (a) and olive oil (b) predictions resulting from the best calibration model




Fig. 4: Output of sPLS-DA processing for grape seed oil authentication, including background prediction plot (a), AUC-ROC plot (b), contribution plot (c), and 3D individual plot (d)

Sample discrimination using sPLS-DA

Sparse Partial Least Squares Discriminant Analysis (sPLS-DA) is a development of Sparse Partial Least Squares (sPLS) regression, which is applied for classification and discrimination purposes [21]. sPLS-DA, which is developed for second derivative spectra accompanied by graphical visualization (fig. 4), is stated to be more effective for the authentication technique in this research [14].

DISCUSSION

The FTIR spectrum of the three oils appears quite similar. However, careful examination of the FTIR spectrum of the sample shows several differences in the peak profile in the area 1743-718 cm^{-1} : OO has a dominant peak at 1160 cm^{-1} while GO has a dominant peak at 1743 cm^{-1} , which marks pulling vibration (stretching vibration) of the carbonyl group (C=O) [22]. The absorption intensity at 2922 and 1743 cm^{-1} is related to the presence of ester groups and aliphatic chains found in triglycerides [10]. Other areas at 1376, 1160, and 1099 cm^{-1} are related to bending and stretching vibrations by the -C-H-C-O-CH₂- and-CH₂ functional groups in fatty acids [23]. This frequency area was further optimized to measure GO levels in OO with the help of multivariate calibration [24].

The main component analysis (fig. 2) used from the original spectrum is presented in the data Scree Plot, Variable Plot, and Individual Plot of the PCA model. Scree plots are used in visual assessments that show how much variation each component or dimension has; variable plots are useful in evaluating the constructing variables of the main components through the angles formed between vectors, while individual plots show individual positioning of data on a two-dimensional display for the first two components [9].

In the fig. 2, the individual plot produces the separation of pure GO, OO, and GO+OO oils. The total variance generated to create a two-dimensional PCA model is 93.8%, which is obtained from Dim1 (69.5%) and Dim2 (24.3%). Since the spectral data were obtained from twelve main peaks to construct the PCA, it is interesting to evaluate the contribution of each original variable. The variable plot shows the load of each variable, as well as its contribution profile. The absorbance numbers at 1743, 1653, 1238, 1099 and 718 cm^{-1} are the main contributors in the construction of the PCA model. In

conclusion, the vibrational bands in these areas provide specific fingerprint properties for GO and OO. The vibration bands close to 1653 and 1099 cm^{-1} are generated from the GO content. Meanwhile, the OO content contributes to the existence of bands close to 1099 and 718 cm^{-1} .

The selected spectrum type is the second derivative spectrum in the PLS model for both GO and OO. In table 1, it can be seen that for R_{int} , the value is high, while for RMSEC, the value is low; this indicates that FTIR spectroscopy combined with the PLS method is precise and accurate for GO and OO quantities [25].

The prediction model generated from a training data set refers to the calibration model in chemometrics. R_{int} and RMSEC values are determined to evaluate this model. The performance of cross-validation as internal validation of the training data set is reviewed using the leave-one-out technique. The R_{cv} and RMSECV values indicate the quality of the model developed when the internal validation model is applied. Considering that the construction of the external validation model comes from an independent data set or test data set, the model's predictive ability is verified quantitatively through the R_{ext} and RMSEP values (fig. 3).

The sPLS-DA model on the background prediction plot provides visualization and discrimination markers for each group. The sample categorization classification is developed using the maximum distance approach model [26]. For the GO, OO, and GO+OO samples, the AUC-ROC plot provides an illustration of the ability of the sPLS-DA model to discriminate. The AUC curve represents the degree of model separation, while the ROC curve represents the probability of discrimination model. GO and OO are separated into another class because the model results show a value of 1 or 100%, while the value for GO+OO is 0.994. These results indicate that the chance of the sPLS-DA model to differentiate GO+OO from the others is 99.4%. The contribution plot presents the contribution of each real variable to constructing the components of the discrimination model. The constructor of the sPLS-DA model with the highest contribution values for GO, OO and GO+OO were 1728, 1737 and 1734 cm^{-1} , respectively. The variable with the highest contribution is produced from a narrow band above 1700 cm^{-1} ; in other words, the stretching vibration of the carbonyl group can be characterized in this area [27].

CONCLUSION

This research developed a rapid detection technique for counterfeiting grape seed face oil using the Fourier Transform Infrared (FTIR) spectroscopy method combined with chemometric techniques. Exploratory data analysis was carried out using the PCA technique, followed by constructing a multivariate calibration model using PLS and PCR, as well as the discrimination using sPLS-DA. The best multivariate calibration model was obtained from second derivative spectra processed using PLS for both GO and OO. The SPLS-DA model was successfully constructed and discriminated with AUC-ROC values of 1.000, 1.000, and 0.994 for GO, OO, and GO+OO, respectively.

FUNDING

This research was funded by Institute for Research and Community Services, Universitas Sanata Dharma with the scheme of a Magister-Doctor Research Grant [Contract No. 019 No. 019 Penel./LPPM-USD/III/2024].

AUTHORS CONTRIBUTIONS

A. P. conducted all sample preparation, chemometrics techniques development, data interpretation and initiated the original draft of the manuscript. W. A. and A. W. carried out the data acquisition using FTIR spectroscopy. E. P. I. and F. D. O. R. conceptualized the research project. F. D. O. R. reviewed and edited the manuscript. All authors have approved the final version of the manuscript.

CONFLICTS OF INTERESTS

All authors have none to declare

REFERENCES

- Barros C, Barros RB. Natural and organic cosmetics: definition and concepts. *Journal of Cosmetology & Trichology*. 2020;6(2):1-9. doi: 10.20944/preprints202005.0374.
- Mursyidi A. The role of chemical analysis in the halal authentication of food and pharmaceutical products. *J Food Pharm Sci*. 2013;1:1-4. doi: 10.14499/jfps.
- Marlina PR, Pradita R, Lucida H, Sudji IR, Salsabila HN, Elida N. Formulation of mesenchymal stem cell secretome as antiaging cream. *Int J App Pharm*. 2023;15(1):45-50. doi: 10.22159/ijap.2023.v15n1.47506.
- Okereke JN, Udebuani AC, Ezeji EU, Ohasi KO, Nnoli MC. Possible health implications associated with cosmetics: a review. *Sci J Public Health*. 2015;3(5-1):58-63. doi: 10.11648/j.sjph.s.2015030501.21.
- Goncalves S, Gaivao I. Natural ingredients common in the trase-os-montes region (portugal) for use in the cosmetic industry: a review about chemical composition and antigenotoxic properties. *Molecules*. 2021;26(17):5255. doi: 10.3390/molecules26175255, PMID 34500687.
- Maru AD, Lahoti SR. Formulation and evaluation of moisturizing cream containing sunflower wax. *Int J Pharm Pharm Sci*. 2018;10(11):54-9. doi: 10.22159/ijpps.2018v10i11.28645.
- Maru AD, Lahoti SR. Formulation and evaluation of ointment containing sunflower wax. *Asian J Pharm Clin Res*. 2019;12(8):115-20. doi: 10.22159/ajpcr.2019.v12i8.33199.
- Astuti F, Saifulah TN, Wulanjati MP, Yosephine AD, Ardianti D. Basil essential oil (*Ocimum basilicum* L.) activities on streptococcus mutans growth biofilm formation and degradation and its stability in microemulsion mouthwash formula. *Int J Pharm Clin Res*. 2016;8(1):26-32. doi: 10.22146/tradmedj.8036.
- Astuti A, Fitri N. Antiaging serum formulation of black pepper (*Piper nigrum* L.) essential oil and antioxidant activity test using the DPPH method. *Asian J Innov Entrep*. 2020;5(1):1-11.
- Sumaiyah LBM, M. Leisyah B. The effect of antioxidant of grapeseed oil as skin anti-aging in nanoemulsion and emulsion preparations. *Rasiyan J Chem*. 2019;12(3):1185-94. doi: 10.31788/RJC.2019.1235337.
- Garavaglia J, Markoski MM, Oliveira A, Marcadenti A. Grape seed oil compounds: biological and chemical actions for health. *Nutr Metab Insights* 2016;9:59-64. doi: 10.4137/NMI.S32910, PMID 27559299.
- Vanstone N, Moore A, Martos P, Neethirajan S. Detection of the adulteration of extra virgin olive oil by near infrared spectroscopy and chemometric techniques. *Food Qual Saf*. 2018;2(4):189-98. doi: 10.1093/fqsafe/fyy018.
- Riyanto AB, Riyanto S, Lukitaningsih E, Rohman A. Analysis of sunflower oil in ternary mixture with grapeseed oil and candlenut oil in the ternary mixture system using FTIR spectroscopy and chemometrics. *Food Res*. 2020;4(5):1726-31. doi: 10.26656/fr.2017.4(5).023.
- Riswanto FD, Windarsih A, Putri DC, Gani MR. An integrated authentication analysis of Citrus aurantium L. essential oil based on FTIR spectroscopy and chemometrics with tuning parameters. *Indonesian J Pharm*. 2023;34(2):205-17. doi: 10.22146/ijp.5225.
- Sahoo MR, Varrier RR, Rajendran A. Analysis and chemical profiling of honey using 1H-NMR spectroscopy FTIR spectroscopy and TLC using various chromogenic reagents for derivatization. *Int J Pharm Pharm Sci*. 2023;15(4):33-8. doi: 10.22159/ijpps.2023v15i4.47292.
- Rohman A, Dzulfianto A, Riswanto FD. The employment of UV-spectroscopy combined with multivariate calibration for analysis of paracetamol propyphenazone and caffeine. *Indonesian J Pharm*. 2017;28(4):191. doi: 10.14499/indonesianjpharm28iss4pp191.
- Huang F, Song H, Gao L, Guang P, Yang X, Li L. Detection of adulteration in Chinese honey using NIR and ATR-FTIR spectral data fusion. *Spectrochim Acta A Mol Biomol Spectrosc*. 2020;235:118297. doi: 10.1016/j.saa.2020.118297, PMID 32248033.
- Rohman A, Che Man YB, Ali ME. The authentication of virgin coconut oil from grape seed oil and soybean oil using FTIR spectroscopy and chemometrics. *Int J App Pharm*. 2019;11(2):259-63. doi: 10.22159/ijap.2019v11i2.31758.
- Bayne CK. Practical guide to chemometrics. *Technometrics*. 1995;37(2):230-1. doi: 10.1080/00401706.1995.10484310.
- Hamdan MR, Ismail Z, Ahmad MN. Assessment of herbal medicines by chemometrics assisted interpretation of FTIR spectra assessment of herbal medicines by chemometrics assisted interpretation of FTIR spectra; 2004.
- Calvini R, Ulrici A, Amigo JM. Practical comparison of sparse methods for classification of arabica and robusta coffee species using near infrared hyperspectral imaging. *Chemom Intell Lab Syst*. 2015;146:503-11. doi: 10.1016/j.chemolab.2015.07.010.
- Nurrulhidayah AP, Man YB, Al Kahtani HA, Rohman A. Application of FTIR spectroscopy coupled with chemometrics for authentication of *Nigella sativa* seed oil. *Spectroscopy*. 2011;25(5):243-50. doi: 10.1155/2011/470986.
- Amir JR, Jamwal R, Kumari S, Dhauhanija AS, Balan B, Singh DK. Application of ATR-FTIR spectroscopy along with regression modelling for the detection of adulteration of virgin coconut oil with paraffin oil. *LWT Food Sci Technol*. 2020;118:2019.108754. doi: 10.1016/j.lwt.2019.108754.
- Rohman A, Man YB. Fourier transform infrared (FTIR) spectroscopy for analysis of extra virgin olive oil adulterated with palm oil. *Food Res Int*. 2010;43(3):886-92. doi: 10.1016/j.foodres.2009.12.006.
- Rohman A, Setyaningrum DL, Riyanto S. FTIR spectroscopy combined with partial least square for analysis of red fruit oil in ternary mixture system. *Int J Spectrosc*. 2014;2014:1-5. doi: 10.1155/2014/785914.
- Rohart F, Gautier B, Singh A, Le Cao KA. mixOmics: an R package for omics feature selection and multiple data integration. *PLoS Comput Biol*. 2017;13(11):e1005752. doi: 10.1371/journal.pcbi.1005752, PMID 29099853.
- Cebi N, Taylan O, Abusurrah M, Sagdic D. Detection of orange essential oil isopropyl myristate and benzyl alcohol in lemon essential oil by FTIR spectroscopy combined with chemometrics. *Foods*. 2020;10(1):1-14. doi: 10.3390/foods10010027, PMID 33374136.