

# Indonesian Journal of Pharmacy



[HOME](#) [ABOUT](#) [PEOPLE](#) [SUBMISSIONS](#) [ISSUE](#) [PREVIOUS WEBSITE](#)



Indonesian Journal of Pharmacy (IJP) is a leading international, peer-reviewed journal in Pharmaceutical Science in Indonesia. The journal is published quarterly online by the Faculty of Pharmacy, Universitas Gadjah Mada. We provide open access of published articles to support greater exchange of scientific knowledge.

IJP was established in 1972 and the online publication of the journal was initiated in 2008. Since 2012, the journal has been published in English by the Faculty of Pharmacy in collaboration with Indonesian Pharmacists Association (IAI). IJP has been accredited by the Directorate General of Higher Education (DGHE) DIKTI No. 30/E/KPT/2018. IJP has also been indexed by SCOPUS since 2018.

ISSN (electronic): 2338-9486  
ISSN (printed) : 2338-9427

## CURRENT ISSUE

**Vol 35 No 3, 2024**



**PUBLISHED:** 2024-09-26

## REVIEW ARTICLE

### **Beta-Blocker in Heart Rate Control and Cardio Protection: The Role of ADRB1 Variants and HCN4 Regulation – A Systematic Review**

Rasmaya Niruri, Zullies Ikawati, Agung Endro Nugroho, Habibie Arifianto  
375–391

 PDF

 Abstract views: 60 |  views: 0

## Clinical Decision Support Systems to Identify Drug-Related Problems in Diabetes Mellitus Patients: A Systematic Review

Niken Larasati, Satibi Satibi, Susi Ari Kristina, Lutfan Lazuardi

392–408



PDF

Abstract views: 102 | PDF views: 0

## RESEARCH ARTICLE

---

### Phytochemical, Nutraceutical and Pharmacological Aspects of the Philippine native *Acalypha angatensis* Blanco, Fl. Filip.

Regina Lourdes Hipol, Hilda Wayas, Florence Mae Bacuyag, Madonna Daquigan, Roland Hipol, Jedida Cabanlong

409–424



PDF

Abstract views: 36 | PDF views: 0

### Isolation of Nutmeg Essential Oil (*Myristica fragrans* houtt) From Aceh Indonesia and Their Antioxidant and Antibacterial Activities

Khairan Khairan, Meutia Faradilla, Binawati Ginting

425–436



PDF

Abstract views: 53 | PDF views: 0

### Gut Microbiota Dynamics and Phenotypic Changes Induced by Tetracycline in *Drosophila melanogaster*

Zhavira Pradiny Saadjad, Sartini Sartini, Rizky Chairatunnisa, Nadila Pratiwi Latada, Mukarram Mudjahid, Firzan Nainu

437–450



PDF

Abstract views: 92 | PDF views: 0

### PyPLIF HIPPOS-aided Construction and Retrospective Validation of Structure-Based Virtual Screening Protocol Targeting VEGFR2

Supanji Supanji, Ayudha Bahana Perdamaian, Titi Marsifah, Riris Istighfari Jenie, Muthi Ikawati, Dewi Kartikawati Paramita, Enade Perdana Istyastono

451–458



PDF

Abstract views: 81 | PDF views: 0

### Comparative Cellular and In Vivo Anti-cancer studies of Doxorubicin Liposomes Prepared with Different Types of Phospholipids

Andang Miatmoko, Devy Maulidya Cahyani, Kumi Kawano, Yoshiyuki Hattori

459–466



PDF

Abstract views: 139 | views: 0

### **Zinc Sulfate and $\alpha$ -tocopherol Supplementation Enhance Reproductive Performance in Male Albino Rats (*Rattus norvegicus*) With Lead Acetate Toxicity**

Ragil Angga Prastiya, Trilas Sardjito, Bodhi Agustono, Faisal Fikri, Amung Logam Saputro, Wurlina Wurlina, Basmala Launa Dewi, Sherly Mega Pratiwi, Aras Prasetyo Nugroho, Ikechukwu Benjamin Moses

467–480



PDF

Abstract views: 31 | views: 0

### **Attenuation of TNF- $\alpha$ and Iron Levels in Renal Hemosiderosis by *Phaleria macrocarpa* (Scheff.) Boerl Extract in a Rat Iron Overload Model**

Nadia Larasinta Heriatmo, Ari Estuningtyas, Vivian Soetikno, Ernie H. Poerwaningsih, Kusmardi Kusmardi

481–490



PDF

Abstract views: 67 | views: 0

### **Physico-Chemical and Structural Characterization of Mixed Natural Polymer Hydrogels Under Simulated Gastrointestinal Fluid**

Agustina Ari Murti Budi Hastuti, Khadijah Zai, Puput Risdanareni, Rahmi Karolina, Irfan Mustafa

491–500



PDF

Abstract views: 46 | views: 0

### **Optimizing the Formula of Polymeric-Based Aripiprazole Nanosuspension Using Response Surface Methodology for Intranasal Drug Delivery**

Nabila Fatin Aisiah, Silvia Surini

501–511



PDF

Abstract views: 39 | views: 0

### **Simultaneous Quantification of Ramipril and Ramiprilat in Drug Formulations: A Clinical LC-MS/MS Study**

Determination of Ramipril and Ramiprilat

Khaled Omari, Ahmed Abu-Awwad, Basel Arafat, Gianina Macovei, TLucretia Boltez, Zaher Abdel Baki, Eyad Mallah, Tawfiq Arafat

512–520





PDF

Abstract views: 60 | views: 0 views: 0

### **Formulation of Lotion with Ethanol Extract of *Curcuma mangga* Val. as UV B Sunscreen and Activity Test using UV-Vis Spectrophotometry**

Nindya Kusumorini, Abdul Karim Zulkarnain  
521-530

 PDF

 Abstract views: 97 |  views: 0

### **The Role of Growth Differentiation Factor-15 in the Diagnosis of Patients with Chronic Angina Undergoing Coronary Angiography**

Aseel Butti, Wassan Abdul Kareem Abbas, Ahmed Yousif Hasan  
531-539



 PDF

 Abstract views: 16 |  views: 0

### **Validation of An HPLC Method for the Determination of Some B-Lactams Antibiotics Using A Green Analytical Technique**

Thoai Dang Nguyen, Anh Tran Thi Phuong Phuong, Ky Truong Quoc  
540-548

 PDF

 Abstract views: 51 |  views: 0


## **SHORT COMMUNICATION**

---

### **Analytical Method Validation and Simultaneous Determination of Ketotifen Fumarate and Cyproheptadine Hydrochloride in Divided Powder by RP-HPLC Method**

Melynia Sintha Dewi, Florentinus Dika Octa Riswanto, Sri Hartati Yuliani  
549-556

 PDF

 Abstract views: 111 |  views: 0

**VIEW ALL ISSUES** [➤](#)

---

Welcome to our new website all readers, authors, and reviewers of the Indonesian Journal of Pharmacy. This website is based on OJS version 3 and has been active since September 1, 2020. We wish the best participation and contribution from all the readers, authors, and reviewers. Thank you.



#### JOURNAL MENU

[Aims & Scope](#)



[Editorial Board](#)[Publication Ethics](#)[Editorial Policies](#)[Instructions for Authors](#)[Article Processing Charge](#)[Peer Review Process](#)[Indexing & Archiving](#)[Journal Statistics](#)[Journal History](#)[Editorial Office](#)[Article In Press](#)[Publication Process](#)[Template Document](#)

## INFORMATION

---

[For Readers](#)[For Authors](#)[For Librarians](#)

## MOST READ LAST WEEK

---

Advances in Quantitative and Qualitative Assessment of Collagen in Cutaneous Wound Healing

👁 641

Formulation and Evaluation of Luliconazole nanosponge gel using Experimental design

👁 496

The Development of Plant-Based Jelly Candy for Chrysanthemum indicum L. Flower Extract and Evaluation of The Antioxidant Activity

👁 414

Antidiabetic Activity of Averrhoa bilimbi L. Fruit Extracts and the Identification of Active Compounds Using LC-MS and In silico Methods

👁 378

Formulation And Optimization Peel-Off Gel Mask with Polyvinyl Alcohol and Gelatin Based Using Factorial Design from Banana Peel Flour (Musa paradisiaca L) As Antioxidant

👁 339

## CURRENT ISSUE

---

ATOM 1.0

RSS 2.0

RSS 1.0

### Editorial Office:

FACULTY OF PHARMACY  
UNIVERSITAS GADJAH MADA  
Jl. Kaliurang Km.4 Sekip Utara  
Yogyakarta 55281



This work is licensed under a [Creative Commons Attribution 2.0 Generic License](#).

Indonesian Journal of Pharmacy is indexed by :

[SCOPUS](#), [DIMENSION](#), [Google Scholar](#), [SINTA](#), [DOAJ](#)

[View My Stats](#)



Platform &  
workflow by  
**OJS / PKP**



[HOME](#) [ABOUT](#) [PEOPLE](#) [SUBMISSIONS](#) [ISSUE](#) [PREVIOUS WEBSITE](#)

[HOME](#) / [Editorial Team](#)

### Editor In Chief

[Dr. Marlyn Dian Laksitorini](#), Faculty of Pharmacy, Universitas Gadjah Mada Indonesia.

### Editorial Board

[Prof. Dr. Veeresh P. Veerapur](#), Sree Siddaganga College of Pharmacy, Pharmaceutical Chemistry Department, India

[Prof. Dr. Lee E. Kirsch](#), University of Iowa, Division of Pharmaceutics and Translational Therapeutics, United States

[Prof. Dr. Jeroen Kool](#), Vrije Universiteit Amsterdam, Division of BioAnalytical Chemistry, Netherlands

[Dr. Saikat Kumar Basu](#), University of Lethbridge, Department of Biological Sciences, Canada

[Dr. Joseph David Francis Tucci](#), La Trobe University, School of Pharmacy and Applied Science, Australia

[Dr. Chuda Chittasupho](#), Srinakharinwirot University, Department of Pharmaceutical Technology, Thailand

[Dr. Montarat Thavorncharoensap](#), Faculty of Pharmacy, Department of Pharmacy, Mahidol University, Thailand

[Dr. Karuna Shanker](#), Central Institute of Medicinal and Aromatic Plants India, Department of Analytical Chemistry, India

[Dr. Jun An](#), Sun Yat-Sen University, Department of Cardiothoracic Surgery, China

[Dr. Mohammed Emamussalehin Choudhury](#), Department of Pharmacology, Bangladesh Agriculture University, Bangladesh

[Dr. Abdul Wahab](#), Department of Pharmacy, Kohat University of Science and Technology (KUST), Pakistan

[Dr. Tony Hadibarata](#), Curtin University Sarawak Malaysia, Department of Environmental Engineering, Malaysia

[Dr. Shahin Gavanji](#), Department of Biotechnology, Faculty of Advanced Sciences and Technologies, University of Isfahan, Isfahan, Iran

[Dr. Muhammad Shahzad Aslam](#), Xiamen University, Malaysia



### JOURNAL MENU

[Aims & Scope](#)

[Editorial Board](#)

[Publication Ethics](#)

[Editorial Policies](#)

[Instructions for Authors](#)

[Article Processing Charge](#)

[Peer Review Process](#)

[Indexing & Archiving](#)

[Journal Statistics](#)

[Journal History](#)

[Editorial Office](#)

[Article In Press](#)

[Publication Process](#)

[Template Document](#)



## INFORMATION

---

[For Readers](#)

[For Authors](#)

[For Librarians](#)

## MOST READ LAST WEEK

---

Advances in Quantitative and Qualitative Assessment of Collagen in Cutaneous Wound Healing

👁 641

Formulation and Evaluation of Luliconazole nanosponge gel using Experimental design

👁 496

The Development of Plant-Based Jelly Candy for Chrysanthemum indicum L. Flower Extract and Evaluation of The Antioxidant Activity

👁 414

Antidiabetic Activity of Averrhoa bilimbi L. Fruit Extracts and the Identification of Active Compounds Using LC-MS and In silico Methods

👁 378

Formulation And Optimization Peel-Off Gel Mask with Polyvinyl Alcohol and Gelatin Based Using Factorial Design from Banana Peel Flour (Musa paradisiaca L) As Antioxidant

👁 339

## CURRENT ISSUE

---

ATOM 1.0

RSS 2.0

RSS 1.0

### Editorial Office:

FACULTY OF PHARMACY  
UNIVERSITAS GADJAH MADA  
Jl. Kaliurang Km.4 Sekip Utara  
Yogyakarta 55281



This work is licensed under a [Creative Commons Attribution 2.0 Generic License](#).

Indonesian Journal of Pharmacy is indexed by :

[SCOPUS](#), [DIMENSION](#), [Google Scholar](#), [SINTA](#), [DOAJ](#)

[View My Stats](#)



Platform &  
workflow by  
**OJS / PKP**

## Analytical Method Validation and Simultaneous Determination of Ketotifen Fumarate and Cyproheptadine Hydrochloride in Divided Powder by RP-HPLC-UV Method

Melynia Sintha Dewi<sup>1</sup>, Florentinus Dika Octa Riswanto<sup>2\*</sup> and Sri Hartati Yuliani<sup>1</sup>

1. Division of Pharmaceutical Technology, Faculty of Pharmacy, Campus III Paingan, Universitas Sanata Dharma, Maguwoharjo, Sleman, Yogyakarta 55282, Indonesia
2. Division of Pharmaceutical Analysis and Medicinal Chemistry, Faculty of Pharmacy, Campus III Paingan, Universitas Sanata Dharma, Maguwoharjo, Sleman, Yogyakarta 55282, Indonesia

### Article Info

Submitted: 26-05-2023

Revised: 27-10-2023

Accepted: 01-11-2023

\*Corresponding author  
Florentinus Dika Octa R

Email:  
dikaocta@usd.ac.id

### ABSTRACT

Ketotifen fumarate and cyproheptadine HCl on divided powder dosage form are the most popular prescription for children at "X" hospital in Yogyakarta. The interest for studying this divided powder was related to the quality test towards human health. If the drug doesn't meet the requirement of good quality drug, it can cause damage to the quality of life. Hence, it was important to develop a validated analytical method to quantify content of active drug from samples. The samples were prepared by a pharmacist at "X" hospital in Wonosari, Yogyakarta. Tablets from each compound were blended and divided by pulverizer into powder bags then were sealed. In this study, a reversed phase HPLC Ultraviolet (RP-HPLC-UV) method was optimized and validated then applied for determining the content of ketotifen fumarate and cyproheptadine HCl in divided powder. The optimal condition for HPLC separation with mobile phase of methanol were 0.04M Na<sub>2</sub>HPO<sub>4</sub> buffer (60:40) and flow rate 0.8 mL/min. According to AOAC International 2019, the developed RP-HPLC-UV method was good and met the requirement of selectivity and sensitivity. The calibration curve formulas of ketotifen fumarate and cyproheptadine hydrochloride were  $y = 27714x - 79111$  ( $r=0.994$ ) and  $y = 26324x + 72581$  ( $r=0.993$ ) in the range of 10.9-54.5 and 10.6-53.0 µg/mL, respectively. Accuracy and intermediate precision were assessed on both standard solution and samples spiked with standard solution of ketotifen fumarate and cyproheptadine hydrochloride. It was found that both standard solution and the spiked samples solution met the requirement of recovery of 80-110% and RSD of less than 7.3% for intraday and interday analysis. Content of ketotifen fumarate and cyproheptadine hydrochloride in sample (n=7) were  $16.574 \pm 0.098$  and  $30.640 \pm 0.035$  µg/mL, respectively. It can be concluded that the RP-HPLC-UV method was successfully validated then applied in the determination of ketotifen fumarate and cyproheptadine HCl in divided powder.

**Keywords:** analytical; cyproheptadine hydrochloride; ketotifen fumarate; HPLC; validation

### INTRODUCTION

Allergic diseases are the most common chronic conditions that affect the quality of patient's life (Kuna *et al.*, 2016). Asthma is the most prevalent chronic respiratory disease worldwide. This disease affecting more than 300 million people of all ethnic groups at all ages (Soriano *et al.*, 2017). It is the most common chronic disease especially in

children, imposing an increasingly consistent burden on the health system (Asher & Pearce, 2014). The recent Global Asthma Study (GAN) phase reported that the global prevalence of asthma was 11% in children aged 6–7 years and 9.1% at 13–14 years (Marcos *et al.*, 2022). Asthma has accounted for missed school days among children in Asia-Pacific (16–61%),



Europe (34–68%) and the United States (43%) and has been affected to lower academic achievement (Triasih *et al.*, 2023). Antihistamines are the most commonly used in the therapy of allergies (Kuna *et al.*, 2016). First and second generation of antihistamines providing effective relief of allergic conditions such as ketotifen fumarate and cyproheptadine hydrochloride (El-Kommos *et al.*, 2015). Ketotifen fumarate has been administered to prevent allergic disease such as asthma and anaphylactic reaction (Moreno *et al.*, 2020). Cyproheptadine has been used in chronic allergic and pruritic conditions (Chakraborty, 2019; Vardanyan, 2017). As the most popular prescription for children at “X” hospital in Yogyakarta, ketotifen fumarate and cyproheptadine hydrochloride were chosen to be analyzed in this study. It was stated in previous studies that the prescription of ketotifen fumarate and cyproheptadine hydrochloride from “X” hospital in Yogyakarta were frequently prescribed with the total amount of 87 from 291 prescription (the five most prescribed formula) (Yuliani *et al.*, 2020). However, there were limited studies reported the analytical method development to ensure the quality of compounded drugs, especially divided powder. Hence, it is important to develop a validated analytical method to analyze ketotifen fumarate and cyproheptadine hydrochloride content in divided powder to ensure the quality and stability of this compounded drug.

High performance liquid chromatography (HPLC) method was commonly applied for the separation, quantification, and identification of compounds present in a mixture (Basharat *et al.*, 2021). Several compounded preparation analysis including ketotifen fumarate or cyproheptadine hydrochloride has been using this HPLC method (Abdelrahman *et al.*, 2021; Aldewachi & Omar, 2022; Gebretsadik *et al.*, 2023; Kabra *et al.*, 2014; Kesharbai, 2014; Mohammed *et al.*, 2021; Muralidharan *et al.*, 2012; Peikova *et al.*, 2022; Pozharani *et al.*, 2022; Prasad *et al.*, 2022; Rajput & Sathe, 2019; Sahai & Devanna, 2021; Sethiya & Rathore, 2022; Tokumura *et al.*, 2018; Yilmaz *et al.*, 2022). However, studies analyzing ketotifen fumarate and cyproheptadine hydrochloride in divided powder combination are still limited. An analytical method validation should be performed to ensure that the analytical method is applicable to achieve the appropriate results (Mennickent & de Diego, 2019). The aim of this study was to validate

the analytical method of RP-HPLC-UV and apply the validated method to quantify content of ketotifen fumarate and cyproheptadine hydrochloride in divided powder. The RP-HPLC-UV method was validated in several validation parameters including selectivity, linearity and range, precision, accuracy, sensitivity such as LOD and LOQ according to the AOAC Official Methods of Analysis (AOAC International, 2019).

## **MATERIALS AND METHODS**

### **Materials and Chemicals**

The reference standards of ketotifen fumarate and cyproheptadine hydrochloride were purchased from Indonesian Food and Drug Authority (FDA). The solvent used in this research was methanol for liquid chromatography grade (LiChrosolv®). The sodium hydrogen phosphate for buffer solution was purchased from Merck Millipore. Ketotifen fumarate and cyproheptadine hydrochloride tablets were obtained from a drugstore in Yogyakarta, Indonesia.

### **Instrumentation and Software**

A system of HPLC Shimadzu LC-2010 with UV/Vis detector equipped with C18 column Luna Phenomenex® (250×4.6 mm, i.d. 5 µm) endcapped was used in this study. Analytical balance OHAUS® PA 413, Gast® vacuum pump, Retsch® ultrasonicator, sterile syringe filter with a 0.45 µm pore size hydrophilic PTFE membrane (Merck Millipore), and a set of Socorex® micropipettes were also used in this study. Data from HPLC instrument was subsequently analyze using Excel (Microsoft Inc., USA) for generating linear calibration model and other data calculations.

### **HPLC Conditions**

An isocratic RP-HPLC-UV method was applied with the mobile phase of methanol:0.04 M Na<sub>2</sub>HPO<sub>4</sub> buffer pH 4.8 (60:40) and flow rate was adjusted to 0.8 mL/min. Column temperature was set to default 26.5°C. Wavelength detection was optimized and adjusted at 287 nm. Volume injection for each sample was 10 µL.

### **Preparation of standard solution**

Accurate weights of 10.6 and 10.9 mg for ketotifen fumarate and cyproheptadine hydrochloride, respectively, were put into different 10 mL volumetric flask. The concentration of ketotifen fumarate was 1060 ppm and cyproheptadine hydrochloride 1090 ppm.

### Preparation of sample solution

The preparation of sample solution was adapted from Kesharbai, 2014. An accurate weight of 225 mg of divided powder sample were placed in beaker glass and diluted by methanol into the volume of 10.0 mL volumetric flask. The working sample solution was prepared by transferring a 1.5 mL sample diluted solution into a 5 mL volumetric flask followed by dilution with methanol to volume. This solution was filtered using a sterile PTFE membrane before injection into the HPLC system.

### Preparation of spiked sample solution

An accurate weighed of 225 mg ketotifen fumarate and cyproheptadine hydrochloride divided powder were placed in beaker glass and diluted by methanol into the volume of 10.0 mL volumetric flask, respectively. The spiked sample solution was prepared by transferring 85,95,105  $\mu$ L cyproheptadine HCl and 110, 130,150  $\mu$ L ketotifen fumarate solution into a 5 mL volumetric flask followed by dilution with methanol to volume. This solution was filtered using a sterile PTFE membrane before injection into the HPLC system.




Figure 1. HPLC optimization using three variations of mobile phase for mixture standard solution of ketotifen fumarate (Ket) and cyproheptadine hydrochloride (Cyp). Wavelength detection at 287 nm. Notes: A: mobile phase = methanol : redistilled water : acetic acid (45:40:15), flow rate = 0.8 mL/min; B: mobile phase = buffer : methanol (65:35); flow rate = 0.8 mL/min; and C: mobile phase = buffer : methanol (60:40), flow rate = 0.8 mL/min.

### Optimization of Analytical Method

Optimization of the chromatographic separation was performed by variation of mobile phase. Wavelength detection was set to 287 nm in which two analytes were intersecting wavelengths. The volume injection for each run was 10  $\mu$ L. The composition of each run were methanol : redistilled water : acetic acid (45:40:15), buffer : methanol (35:65), buffer : methanol (40:60) with the same flow rate 0.8 mL/min (Figure 1).

### Validation of Analytical Method

The HPLC method was validated for selectivity, linearity and range, precision, accuracy, sensitivity such as LOD and LOQ according to the AOAC Official Methods of Analysis (AOAC International, 2019).




Figure 2. Chemical structures and UV spectra profiles of ketotifen fumarate (a) and cyproheptadine hydrochloride (b) in methanol solvent

## RESULTS AND DISCUSSION

### Determination of Wavelength Detection

Wavelength detection should be stated at the initial stage of HPLC-UV method development. It was important to choose an optimal wavelength for detecting several compounds in a mixture matrix (Figure 2). In this study, detection of the analytes was performed at 287 nm. At this wavelength, ketotifen fumarate and cyproheptadine hydrochloride can be optimally detected. The extensive overlapping spectra of analytes may become the limitation while applying the UV spectroscopy for a simultaneous quantification (Dzulianto *et al.*, 2017). However, by applying HPLC method, these two analytes can be detected simultaneously at the same wavelength and resulted separation profile for each analyte due to the interaction of the analyte itself both with mobile and stationary phase at the various interaction strength (Snyder *et al.*, 2010).

Table I. Results of system suitability test (n=6)

Analytes	Retention Time		Area	
	Mean	RSD (%)	Mean	RSD (%)
Ketotifen Fumarate	3.475	1.551	271556.8	0.973
Cyproheptadine Hydrochloride	8.373	1.114	584726	0.887




Figure 3. HPLC separation profiles ketotifen fumarate and cyproheptadine hydrochloride, divided powder sample, and divided powder sample with standard addition of ketotifen fumarate and cyproheptadine hydrochloride. Mobile phase: methanol:buffer (60:40 v/v). Column: C18 Luna Phenomenex® (250×4.6 mm, i.d. 5 µm) endcapped. Flowrate: 0.8 mL/min. Column temperature: 26.5 °C. Wavelength detection at 287 nm. Volume injection: 10 µL

**System Suitability Test**

The system suitability test is necessary to verify and ensure the continued performance of systems and analysis methods. HPLC separation properties are retention time and area. The relative standard deviation of retention time and area met the acceptance criteria (RSD ≤ 2%) (Snyder *et al.*, 2010) (Table I).

**Selectivity**

Selectivity is a method to quantify the analyte from interference. This method can be held by a peak purity test. The peak should be single without another analyte. This HPLC condition met the criteria of resolution > 1.5, tailing factor < 2.0, theoretical plate number > 2000, and resulted minimum deviation (≤ 2%) both for retention

time and peak area for each analyzed compound (AOAC International, 2019). The HPLC separation profiles of ketotifen fumarate and cyproheptadine hydrochloride, divided powder sample, and divided powder sample spiked with standard of ketotifen fumarate and cyproheptadine hydrochloride (Figure 3). Retention time of ketotifen fumarate and cyproheptadine hydrochloride were 3.444 and 7.981 min, respectively.

**Linearity and Range**

The data of peak areas versus concentration of standard solutions series from each compound were plotted to generate multiple point calibration curves for quantifying the content of ketotifen fumarate and cyproheptadine hydrochloride.

Table II. Accuracy and Precision Study of Standard Ketotifen Fumarate and Cyproheptadine Hydrochloride Solutions (n=3).

Intraday						
Compounds	Level	Added amount (µg/mL)	Found amount (µg/mL)	SD	RSD (%)	Recovery (%)
Ketotifen Fumarate	Low	10.9	12.6	0.02	0.17	86.34
	Medium	21.8	21.2	0.06	0.31	102.67
	High	32.7	32.7	0.55	1.69	99.89
Cyproheptadine Hydrochloride	Low	10.6	12.2	0.03	0.27	86.29
	Medium	21.2	20.7	0.03	0.15	102.19
	High	31.8	31.6	0.05	0.17	100.52
Interday						
Ketotifen Fumarate	Low	10.9	12.7	0.08	0.65	85.64
	Medium	21.8	21.2	0.08	0.41	102.48
	High	32.7	32.5	0.22	0.70	100.62
Cyproheptadine Hydrochloride	Low	10.6	12.3	0.05	0.47	85.55
	Medium	21.2	20.8	0.12	0.58	101.67
	High	31.8	31.9	0.56	1.74	99.65

Table III. Accuracy and Precision Study of Spiked Samples (n=3)

Intraday						
Compounds	Level	Added amount (µg/mL)	Found amount (µg/mL)	SD	RSD (%)	Recovery (%)
Ketotifen Fumarate	Low	15	15.52	0.13	0.86	103.48
	Medium	13	11.35	0.18	1.59	87.34
	High	11	11.19	0.15	1.37	101.73
Cyproheptadine Hydrochloride	Low	21	22.05	0.21	0.96	105.04
	Medium	19	20.90	0.29	1.42	110.01
	High	17	17.50	1.69	0.86	102.95
Interday						
Ketotifen Fumarate	Low	15	15.58	0.15	0.99	103.88
	Medium	13	11.36	0.17	1.50	87.44
	High	11	10.49	0.40	3.90	95.41
Cyproheptadine Hydrochloride	Low	21	22.08	0.17	0.77	105.17
	Medium	19	20.8	0.26	1.24	109.60
	High	17	17.71	0.24	1.38	104.19

Table IV. Results of Determination of Ketotifen Fumarate and Cyproheptadine Hydrochloride in Divided Powder

Replication	Analyte concentration (µg/mL)	
	Ketotifen Fumarate	Cyproheptadine Hydrochloride
1	16.56	30.62
2	16.56	30.57
3	16.60	30.62
4	16.64	30.64
5	16.36	30.65
6	16.63	30.68
7	16.63	30.66
Mean (µg/mL)	16.57	30.64
SD	0.09	0.03
RSD (%)	0.59	0.11



Calibration curve equation of ketotifen fumarate and cyproheptadine hydrochloride were  $y = 27714x - 79111$  ( $r=0.9945$ ) and  $y = 26324x + 72581$  ( $r=0.9935$ ), respectively. These curves were linear in the range of 10.9-54.5 and 10.6-53.0  $\mu\text{g/mL}$  for ketotifen fumarate and cyproheptadine hydrochloride, respectively.

#### **Limit of Detection and Limit of Quantification**

LOD and LOQ as the parameters of sensitivity were determined using the standard deviation approach (Miller *et al.*, 2018). LOD for ketotifen fumarate and cyproheptadine hydrochloride were 2.889 and 2.615  $\mu\text{g/mL}$ , respectively. LOQ for ketotifen fumarate and cyproheptadine hydrochloride were 9.6312 and 8.7196  $\mu\text{g/mL}$ , respectively.

#### **Accuracy and Precision of Standard Solutions**

Accuracy and intermediate precision of standard solution were assessed by determining concentration of ketotifen fumarate and cyproheptadine hydrochloride at three concentration levels (low, medium, high). Each concentration level was analyzed triplicate within three different days (interday) and at the same day (intraday). It was found that the developed RP-HPLC-UV method met the acceptance criteria of  $\text{RSD} < 7.3\%$  and recovery in the range of 80-110% (AOAC International, 2019) (Table II). It can be stated that this analytical method was accurate and precise for analyzing standard solution mixture of standard ketotifen fumarate and cyproheptadine hydrochloride both for intraday and interday evaluations.

#### **Accuracy and Precision of Spiked Samples**

Accuracy and intermediate precision of spiked samples were also performed in this study. Blank sample solutions were added by standard solution of ketotifen fumarate and cyproheptadine hydrochloride at three concentration levels. Each concentration level was analyzed triplicate within three different days (interday). Table III presented the results of accuracy and precision study of sample solutions with the addition of ketotifen fumarate and cyproheptadine hydrochloride standards. It was found that the developed RP-HPLC-UV method met the acceptance criteria of  $\text{RSD} < 7.3\%$  and recovery in the range of 80-110% (AOAC International, 2019). It can be stated that this analytical method was accurate and precise for analyzing divided powder samples with the

presence of standard ketotifen fumarate and cyproheptadine hydrochloride both for intraday and interday evaluations.

#### **Determination of Ketotifen Fumarate and Cyproheptadine Hydrochloride in Divided Powder**

The validated RP-HPLC-UV method was applied for determining content of ketotifen fumarate and cyproheptadine hydrochloride in divided powder samples. Table IV presented the results of determination of ketotifen fumarate and cyproheptadine hydrochloride in divided powder sample. The precision of method was assessed and proven by the percentage of RSD less than 7.3% in seven replicates of determination (AOAC International, 2019).

#### **CONCLUSION**

A validated isocratic RP-HPLC-UV method was successfully developed. This method was selective, linear, sensitive, accurate, and precise to quantitatively analyze ketotifen fumarate and cyproheptadine hydrochloride in different samples of divided powder combination to provide contextual information in hospital and drugstore. This method could be applied for routine analysis quality of drug in Indonesia. The good quality of drugs can directly impact to patient's health.

#### **ACKNOWLEDGMENTS**

This research was financially supported by "Program Penelitian Skema Magister Doktor" from the Institute for Research and Development of Society, Sanata Dharma University (No.12 Penel./LPPM-USD/II/2023).

#### **CONFLICT OF INTEREST**

The authors declare no conflict of interest.

#### **REFERENCES**

- Abdelrahman, M. M., Abdelaleem, E. A., Ali, N. W., & Emam, R. A. (2021). Development and validation of stability indicating high-performance liquid chromatographic method for determination of cyproheptadine hydrochloride, its impurity and degradation product. *Journal of Chromatographic Science*, 59(2), 128-133. <https://doi.org/10.1093/chromsci/bmaa085>
- Aldewachi, H., & Omar, T. A. (2022). Development of HPLC Method for Simultaneous Determination of Ibuprofen and

- Chlorpheniramine Maleate. *Scientia Pharmaceutica*, 90(53), 1-9. <https://doi.org/10.3390/scipharm90030053>
- AOAC International. (2019). *Official Methods Of Analysis*.
- Asher, I., & Pearce, N. (2014). Global burden of asthma among children. *International Journal of Tuberculosis and Lung Disease*, 18(11), 1269-1278. <https://doi.org/10.5588/ijtld.14.0170>
- Basharat, R., Kotra, V., Loong, L. ., Mathews, A., Kanakal, M. ., & Devi, C. B. . (2021). Ultra Performance Liquid Chromatography (Mini-Review) RABIA. *Oriental Journal of Chemistry*, 37(4), 847-857. <https://doi.org/10.13005/ojc/370411>
- Chakraborty, A. (2019). Role of Cyproheptadine in Various Psychiatric Conditions. *Indian Journal of Private Psychiatry*, 13(2), 58-61. <https://doi.org/10.5005/jp-journals-10067-0034>
- Dzulianto, A., Riswanto, F. D. O., & Rohman, A. (2017). The employment of UV-spectroscopy combined with multivariate calibration for analysis of paracetamol, propyphenazone and caffeine. *Indonesian Journal of Pharmacy*, 28(4), 191-197. <https://doi.org/10.14499/indonesianjpharm28iss4pp191>
- El-Kommos, M. E., El-Gizawy, S. M., Atia, N. N., & Hosny, N. M. (2015). Analysis of commonly prescribed non-sedating antihistamines. *Analytical Chemistry Research*, 3, 1-12. <https://doi.org/10.1016/j.ancr.2014.11.003>
- Gebretsadik, H., Kahsay, G., Eticha, T., & Gebretsadikan, T. (2023). A Validated New RP-HPLC Method for Simultaneous Determination of Amoxicillin, Ampicillin and Cloxacillin in Pharmaceutical Formulations. *Acta Chromatographica*, 35(2), 193-203. <https://doi.org/10.1556/1326.2022.01043>
- Kabra, P. V., Nargund, L. V. G., & Murthy, M. S. (2014). Development and validation of a stability-indicating LC-UV method for simultaneous determination of ketotifen and cetirizine in pharmaceutical dosage forms. *Tropical Journal of Pharmaceutical Research*, 13(7), 1141-1147. <https://doi.org/10.4314/tjpr.v13i7.19>
- Kesharbai, C. N. (2014). Rp-Hplc Method Development and Validation of Ketotifen Fumarate and Theophylline in Bulk and Combined Dosage Form. *International Research Journal of Pharmacy*, 5(5), 407-411. <https://doi.org/10.7897/2230-8407.050585>
- Kuna, P., Jurkiewicz, D., Czarnecka-Operacz, M. M., Pawliczak, R., Woron, J., Moniuszko, M., & Emeryk, A. (2016). The role and choice criteria of antihistamines in allergy management - expert opinion. *Postepy Dermatologii i Alergologii*, 33(6), 397-410. <https://doi.org/10.5114/pdia.2016.63942>
- Marcos, L. G., Asher, M. I., Pearce, N., Ellwood, E., Bissell, K., Chiang, C. Y., Sony, A. E., & Ellwood, P. (2022). The burden of asthma, hay fever and eczema in children in 25 countries: GAN Phase I study. *The European Respiratory Journal*, 60(3). <https://doi.org/10.1183/13993003.02286-2021>
- Mennickent, S., & de Diego, M. (2019). Analytical Method Validation as the First Step in Drug Quality Control. *Quality Management and Quality Control - New Trends and Developments*, 1-21. <https://doi.org/10.5772/intechopen.82826>
- Miller, J. N., Miller, J. C., & Miller, R. D. (2018). *Statistics and Chemometrics for Analytical Chemistry* (Seventh Ed). Pearson Education Limited.
- Mohammed, O. J., Hamzah, M. J., & Saeed, A. M. (2021). RP - HPLC Method Validation for Simultaneous Estimation of Paracetamol and Caffeine in Formulating Pharmaceutical Form. *Research J. Pharm and Tech.*, 14(9), 4743-4748. <https://doi.org/10.52711/0974-360X.2021.00825>
- Moreno, D., Sobarzo, C.M., Lustig, L., Pena, M.G.R., Guazzone, V. A. (2020). Effect of ketotifen fumarate on experimental autoimmune orchitis and torsion of the spermatic cord. *Asian Journal of Andrology*, 22(1), 112-117. <https://doi.org/10.4103/aja.aja>
- Muralidharan, S., Han, L. B., Ming, J. L. Y., Kartigayam, S., & Dhanaraj, S. A. (2012). Simple and accurate estimation of ketotifen fumarate by RP-HPLC. *International Journal of Pharmaceutical, Chemical and Biological Sciences*, 2(3), 392-396.
- Peikova, L., Tzankova, D., Smerikarova, M., Balkanski, S., & Zlatkov, A. (2022). Development of RP-HPLC Methods for the Analysis of Dexamethasone and Levofloxacin Alone and in Combinations



- used in the Therapy of Covid-19. *Pharmacia*, 69(4), 1075–1080. <https://doi.org/10.3897/pharmacia.69.e97779>
- Pozharani, L. B., Burgaz, E. ., & Baloglu, E. (2022). Development and Validation of an RP-HPLC Method for Simultaneous Determination of Curcumin and Metronidazole in Combined Dosage Form. *Istanbul J Pharm*, 52(3), 250–257. <https://doi.org/10.26650/IstanbulJPharm.2022.1005453>
- Prasad, P. R., Karunakar, D., & Devi, B. R. (2022). Analytical Method Development and Validation of Allopurinol and Related Substances by using RP-HPLC in Bulk Form. *International Journal of Health Sciences*, 6(2), 10447–10470. <https://doi.org/10.53730/ijhs.v6nS2.7809>
- Rajput, S. J., & Sathe, M. A. (2019). New bioanalytical HPLC method for the determination of cyproheptadine hydrochloride in human plasma and its application to rat pharmacokinetic study. *Indian Journal of Pharmaceutical Education and Research*, 53(3), S338–S346. <https://doi.org/10.5530/ijper.53.3s.104>
- Sahai, M., & Devanna, N. (2021). Validated Stability Indicating HPLC Approach for Quantifying Tricholine Citrate and Cyproheptadine Simultaneously in Syrup Forms. *International Journal of Applied Pharmaceutics*, 13(3), 207–213. <https://doi.org/10.22159/ijap.2021v13i3.40871>
- Sethiya, A., & Rathore, R. P. . (2022). RP-HPLC Method Development and Validation for the Estimation of Artemisinin in Bulk Drug. *International Journal of Health Sciences*, 6(2), 14037–14045. <https://doi.org/10.53730/ijhs.v6nS2.8684>
- Snyder, L., Kirkland, J.J., Dolan, J. . (2010). Introduction to Modern Liquid Chromatography, Third Edition. In *Journal of the American Society for Mass Spectrometry* (Vol. 22, Issue 1). <https://doi.org/10.1007/s13361-010-0021-8>
- Soriano, J. B., Abajobir, A. A., Abate, K. H., Abera, S. F., Agrawal, A., Ahmed, M. B., Aichour, A. N., Aichour, I., Eddine Aichour, M. T., Alam, K., Alam, N., Alkaabi, J. M., Al-Maskari, F., Alvis-Guzman, N., Amberbir, A., Amoako, Y. A., Ansha, M. G., Antó, J. M., Asayesh, H., ... Vos, T. (2017). Global, regional, and national deaths, prevalence, disability-adjusted life years, and years lived with disability for chronic obstructive pulmonary disease and asthma, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *The Lancet Respiratory Medicine*, 5(9), 691–706. [https://doi.org/10.1016/S2213-2600\(17\)30293-X](https://doi.org/10.1016/S2213-2600(17)30293-X)
- Tokumura, T., Yasumoto, A., & Kurita, T. (2018). A Validated HPLC Ketotifen Fumarate Assay Method for Cleaning Validation on an Automatic Packaging Machine. *Scholars Academic Journal of Pharmacy (SAJP)*, 460–463. <https://doi.org/10.21276/sajp.2018.7.11.1>
- Triasih, R., Setyowireni, D., Nurani, N., & Setyati, A. (2023). Prevalence, Management, and Risk Factor of Asthma Among School-Age Children in Yogyakarta, Indonesia. *Journal of Asthma and Allergy*, 16(January), 105–106. <https://doi.org/10.2147/JAA.S404674>
- Vardanyan, R. (2017). Piperidin-4-Ylidene Substituted Tricyclic Compounds. In *Piperidine-Based Drug Discovery*. <https://doi.org/10.1016/b978-0-12-805157-3.00006-5>
- Yilmaz, B., Bayrak, B., & Kadioglu, Y. (2022). Development and Validation of HPLC Method for the Determination of Flurbiprofen in Pharmaceutical Preparations. *International Journal of PharmATA*, 2(3), 72–81.
- Yuliani, S. H., Putri, D. C. A., Widayati, A., & Abiyoga, B. (2020). Compounding practice in a developing country: A case study of divided powder in Indonesia. *Research Journal of Pharmacy and Technology*, 13(12), 6231–6237. <https://doi.org/10.5958/0974-360X.2020.01086.0>