

# Journal of Research in Pharmacy

An international open-access journal of pharmacy and pharmaceutical sciences  
Formerly published as Marmara Pharmaceutical Journal

(<https://jrespharm.com/>)

**JRP**  
ISSN : 1309-0801



[Current Issue \(https://jrespharm.com/content.php?id=102\)](https://jrespharm.com/content.php?id=102)

[Articles in Press \(https://jrespharm.com/content.php?id=46\)](https://jrespharm.com/content.php?id=46)

[Archive \(https://jrespharm.com/archive.php\)](https://jrespharm.com/archive.php)

[Search \(https://jrespharm.com/search.php\)](https://jrespharm.com/search.php)

[Author Benefits \(https://jrespharm.com/static.php?id=11\)](https://jrespharm.com/static.php?id=11)

## Editor-in-Chief

Hatice Kübra Elçioğlu

## Vice Editors

Levent Kabasakal

Esra Tatar

## Online ISSN

2630-6344

## Publisher

Marmara University

## Frequency

Bimonthly (Six issues / year)

## Abbreviation

J.Res.Pharm.

## Former Name

Marmara Pharmaceutical Journal

Other Issues :

[◀ Previous \(content.php?id=53\)](https://jrespharm.com/content.php?id=53)

[Next ▶ \(content.php?id=54\)](https://jrespharm.com/content.php?id=54)

Journal of Research in Pharmacy

Articles in Press

[🔍](#) Articles (22)

Articles

[Clear](#)

[View Selected Abstracts](#)

**Development of natural acne mask formulations from rose pomace waste (abstract.php?id=1770) Articles**

Pages 001-010

Ahmet Arif KURT, Bashar IBRAHIM

DOI : 10.29228/jrp.2022.00

[Abstract \(abstract.php?lang=en&id=1770\)](#)

[Full Text PDF \(pdf.php?&id=1770\)](#)

[Similar Articles \(similar.php?&id=1770\)](#)

[E-mail to Author \(mailto:basharibrahim@sdu.edu.tr\)](mailto:basharibrahim@sdu.edu.tr)

**The impact of carvedilol on level of interleukin-6, superoxide dismutase, elastin levels and epidermal thickness in experimentally aging induced mice model (abstract.php?id=1771) Articles**

Pages 011-019

Elaf Mahmood SHIHAB, Haitham Mahmood KADHIM, Samer Salim SHAHOOTH

DOI : 10.29228/jrp.2022.00

[Abstract \(abstract.php?lang=en&id=1771\)](#)

[Full Text PDF \(pdf.php?&id=1771\)](#)

[Similar Articles \(similar.php?&id=1771\)](#)

[E-mail to Author \(mailto:Elaf.mahmood@esraa.edu.iq\)](mailto:Elaf.mahmood@esraa.edu.iq)

**The effect of vitamin D and paricalcitol on protein disulfide isomerase (abstract.php?id=1772) Articles**

Pages 020-029

Muhammed Murat KÖKSAL, Turgut ŞEKERLER, Azize ŞENER

DOI : 10.29228/jrp.2022.00

[Abstract \(abstract.php?lang=en&id=1772\)](#)

[Full Text PDF \(pdf.php?&id=1772\)](#)

[Similar Articles \(similar.php?&id=1772\)](#)

[E-mail to Author \(mailto:turgut.sekerler@marmara.edu.tr com\)](mailto:turgut.sekerler@marmara.edu.tr)

**Antioxidant activity, chemical composition, and therapeutic activity of essential oils from certain Lamiaceae species in Türkiye (abstract.php?id=1773) Articles**

Pages 030-051

DOI : 10.29228/jrp.2022.00

[Abstract \(abstract.php?lang=en&id=1773\)](#)

[Full Text PDF \(pdf.php?&id=1773\)](#)

[Similar Articles \(similar.php?&id=1773\)](#)

[E-mail to Author \(mailto:bgurbuz@marmara.edu.tr\)](mailto:bgurbuz@marmara.edu.tr)

**Formulation and characterisation of curcumin loaded PLGA-Tf nanoparticle for increasing the availability of drug in the brain for the management of parkinson's disease (abstract.php?id=1774) Articles**

*Pages 052-064*

Deepika VERMA,Kumud UPADHYAYA

DOI : 10.29228/jrp.2022.00

[Abstract \(abstract.php?lang=en&id=1774\)](#)

[Full Text PDF \(pdf.php?&id=1774\)](#)

[Similar Articles \(similar.php?&id=1774\)](#)

[E-mail to Author \(mailto:deepikadlverma@gmail.com\)](mailto:deepikadlverma@gmail.com)

**Production and intracellular trafficking of SARS CoV-2 spike protein in insect cells infected with recombinant baculovirus (abstract.php?id=1775) Articles**

*Pages 065-074*

Kadir TURAN

DOI : 10.29228/jrp.2022.00

[Abstract \(abstract.php?lang=en&id=1775\)](#)

[Full Text PDF \(pdf.php?&id=1775\)](#)

[Similar Articles \(similar.php?&id=1775\)](#)

[E-mail to Author \(mailto:kadirturan@marmara.edu.tr\)](mailto:kadirturan@marmara.edu.tr)

**Determination and data analysis for Pb, Ni, Fe, Mn and Co elements in serum of PCOS patients (abstract.php?id=1776) Articles**

*Pages 075-080*

Salam A.H. AL-AMERI,Amer Hasan ABDULLAH,Muntadher H. CHALLOOB

DOI : 10.29228/jrp.2022.00

[Abstract \(abstract.php?lang=en&id=1776\)](#)

[Full Text PDF \(pdf.php?&id=1776\)](#)

[Similar Articles \(similar.php?&id=1776\)](#)

[E-mail to Author \(mailto:muntadther.hussein@uomustansiriyah.edu.iq\)](mailto:muntadther.hussein@uomustansiriyah.edu.iq)

**The effect of the ethanolic extract of *Laportea decumana* (Roxb.) Wedd. on the inflammatory, proliferative and maturation stages of wound healing of an acute injury using a rat model (abstract.php?id=1777) Articles**

*Pages 081-090*

Dwi Novrianty BUSAERI, Yulia Yusrini DJABIR, Rina AGUSTINA, Gemini ALAM, Djulfikri MEWAR, Syaadatun NADIAH

DOI : 10.29228/jrp.2022.00

[Abstract \(abstract.php?lang=en&id=1777\)](#)

[Full Text PDF \(pdf.php?&id=1777\)](#)

[Similar Articles \(similar.php?&id=1777\)](#)

[E-mail to Author \(mailto:yulia.yusrini@unhas.ac.id\)](mailto:yulia.yusrini@unhas.ac.id)

**Optimisation and validation of end-point PCR for the detection of porcine DNA in anti-ageing cream products containing collagen (abstract.php?id=1778) Articles**

*Pages 091-102*

Novi YANTIH, Era WIDIANINGSIH, Sri SURATI, Andi ASNAYANTI

DOI : 10.29228/jrp.2022.00

[Abstract \(abstract.php?lang=en&id=1778\)](#)

[Full Text PDF \(pdf.php?&id=1778\)](#)

[Similar Articles \(similar.php?&id=1778\)](#)

[E-mail to Author \(mailto:yantih.novi@univpancasila.ac.id\)](mailto:yantih.novi@univpancasila.ac.id)

**Evaluation of anti-ulcer potential of *Viola odorata* extract by in-vitro models and ethanol-induced gastric ulcer in rats (abstract.php?id=1779) Articles**

*Pages 103-114*

Jignesh KANSAGRA, Tejas GANATRA

DOI : 10.29228/jrp.2022.00

[Abstract \(abstract.php?lang=en&id=1779\)](#)

[Full Text PDF \(pdf.php?&id=1779\)](#)

[Similar Articles \(similar.php?&id=1779\)](#)

[E-mail to Author \(mailto:jigneshk218@gmail.com\)](mailto:jigneshk218@gmail.com)

**Development and optimization of itraconazole-loaded topical gel using DESM technique (abstract.php?id=1780) Articles**

*Pages 115-122*

Biswajit BANERJEE, Bikram GHOSH, Tripti SHARMA, Arijit MONDAL, Ipsa PADHY

DOI : 10.29228/jrp.2022.00

[Abstract \(abstract.php?lang=en&id=1780\)](#)

[Full Text PDF \(pdf.php?&id=1780\)](#)

[Similar Articles \(similar.php?&id=1780\)](#)

[E-mail to Author \(mailto:biswajit.banerjee.pharm@gmail.com\)](mailto:biswajit.banerjee.pharm@gmail.com)

**Design optimization and evaluation of polyvinyl alcohol based oral thin film of apixaban (abstract.php?id=1781) Articles**

*Pages 123-136*

Mohammad Adnan RAZA, Abhishek KANUGO

[Abstract \(abstract.php?lang=en&id=1781\)](#)

[Full Text PDF \(pdf.php?&id=1781\)](#)

[Similar Articles \(similar.php?&id=1781\)](#)

[E-mail to Author \(mailto:abhi.kanugo09@gmail.com\)](mailto:abhi.kanugo09@gmail.com)

**Analyzing genetic and epigenetic HORMAD alterations in breast cancer resistance and metastatic events (abstract.php?id=1782) Articles**

*Pages 137-150*

Adam HERMAWAN, Herwandhani PUTRI

DOI : 10.29228/jrp.2022.00

[Abstract \(abstract.php?lang=en&id=1782\)](#)

[Full Text PDF \(pdf.php?&id=1782\)](#)

[Similar Articles \(similar.php?&id=1782\)](#)

[E-mail to Author \(mailto:adam\\_apt@ugm.ac.id\)](mailto:adam_apt@ugm.ac.id)

**Chemometrics-assisted fingerprinting profiling of Mangosteen extracts from Sulawesi island using Fourier Transform Infrared (FTIR) Spectroscopy (abstract.php?id=1783) Articles**

*Pages 151-158*

Budiman YASIR, Muhammad AZWAR AR, Asril BURHAN, Khairuddin KHAIRUDDIN, Gemini ALAM, Irnawati IRNAWATI, Abdul ROHMAN

DOI : 10.29228/jrp.2022.00

[Abstract \(abstract.php?lang=en&id=1783\)](#)

[Full Text PDF \(pdf.php?&id=1783\)](#)

[Similar Articles \(similar.php?&id=1783\)](#)

[E-mail to Author \(mailto:abdul\\_kimfar@ugm.ac.id\)](mailto:abdul_kimfar@ugm.ac.id)

**Properties of Chlorogenic Acid as a Skin Fibroblast Antiaging Agent with Apoptosis Inhibition and Free Radical Scavenging Activities (abstract.php?id=1784) Articles**

*Pages 159-167*

Ermi GIRSANG, I Nyoman Ehrich LISTER, Chrismis Novalinda GINTING, Wahyu WIDOWATI, Afif YATI, Hanna Sari Widya KUSUMA, Rizal AZIS

DOI : 10.29228/jrp.2022.00

[Abstract \(abstract.php?lang=en&id=1784\)](#)

[Full Text PDF \(pdf.php?&id=1784\)](#)

[Similar Articles \(similar.php?&id=1784\)](#)

[E-mail to Author \(mailto:ermigirsang@unprimdn.ac.id\)](mailto:ermigirsang@unprimdn.ac.id)

**Chitosan-based dry oral emulsion for isradipine delivery: Eco-friendly and surfactant free approach (abstract.php?id=1785) Articles**

*Pages 168-176*

Zahraa M. NAJI, Fatima J. JAWAD

DOI : 10.29228/jrp.2022.00

[Abstract \(abstract.php?lang=en&id=1785\)](#)

[Full Text PDF \(pdf.php?&id=1785\)](#)

[Similar Articles \(similar.php?&id=1785\)](#)

[E-mail to Author \(mailto:drfatimajalal@copharm.uobaghdad.edu.iq\)](mailto:drfatimajalal@copharm.uobaghdad.edu.iq)

**Micronucleus evaluation of remifentanil exposure (abstract.php?id=1786) Articles**

*Pages 177-181*

Öge ARTAGAN

DOI : 10.29228/jrp.2022.00

[Abstract \(abstract.php?lang=en&id=1786\)](#)

[Full Text PDF \(pdf.php?&id=1786\)](#)

[Similar Articles \(similar.php?&id=1786\)](#)

[E-mail to Author \(mailto:ogebasoglan.mu.edu.tr\)](mailto:ogebasoglan.mu.edu.tr)

**Alpha-1-acid glycoprotein and Pro-B-natriuretic peptide evaluation in patients with secondary pulmonary hypertension and end-stage kidney diseases receiving hemodialysis (abstract.php?id=1787) Articles**

*Pages 182-189*

Hanan Safaa AL-WAHEED,Raid J. Mohammed AL-TIMIMI,Arif Sami MALIK,Ali Ghafil AL-ZAYDI

DOI : 10.29228/jrp.2024.00

[Abstract \(abstract.php?lang=en&id=1787\)](#)

[Full Text PDF \(pdf.php?&id=1787\)](#)

[Similar Articles \(similar.php?&id=1787\)](#)

[E-mail to Author \(mailto:hanansafaa37@gmail.com\)](mailto:hanansafaa37@gmail.com)

**In silico approach of gambier extract for Diabetes Mellitus and multivariate optimization of ultrasound-assisted extraction rich polyphenol using full factorial design (abstract.php?id=1788) Articles**

*Pages 190-209*

Shaum SHIYAN,Galih PRATIWI,Sri HANDAYANI,Joko Tri WIBOWO,Indah SOLIHAAH,Hermansyah HERMANSYAH,Laras NOVITASARI,Athirah Azelia MARSYA,Nahla Akila FIKRIA,Susilawati SUSILAWATI,Puguh INDRASETIAWAN

DOI : 10.29228/jrp.2022.00

[Abstract \(abstract.php?lang=en&id=1788\)](#)

[Full Text PDF \(pdf.php?&id=1788\)](#)

[Similar Articles \(similar.php?&id=1788\)](#)

[E-mail to Author \(mailto:shaumshiy@unsri.ac.id\)](mailto:shaumshiy@unsri.ac.id)

**Design and development of Amphotericin-B loaded transungual drug delivery system for the effective management of onychomycosis (abstract.php?id=1789) Articles**

*Pages 210-221*

Yashwant GIRI,Amulyaratna BEHERA,Biswaranjan MOHANTY,Sk HABIBULLAH,Barsha PRIYADARSHINI,Gurudutta PATTNAIK,B JYOTIRMAYEE

DOI : 10.29228/jrp.2022.00

[Abstract \(abstract.php?lang=en&id=1789\)](#)

[Full Text PDF \(pdf.php?&id=1789\)](#)

[Similar Articles \(similar.php?&id=1789\)](#)

[E-mail to Author \(mailto:amulya.slmp@gmail.com\)](mailto:amulya.slmp@gmail.com)

**Study on ethnopharmacy of medicinal plants for antidiarrheal purposes among the Osing Tribe using Near Infrared Spectroscopy and chemometrics technique (abstract.php?id=1790) Articles**

*Pages 222-229*

Khrisna Agung CENDEKI AWAN,Dhina Ayu SUSANTI,Nina Wiji ANI,Florentinus Dika Octa RI SWANTO

DOI : 10.29228/jrp.2024.00

[Abstract \(abstract.php?lang=en&id=1790\)](#)

[Full Text PDF \(pdf.php?&id=1790\)](#)

[Similar Articles \(similar.php?&id=1790\)](#)

[E-mail to Author \(mailto:dikaocta@usd.ac.id\)](mailto:dikaocta@usd.ac.id)

**Dissolution methods to discriminate in vitro dissolution of poor water soluble weak base drug using three strategies: Acid modification, solvent evaporation and gastroretentive: Ciprofloxacin HCl case (abstract.php?id=1791) Articles**

*Pages 230-241*

Ahmed Ali Hashim AL AMERI,Masar Basim Mohsin MOHAMED

DOI : 10.29228/jrp.2022.00



[Abstract \(abstract.php?lang=en&id=1791\)](abstract.php?lang=en&id=1791)

[Full Text PDF \(pdf.php?&id=1791\)](pdf.php?&id=1791)

[Similar Articles \(similar.php?&id=1791\)](similar.php?&id=1791)

[E-mail to Author \(mailto:ahmedailaarr@uomustansiriyah.edu.iq\)](mailto:ahmedailaarr@uomustansiriyah.edu.iq)

---

Marmara University

[Home \(https://jrespharm.com/\)](https://jrespharm.com/)

[Aims and Scope \(static.php?id=1\)](static.php?id=1)

[Editorial Board \(static.php?id=2\)](static.php?id=2)

[Advisory Board \(content.php?id=102\)](content.php?id=102)

[Table of Contents \(static.php?id=4\)](static.php?id=4)

[Author Guidelines \(static.php?id=7\)](static.php?id=7)

[Abstracting / Indexing \(static.php?id=9\)](static.php?id=9)

[Contact \(static.php?id=8\)](static.php?id=8)

---

# Journal of Research in Pharmacy

An international open-access journal of pharmacy and pharmaceutical sciences  
Formerly published as Marmara Pharmaceutical Journal

(<https://jrespharm.com/>)

**JRP**  
ISSN : 1309-0801



[Current Issue \(https://jrespharm.com/content.php?id=102\)](https://jrespharm.com/content.php?id=102)

[Articles in Press \(https://jrespharm.com/content.php?id=46\)](https://jrespharm.com/content.php?id=46)

[Archive \(https://jrespharm.com/archive.php\)](https://jrespharm.com/archive.php)

[Search \(https://jrespharm.com/search.php\)](https://jrespharm.com/search.php)

[Author Benefits \(https://jrespharm.com/static.php?id=11\)](https://jrespharm.com/static.php?id=11)

## Editor-in-Chief

Hatice Kübra Elçioğlu

## Vice Editors

Levent Kabasakal

Esra Tatar

## Online ISSN

2630-6344

## Publisher

Marmara University

## Frequency

Bimonthly (Six issues / year)

## Abbreviation

J.Res.Pharm.

## Former Name

Marmara Pharmaceutical Journal

## Editorial Board

### Editor-in-Chief

#### Hatice Kübra ELÇİOĞLU

Department of Pharmacology, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
[kubra.elcioglu@marmara.edu.tr](mailto:kubra.elcioglu@marmara.edu.tr) (mailto:kubra.elcioglu@marmara.edu.tr)

### Vice Editors

#### Levent KABASAKAL

Department of Pharmacology, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
[lkabasakal@marmara.edu.tr](mailto:lkabasakal@marmara.edu.tr) (mailto:lkabasakal@marmara.edu.tr)

#### Esra TATAR

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
[etatar@marmara.edu.tr](mailto:etatar@marmara.edu.tr) (mailto:etatar@marmara.edu.tr)

## Analytical Chemistry & Therapeutic Drug Monitoring

### Anil Kumar DWIVEDI

Central Drug Research Institute, Lucknow, India  
[anilcdri@gmail.com](mailto:anilcdri@gmail.com) (mailto:anilcdri@gmail.com)

### Anisa ELHAMILI

Department of Medicinal & Pharmaceutical Chemistry, Faculty of Pharmacy, University of Tripoli, Tripoli, Libya  
[aaelhamili2000@gmail.com](mailto:aaelhamili2000@gmail.com) (mailto:aaelhamili2000@gmail.com)

### Emirhan NEMUTLU

Department of Analytical Chemistry, Faculty of Pharmacy, Hacettepe University, Ankara, Türkiye  
[enemutlu@hacettepe.edu.tr](mailto:enemutlu@hacettepe.edu.tr) (mailto:enemutlu@hacettepe.edu.tr)

### Lorena MEMUSHAJ

Department of Pharmacy, Faculty of Medical Sciences, Aldent University, Tirana, Albania  
[lorena.memushaj@ual.edu.al](mailto:lorena.memushaj@ual.edu.al) (mailto:lorena.memushaj@ual.edu.al)

### Mehmet GÜMÜŞTAŞ

Department of Forensic Toxicology, Institute of Forensic Sciences, Ankara University, Ankara, Türkiye  
[mgumustas@hotmail.com](mailto:mgumustas@hotmail.com) (mailto:mgumustas@hotmail.com)

### Mohd Younis RATHER

Multidisciplinary Research Unit, Government Medical College Srinagar, Srinagar, India  
[younis.rather78@gmail.com](mailto:younis.rather78@gmail.com) (mailto:younis.rather78@gmail.com)

**Pınar TALAY PINAR**

Department of Analytical Chemistry, Faculty of Pharmacy, Yüzüncü Yıl University, Van, Türkiye  
ptalay@gmail.com (mailto:ptalay@gmail.com)

*Biochemistry & Cancer Research***Beyza Ecem ÖZ BEDİR**

Department of Medical Biology, Faculty of Medicine, Ankara Yıldırım Beyazıt University, Ankara, Türkiye  
beoz@ybu.edu.tr (mailto:beoz@ybu.edu.tr)

**Débora DUMMER MEIRA**

Department of Biological Sciences, Nucleus of Human and Molecular Genetics, Federal University of Espírito Santo, Vitória- Espírito Santo, Brazil  
debora.dummer.meira@gmail.com (mailto:debora.dummer.meira@gmail.com)

**Derya ÖZSAVCI**

Department of Biochemistry, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
derya.ozsavci@marmara.edu.tr (mailto:derya.ozsavci@marmara.edu.tr)

**Emine TERZİ**

Department of Medical Biology, Faculty of Medicine, Ankara Yıldırım Beyazıt University, Ankara, Türkiye  
emineterzi1990@hotmail.com (mailto:emineterzi1990@hotmail.com)

**Gülberk UÇAR**

Department of Biochemistry, Faculty of Pharmacy, Hacettepe University, Ankara, Türkiye  
gulberk@hacettepe.edu.tr (mailto:gulberk@hacettepe.edu.tr)

**Haidar A ABDULAMIR**

College of Pharmacy, Al-Maaql University, Basra, Iraq  
h\_al\_attar@yahoo.com (mailto:h\_al\_attar@yahoo.com)

**Hamide Sena ÖZBAY**

Department of Biochemistry, Faculty of Pharmacy, Hacettepe University, Ankara, Türkiye  
senaozbay@hacettepe.edu.tr (mailto:senaozbay@hacettepe.edu.tr)

**İşıl YILDIRIM**

Pharmacy Services Program, Beykent University, Istanbul, Türkiye  
assistant.professor.isil.yildirim@gmail.com (mailto:assistant.professor.isil.yildirim@gmail.com)

**Lokman AYAZ**

Department of Biochemistry, Faculty of Pharmacy, Trakya University, Edirne, Türkiye  
lokmanayaz@yahoo.com (mailto:lokmanayaz@yahoo.com)

**Lynda BOUREBABA**

Department of Experimental Biology, Faculty of Biology and Animal Science, Wrocław University of Environmental and Life Sciences, Wrocław, Poland  
lynda.bourebaba@upwr.edu.pl (mailto:lynda.bourebaba@upwr.edu.pl)

**Nadia M. HAMDY**

Department of Biochemistry, Faculty of Pharmacy, Ain Shams University, Cairo, Egypt  
nadia\_hamdy@pharma.asu.edu.eg (mailto:nadia\_hamdy@pharma.asu.edu.eg)

**Sahar AL-OKBI**

Nutrition and Food Sciences Department, National Research Centre, Cairo, Egypt  
s\_y\_alokbi@hotmail.com (mailto:s\_y\_alokbi@hotmail.com)

**Selma HOUCHI**

Laboratory of Applied Biochemistry, Faculty of Natural and Life Sciences, University Ferhat Abbas, Setif, Algeria  
houchi.selma@univ-setif.dz (mailto:houchi.selma@univ-setif.dz)

*Biotechnology***Ali Demir SEZER**

Department of Pharmaceutical Biotechnology, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
adsezer@marmara.edu.tr (mailto:adsezer@marmara.edu.tr)

**Ammad Ahmad FAROOQI**

Department of Molecular Oncology, Institute of Biomedical and Genetic Engineering (IBGE), Islamabad, Pakistan  
farooqiammadahmad@gmail.com (mailto:farooqiammadahmad@gmail.com)

**Ceyda EKENTOK ATICI**

Department of Pharmaceutical Biotechnology, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
ceyda.ekentok@marmara.edu.tr (mailto:ceyda.ekentok@marmara.edu.tr)

**Fahima DILNAWAZ**

School of Engineering and Technology, Centurion University of Technology and Management, Odisha, INDIA  
fahimadilnawaz@gmail.com (mailto:fahimadilnawaz@gmail.com)

**Murat DOĞAN**

Department of Pharmaceutical Biotechnology, Faculty of Pharmacy, Cumhuriyet University, Sivas, Türkiye  
mdogan@cumhuriyet.edu.tr (mailto:mdogan@cumhuriyet.edu.tr)

**Uğur KARAGÖZ**

Department of Pharmaceutical Biotechnology, Faculty of Pharmacy, Trakya University, Edirne, Türkiye  
ugurkaragoz@trakya.edu.tr (mailto:ugurkaragoz@trakya.edu.tr)

*Clinical and Social Pharmacy & Pharmacoeconomy & Pharmacy Education***Abdikarim Mohammed ABDI**

Department of Clinical Pharmacy, Faculty of Pharmacy, Yeditepe University, Istanbul, Türkiye  
abdikarim.abdi@yeditepe.edu.tr (mailto:abdikarim.abdi@yeditepe.edu.tr)

**Ahmed Hamza AL-SHAMMARI**

Department of Pharmacy, Kut University College, Alkut, Wasit, Iraq  
Ahmedhamzamezaal@gmail.com (mailto:Ahmedhamzamezaal@gmail.com)

**Betül OKUYAN**

Department of Clinical Pharmacy, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
betul.okuyan@marmara.edu.tr (mailto:betul.okuyan@marmara.edu.tr)

**Emre KARA**

Department of Clinical Pharmacy, Faculty of Pharmacy, Hacettepe University, Ankara, Türkiye  
emrekara@hacettepe.edu.tr (mailto:emrekara@hacettepe.edu.tr)

**Ermelinda DURMISHI**

Director, Higher Education and Scientific Research Policies Department, Ministry of Education and Sports, Tirana, Albania  
eridurmishi@yahoo.com (mailto:eridurmishi@yahoo.com)

**Maja ORTNER HADŽIABDIĆ**

Centre for Applied Pharmacy, Faculty of Pharmacy and Biochemistry, University of Zagreb, Zagreb, Croatia  
mortner@pharma.hr (mailto:mortner@pharma.hr)

**Mesut SANCAR**

Department of Clinical Pharmacy, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
mesut.sancar@marmara.edu.tr (mailto:mesut.sancar@marmara.edu.tr)

**Mirela MIRAÇI**

Faculty of Pharmacy, University of Medicine, Tirana, Albania  
mirela.miraci@umed.edu.al (mailto:mirela.miraci@umed.edu.al)

**Nasir IDKAIDEK**

Department of Pharmaceutics and Pharmaceutical Technology, Faculty of Pharmacy and Medical Sciences, Petra University, Amman, Jordan  
nidkaidek@uop.edu.jo (mailto:nidkaidek@uop.edu.jo)

**Tarik CATIĆ**

Department of Pharmacy, Sarajevo School of Science and Technology, Sarajevo, Bosnia and Herzegovina  
tarik.catic@ssst.edu.ba (mailto:tarik.catic@ssst.edu.ba)

**Z.Kübra ÖZDEN YILMAZ**

Department of Clinical Pharmacy, Faculty of Pharmacy, Acıbadem Mehmet Ali Aydınlar University, Istanbul, Türkiye  
zekiye.yilmaz@acibadem.edu.tr (mailto:zekiye.yilmaz@acibadem.edu.tr)

*General Chemistry***Sinem GÖKTÜRK**

Department of Basic Pharmaceutical Sciences, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
sgokturk@marmara.edu.tr (mailto:sgokturk@marmara.edu.tr)

*In Silico Studies***Berna DOĞAN**

Department of Chemistry, Faculty of Science and Letters, Istanbul Technical University, Istanbul, Türkiye  
bernadogan@itu.edu.tr (mailto:bernadogan@itu.edu.tr)

**Gizem TATAR YILMAZ**

Department of Biostatistics and Medical Informatics, Faculty of Medicine, Karadeniz Technical University, Trabzon, Türkiye  
gizemtatar@gmail.com (mailto:gizemtatar@gmail.com)

**Onur SERÇİNOĞLU**

Department of Bioengineering, Faculty of Engineering, Gebze Technical University, Kocaeli, Türkiye  
osercinoglu@gtu.edu.tr (mailto:osercinoglu@gtu.edu.tr)

**Mehmet ÖZBİL**

Department of Bioengineering, Faculty of Engineering, Gebze Technical University, Kocaeli, Türkiye  
mozbil@gtu.edu.tr (mailto:mozbil@gtu.edu.tr)

*Medicinal Chemistry***Bahadır BÜLBÜL**

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Düzce University, Düzce, Türkiye  
bahadir.bulbul@yahoo.com.tr (mailto:bahadir.bulbul@yahoo.com.tr)

**Efe Doğukan DİNCEL**

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Istanbul University, Istanbul, Türkiye  
efe.dincel@istanbul.edu.tr (mailto:efe.dincel@istanbul.edu.tr)

**Entela HALOCI**

Faculty of Pharmacy, University of Medicine, Tirana, Albania  
entela.haloci@umed.edu.al (mailto:entela.haloci@umed.edu.al)

**Göknil Pelin COŞKUN**

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Acıbadem University, Istanbul, Türkiye  
pelin.coskun@acibadem.edu.tr (mailto:pelin.coskun@acibadem.edu.tr)

**Hasan Erdinç SELLİTEPE**

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Karadeniz Technical University, Trabzon, Türkiye  
esellitepe@ktu.edu.tr (mailto:esellitepe@ktu.edu.tr)

**Kerem BURAN**

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, University of Health Sciences, Istanbul, Türkiye  
Kerem.buran@sbu.edu.tr (mailto:Kerem.buran@sbu.edu.tr)

**Simone CARRADORI**

Department of Pharmacy, "G. d'Annunzio" University of Chieti-Pescara, Chieti, Italy  
simone.carradori@unich.it (mailto:simone.carradori@unich.it)

**Somaieh SOLTANI**

Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran  
soltanis@tbzmed.ac.ir (mailto:soltanis@tbzmed.ac.ir)

*Microbiology & Immunology***Shahram KHADEM VATAN**

Department of Medical Parasitology and Mycology, Urmia University of Medical Sciences, Urmia, Iran  
Cellular and Molecular Research Center, Cellular and Molecular Medicine Research Institute, Urmia University of Medical Sciences, Urmia, Iran  
Khademvatan@yahoo.com (mailto:Khademvatan@yahoo.com)

**Demet ERDÖNMEZ**

Department of Pharmaceutical Microbiology, Faculty of Pharmacy, Düzce University, Düzce, Türkiye  
demet.erdonmez@gmail.com (mailto:demet.erdonmez@gmail.com)

**Erkan RAYAMAN**

Department of Pharmaceutical Microbiology, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
erayaman@marmara.edu.tr (mailto:erayaman@marmara.edu.tr)

**Gülgün TINAZ**

Department of Basic Pharmaceutical Sciences, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
gulgun.tinaz@marmara.edu.tr (mailto:gulgun.tinaz@marmara.edu.tr)

**Zahraa AMER HASHIM**

Department of Microbiology and Immunology, College of Pharmacy, Mosul University, Mosul, Iraq  
hashimz@uomosul.edu.iq (mailto:hashimz@uomosul.edu.iq)

*Pharmaceutical Botany & Pharmacognosy & Chemistry of Natural Products***Ahmet EMİR**

Department of Pharmacognosy, Faculty of Pharmacy, Ege University, Izmir, Türkiye  
ahmet.emir@ege.edu.tr (mailto:ahmet.emir@ege.edu.tr)

**Annalisa CHIAVAROLI**

Department of Pharmacology, Faculty of Pharmacy, G. d'Annunzio University of Chieti-Pescara, Chieti, Italy  
annalisa.chiavaroli@unich.it (mailto:annalisa.chiavaroli@unich.it)

**Antoaneta TRENDAFILOVA**

Institute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Sciences, Sofia, Bulgaria  
antoaneta.trendafilova@orgchm.bas.bg (mailto:antoaneta.trendafilova@orgchm.bas.bg)

**Ceren EMİR**

Department of Pharmacognosy, Faculty of Pharmacy, Ege University, Izmir, Türkiye  
ceren.acir@ege.edu.tr (mailto:ceren.acir@ege.edu.tr)

**Claudio FERRANTE**

Department of Pharmacology, Faculty of Pharmacy, G. d'Annunzio University of Chieti-Pescara, Chieti, Italy  
claudio.ferrante@unich.it (mailto:claudio.ferrante@unich.it)

**İlker DEMİRBOLAT**

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Acibadem University, Istanbul, Türkiye  
ilker.demirbolat@acibadem.edu.tr (mailto:ilker.demirbolat@acibadem.edu.tr)

**İ. İrem TATLI ÇANKAYA**

Department of Pharmaceutical Botany, Faculty of Pharmacy, Hacettepe University, Ankara, Türkiye  
iremcankaya@gmail.com (mailto:iremcankaya@gmail.com)

**Laleh KHODAIE**

Department of Pharmacognosy, Faculty of Traditional Medicine, Tabriz University of Medical Sciences, Tabriz, Iran  
khodaiel@gmail.com (mailto:khodaiel@gmail.com)

**Lejla KLEPO**

Department of Chemistry, Faculty of Science, University of Sarajevo, Sarajevo, Bosnia and Herzegovina  
klepolejla@gmail.com (mailto:klepolejla@gmail.com)

**Mirjana MARČETIĆ**

Department of Pharmacognosy, Faculty of Pharmacy, University of Belgrade, Belgrade, Serbia  
mirjana.marctic@pharmacy.bg.ac.rs (mailto:mirjana.marctic@pharmacy.bg.ac.rs)

**Nurettin YAYLI**

Department of Pharmacognosy, Faculty of Pharmacy, Karadeniz Technical University, Trabzon, Türkiye  
yayli@ktu.edu.tr (mailto:yayli@ktu.edu.tr)

**Patrícia RIJO**

Research Center for Biosciences & Health Technologies, Lusofona University, Lisbon, Portugal  
p1609@ulusofona.pt (mailto:p1609@ulusofona.pt)

*Pharmacognosy***Sneha AGRAWAL**

Department of Pharmacognosy, Bharati Vidyapeeth's College of Pharmacy, Navi Mumbai, Maharashtra, India  
sneha.agrawal@bvcop.in (mailto:sneha.agrawal@bvcop.in)

**Turgut TAŞKIN**

Department of Pharmacognosy, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
turguttaskin@marmara.edu.tr (mailto:turguttaskin@marmara.edu.tr)

**Viktorija MAKSIMOVA**

Department of Applied Sciences, Faculty of Medical Sciences, Goce Delcev University, Shtip, Republic of N. Macedonia  
viktorija.maksimova@ugd.edu.mk (mailto:viktorija.maksimova@ugd.edu.mk)

**Vildan ÇELİKSOY**

School of Optometry and Vision Sciences, Cardiff University, Cardiff, UK  
celiksoyv92@gmail.com (mailto:celiksoyv92@gmail.com)

**Vilma TOSKA PAPAJANI**

Department of Pharmacy, University of Medicine, Tirana, Albania  
toskavilma@gmail.com (mailto:toskavilma@gmail.com)

**Zoran ZEKOVIĆ**

Faculty of Technology, University of Novi Sad, Novi Sad, Serbia  
zzekovic@tf.uns.ac.rs (mailto:zzekovic@tf.uns.ac.rs)

*Pharmaceutics***Monika DWIVEDI**

Department of Pharmaceutical Sciences and Technology, Birla Institute of Technology, Mesra, India  
monika.nbri@gmail.com (mailto:monika.nbri@gmail.com)

**Justė Baranauskaitė ORTASŪZ**

Department of Pharmaceutical Technology, Faculty of Pharmacy at Yeditepe University, Istanbul, Türkiye  
Department of Analytical and Toxicological Chemistry, Faculty of Pharmacy, Medical Academy, at the Lithuanian University of Health Sciences, Kaunas, Lithuania  
juste.ortasoz@yeditepe.edu.tr (juste.ortasoz@yeditepe.edu.tr)  
juste.baranauskaite@lsmuni.lt (juste.baranauskaite@lsmuni.lt)

**Afife Büşra UĞUR KAPLAN**

Department of Pharmaceutical Technology, Faculty of Pharmacy, Atatürk University, Erzurum, Türkiye  
afife.busra.ugur@gmail.com (mailto:afife.busra.ugur@gmail.com)

**Rajanikant PATEL**

Granules Pharmaceuticals Inc., Chantilly, VA - 20151, USA  
rajnipharmacy@gmail.com (mailto:rajnipharmacy@gmail.com)

**Burcu ÜNER**

Pharmaceutical and Administrative Sciences, The University of Health Science and Pharmacy in St. Louis, USA  
uner.burcu@yahoo.com (mailto:uner.burcu@yahoo.com)

**Dhanashree P. SANAP**

Department of Pharmaceutics, Bharati Vidyapeeth's College of Pharmacy, Navi Mumbai, India  
dhanashree.sanap@bvcop.in (mailto:dhanashree.sanap@bvcop.in)

**Dinesh KUMAR**

Department of Pharmaceutical Engineering & Technology, Indian Institute of Technology (BHU), Varanasi, India  
dinesh.phe@itbhu.ac.in (mailto:dinesh.phe@itbhu.ac.in)

**Ebru ALTUNTAŞ**

Department of Pharmaceutical Technology, Faculty of Pharmacy, Istanbul University, Istanbul, Türkiye  
ebru.altuntas@istanbul.edu.tr (mailto:ebru.altuntas@istanbul.edu.tr)

**Ela HOTI**

Faculty of Pharmacy, University of Medicine, Tirana, Albania  
ela.hoti@umed.edu.al (mailto:ela.hoti@umed.edu.al)

**Emrah ÖZAKAR**

Department of Pharmaceutical Technology, Faculty of Pharmacy, Atatürk University, Erzurum, Türkiye  
emrahozakar@atauni.edu.tr (mailto:emrahozakar@atauni.edu.tr)

**Enkelejda GOCI**

Pharmacotherapeutic Research Center, Aldent University, Tirana, Albania  
enkelejda.goci@ual.edu.al (mailto:enkelejda.goci@ual.edu.al)

**Kleva SHPATI**

Department of Pharmacy, Albanian University, Tirana, Albania  
k.shpati@albanianuniversity.edu.al (mailto:k.shpati@albanianuniversity.edu.al)

**Sakine TUNCAY TANRIVERDİ**

Department of Pharmaceutical Technology, Faculty of Pharmacy, Ege University, İzmir, Türkiye  
sakine.tuncay@ege.edu.tr (mailto:sakine.tuncay@ege.edu.tr)

**Gülşah GEDİK**

Department of Pharmaceutical Technology, Faculty of Pharmacy, Trakya University, Edirne, Türkiye  
gulsahgedik@trakya.edu.tr (mailto:gulsahgedik@trakya.edu.tr)

**Ongun Mehmet SAKA**

Department of Pharmaceutical Technology and Biotechnology, Faculty of Pharmacy, Ankara University, Ankara, Türkiye  
omsaka@gmail.com (mailto:omsaka@gmail.com)

**Oya KERİMOĞLU**

Department of Pharmaceutical Technology, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
osipahigil@marmara.edu.tr (mailto:osipahigil@marmara.edu.tr)

**Pankaj DWIVEDI**

Pharmaceutical and Administrative Sciences, The University of Health Science and Pharmacy in St. Louis, USA  
dwivedipank@gmail.com (mailto:dwivedipank@gmail.com)

**Rezarta SHKRELI**

Department of Pharmacy, Faculty of Medical Sciences, Aldent University, Tirana, Albania  
rezarta.shkreli@ual.edu.al (mailto:rezarta.shkreli@ual.edu.al)

**Renuka KHATIK**

Washington University in St. Louis, USA  
renukadops@gmail.com (mailto:renukadops@gmail.com)

**Rukiye SEVİNÇ ÖZAKAR**

Department of Pharmaceutical Technology, Faculty of Pharmacy, Atatürk University, Erzurum, Türkiye  
rukiyeso@atauni.edu.tr (mailto:rukiyeso@atauni.edu.tr)

**Saeideh SOLTANI**

Novel Drug Research Center, School of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Sciences, Isfahan, Iran  
soltanisa@pharm.mui.ac.ir (mailto:soltanisa@pharm.mui.ac.ir)

*Pharmacology & Toxicology***Ana V. PEJČIĆ**

Department of Pharmacology and Toxicology, Faculty of Medical Sciences, University of Kragujevac, Kragujevac, Serbia  
anapejic201502@yahoo.com (mailto:anapejic201502@yahoo.com)

**Ayfer BECEREN**

Department of Pharmaceutical Toxicology, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
ayfer.tozan@marmara.edu.tr (mailto:ayfer.tozan@marmara.edu.tr)

**Ayşenur GÜNAYDIN AKYILDIZ**

Department of Pharmaceutical Toxicology, Faculty of Pharmacy, Bezmialem Vakıf University, Istanbul, Türkiye  
gunaydinaysenur@gmail.com (mailto:gunaydinaysenur@gmail.com)

**Ayça TOPRAK SEMİZ**

Vocational School of Health Services, Giresun University, Giresun, Türkiye  
ayca.toprak@giresun.edu.tr (mailto:ayca.toprak@giresun.edu.tr)

**Büşra ERTAŞ**

Department of Pharmacology, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
busra.ertas@marmara.edu.tr (mailto:busra.ertas@marmara.edu.tr)

**Vasudevan MANI**

Department of Pharmacology and Toxicology, College of Pharmacy, Qassim University, Al Qassim, Kingdom of Saudi Arabia  
v.samy@qu.edu.sa (mailto:v.samy@qu.edu.sa)

**Fatiha MISSOUN**

Laboratory of Pharmacognosy and Api-Phytotherapy, University of Mostaganem, Mostaganem, Algeria  
fatiha.missoun@univ-mosta.dz (mailto:fatiha.missoun@univ-mosta.dz)

**Klodiola DHAMO**

Faculty of Technical Medical Sciences, Aldent University, Tirana, Albania  
klodiola.dhamo@ual.edu.al (mailto:klodiola.dhamo@ual.edu.al)

**Long Chiau MING**

School of Medical and Life Sciences, Sunway University, Sunway City, Malaysia  
longchiauming@gmail.com (mailto:longchiauming@gmail.com)

**Merve KABASAKAL**

Department of Medical Pharmacology, Faculty of Medicine, University of Health Sciences, Istanbul, Türkiye  
merve.kabasakal@sbu.edu.tr (mailto:merve.kabasakal@sbu.edu.tr)

**Miloš N. MILOSAVLJEVIĆ**

Department of Pharmacology and Toxicology, Faculty of Medical Sciences, University of Kragujevac, Kragujevac, Serbia  
milosavljevicmilos91@gmail.com (mailto:milosavljevicmilos91@gmail.com)

**Mohammed Jabbar MANNA**

Department of Pharmacology, College of Dentistry, Al-Mustansiriya University, Baghdad, Iraq  
mohammedalmanna@uomustansiriyah.edu.iq (mailto:mohammedalmanna@uomustansiriyah.edu.iq)

**Nurdan TEKİN**

Department of Medical Pharmacology, Faculty of Medicine, University of Health Sciences, Istanbul, Türkiye  
nurdan.tekin@sbu.edu.tr (mailto:nurdan.tekin@sbu.edu.tr)

**Oğuzhan AYDEMİR**

İstanbul Kent Üniversitesi, Eczacılık Fakültesi, Farmakoloji Anabilim Dalı, İstanbul, Türkiye  
aydemir.oguzhan@hotmail.com (mailto:aydemir.oguzhan@hotmail.com)

**Rümeysa KELEŞ KAYA**

Department of Medical Pharmacology, Faculty of Medicine, Sakarya University, Sakarya, Türkiye  
rumeysakeles@sakarya.edu.tr (mailto:rumeysakeles@sakarya.edu.tr)

**Sana REHMAN**

Department of Pharmacology, HIMSR & HAHC Hospital, Jamia Hamdard, New Delhi, INDIA  
drsanarehman2012@gmail.com (mailto:drsanarehman2012@gmail.com)

**Sinan SERMET**

Istinye University Faculty of Medicine, Department of Clinical Sciences and Department of Pharmacology and Clinical Pharmacology, Istanbul, Türkiye  
sinan.sermet@istinye.edu.tr (mailto:sinan.sermet@istinye.edu.tr)

**Ünzile YAMAN**

Department of Pharmaceutical Toxicology, Faculty of Pharmacy, Katip Çelebi University, İzmir, Türkiye  
unzileyaman@gmail.com (mailto:unzileyaman@gmail.com)

**Zarife Nigar ÖZDEMİR KUMRAL**

Department of Physiology, Faculty of Medicine, Marmara University, Istanbul, Türkiye  
znozdemir@marmara.edu.tr (mailto:znozdemir@marmara.edu.tr)

**Zeina ALTHANOON**

Department of Pharmacology and Toxicology, College of Pharmacy, Mosul University, Mosul, Iraq  
dr.zeina@uomosul.edu.iq (mailto:dr.zeina@uomosul.edu.iq)

*Copy Editor***Ayşe Nur HAZAR YAVUZ**

Department of Pharmacology, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
ayse.hazar@marmara.edu.tr (mailto:ayse.hazar@marmara.edu.tr)

**Büşra ERGEN**

Department of Clinical Pharmacy, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
busra.ergen@marmara.edu.tr (mailto:busra.ergen@marmara.edu.tr)

**Elif Beyzanur POLAT**

Department of Pharmacology, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
elif.beyzanur@marmara.edu.tr (mailto:elif.beyzanur@marmara.edu.tr)

**Fatih Taha ÇİFTÇİ**

Department of Clinical Pharmacy, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
fatih.ciftci@marmara.edu.tr (mailto:fatih.ciftci@marmara.edu.tr)

**Müzeyyen AKSOY**

Department of Clinical Pharmacy, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
muzeyyen.aksoy@marmara.edu.tr (mailto:muzeyyen.aksoy@marmara.edu.tr)

**Ömer Faruk ÖZKANLI**

Department of Clinical Pharmacy, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
oozkanli@marmara.edu.tr (mailto:oozkanli@marmara.edu.tr)

**Şeyma GÖZELİZMİR**

Department of Clinical Pharmacy, Faculty of Pharmacy, Marmara University, Istanbul, Türkiye  
seyma.gozelizmir@marmara.edu.tr (mailto:seyma.gozelizmir@marmara.edu.tr)

*Language Editor***Khadija ALJESRI**

Department of Pharmacology, Institute of Health Sciences, Marmara University, Istanbul, Türkiye

*Biostatistics Editor***Gülnaz NURAL BEKİROĞLU**

Department of Biostatistics, Faculty of Medicine, Marmara University, Istanbul, Türkiye  
nural@marmara.edu.tr (mailto:nural@marmara.edu.tr)

Marmara University

Home (<https://jrespharm.com/>)

Aims and Scope (<static.php?id=1>)

Editorial Board (<static.php?id=2>)

Advisory Board (<content.php?id=102>)

Table of Contents (<static.php?id=4>)

Author Guidelines (<static.php?id=7>)

Abstracting / Indexing (<static.php?id=9>)

Contact (<static.php?id=8>)

---



# Study on ethnopharmacy of medicinal plants for antidiarrheal purposes among the Osing Tribe using Near Infrared Spectroscopy and chemometrics techniques

Khrisna Agung CENDEKIAWAN<sup>1,2</sup>, Dhina Ayu SUSANTI<sup>2</sup>, Nina WIJIANI<sup>3,4</sup>, Florentinus Dika Octa RISWANTO<sup>1\*</sup>

<sup>1</sup> Division of Pharmaceutical Analysis and Medicinal Chemistry, Faculty of Pharmacy, Sanata Dharma University, Campus 3 Paingan Maguwoharjo Depok Sleman, Yogyakarta 55282, Indonesia

<sup>2</sup> Pharmacy Study Program, Universitas dr. Soebandi, Jalan dr. Soebandi Nomor 99 Jember, Indonesia

<sup>3</sup> Pharmacy Study Program, STIKES Banyuwangi, Jalan Letkol Istiqolah 109 Banyuwangi, Indonesia

<sup>4</sup> Faculty of Pharmacy, Jember University, Jalan Kalimantan No 37 Jember, Indonesia

\*Corresponding Author. E-mail: dikaocta@usd.ac.id (F.D.O.R); Tel. +6285729071377.

Received: 16 January 2024 / Revised: 28 March 2024 / Accepted: 31 March 2024

**ABSTRACT:** Traditional healing culture has long been known in Indonesia, this is proven by the existence of inscriptions that tell about herbal medicine that has been used for generations. As time goes by, the use of plants as medicine has decreased. This is because the inheritance process is only verbal and there is no documentation about these medicinal plants. This encourages research and preservation of plants as medicine, one of which is ethnopharmacy. The implementation of ethnopharmacy in the Osing tribe regarding the use of medicinal plants as anti-diarrhea can provide knowledge about the existence of plants that contain flavonoid compounds which have the potential to act as anti-diarrhea. This study aimed at verifying the anti-diarrheal attributes of medicinal plant compounds, *Near Infra-Red* (NIR) spectroscopy and chemometric techniques were employed. Leveraging the established principle of flavonoid compounds as potent antioxidants capable of stabilizing the digestive tract, we employed flavonoid standards as a reference benchmark for assessing the anti-diarrheal efficacy of these compounds. This approach facilitated a comprehensive evaluation to verify the extent of their anti-diarrheal potential. The results of this research found that the guava leaf plant is widely used by the Osing tribe as an anti-diarrhea. Next, guava leaves were analyzed using *Near Infra Red* (NIR) spectroscopy and chemometric techniques, and test results were obtained in the form of guava leaf samples from the Osing tribe that contain flavonoids so they can be used for antidiarrheal therapy.

**KEYWORDS:** Ethnopharmacy; antidiarrhea; spectroscopy; chemometrics.

## 1. INTRODUCTION

Within the Indonesian society, medicinal plants occupy a distinct position, representing a cultural heritage in the domain of health and standing out as a well-established cultural legacy renowned for its therapeutic efficacy. Consequently, these medicinal plants have been passed down through generations and frequently employed by the Indonesian population as integral elements of traditional medicine essential for the community [1]. The Osing region, situated around the Ijen Mountains and Banyuwangi Regency in East Java Province, exhibits a distinct cultural identity that encourages researchers seeking to explore this area [2]. While numerous publications delve into the religious rituals practiced by the Osing community, limited documentation exists regarding the essential knowledge of medicinal plant usage. This documentation serves as an important resource, providing pivotal insights and serving as a foundational exploration for the discovery of potential new medicines [3,4].

To explore the indigenous community's knowledge regarding the use of medicinal plants, an approach rooted in ethnopharmacy can be employed. Ethnopharmacy constitutes an interdisciplinary field focused on the utilization of medicinal plants within the context of a specific society's healing culture [5]. While extensive research exists on ethnopharmacy, particularly concerning various ethnic groups in Indonesia, a limited number of research has been directed towards the Osing community. Therefore, this study serves as an initial gateway to unveil the traditional medicinal practices of the Osing ethnic group in East Java, offering a vital starting point for further investigation [6].

**How to cite this article:** Cendekiawan KA, Susanti DA, Wijiani, N, Riswanto, FDO. Study on ethnopharmacy of medicinal plants for antidiarrheal purposes among the Osing Tribe using Near Infrared Spectroscopy and chemometrics techniques. J Res Pharm. 2025; 29(1): 222-229.






The detection of compound presence often involves employing analytical methodologies namely *Near Infra Red* (NIR) spectroscopy and chemometrics [7]. NIR spectroscopy operates on electromagnetic wavelengths beyond the visible light spectrum (specifically within the 700-3000 nm range), enabling for the identification of compounds in a given preparation or sample [8]. Its widespread application extends notably to the analysis of organic compounds. Additionally, chemometric techniques serve as a supporting approach for quantitative analysis, facilitating further exploration of the variations among constituent elements within a sample, thereby enabling precise measurement of compound quantities [9,10]. These techniques were employed to ensure validation and safety for consumers of medicinal plants to optimize the expected therapeutic outcomes [11].

Based on the aforementioned explanation, the researcher was encouraged to conduct an investigation focusing on the ethnopharmacology of anti-diarrheal medicinal plants within the Osing ethnic community. This study aims to employ *Near Infra Red* spectroscopy and chemometric techniques as instrumental tools in the research process.

## 2. RESULTS

Through ethnopharmacological investigations, it was observed that guava leaves are prominently used within the Osing ethnic group to manage diarrheal conditions. Subsequent phytochemical screening identified the presence of quercetin flavonoids in the extract [12]. The detailed outcomes are presented in Table 1.

**Table 1.** Phytochemical Screening of Guava Leaf Extract was conducted through several tests based on the suspected compound content within the extract using specific reagents.

Phytochemical Test	Reagent	Observation	Result
Alkaloid	• Extract + Mayer		+
	• Extract + Bouchardat		+
	• Extract + Mayer		+
Flavonoid	• Extract + Mg powder + amyl alcohol		+
	• Extract + concentrated HCl		+
Tannin or Phenolic	• Extract + FeCl <sub>3</sub>		+
Saponin	• Extract + hot water (shaken vertically for 10 minutes)		-
Triterpenoid or Steroid	• Extract + Liebermann Bouchard		+


Description:

+ : Contains secondary metabolite compounds

- : Does not contain secondary metabolite compounds


The next stage involves preparing simulated samples of guava leaf herbal material, followed by analysis using *Near Infra Red* (NIR) and chemometric techniques to establish a chemometric classification model. This model was then applied to guava leaf herbal material samples used by the Osing community in the village of Banjar, Licin District, Banyuwangi Regency.

A classification model was established using a training dataset. The spectra from each dataset were utilized to construct a chemometric classification model, specifically Linear Discriminant Analysis (LDA) [13]. From the results of the infrared scan in Figure 1, the entire infrared spectra generated from both pure guava leaf herbal material and the mixture in the training set and test set showed similar absorption patterns, differing only in the quantitative values of their respective spectrum absorbances. The outcomes demonstrated a mapping illustrating the differentiation between pure guava leaf herbal material and mixtures. Meanwhile, the prediction table exhibited the classification of samples into the established categories.




**Figure 1.** Visualization of the infrared spectra of the training set (a) and test set (b) of guava leaf simplicia

Subsequently, the establishment of a classification model was undertaken utilizing the training set data. The spectra derived from each dataset within the training set were employed to formulate a chemometric classification model, specifically employing Linear Discriminant Analysis (LDA). Consequently, a systematic evaluation was conducted to discern the most optimal model that establishes a meaningful association between categories and discriminants. The results, in the form of mapping, demonstrate a distinct segregation between the categories of pure and mixed guava leaf herbal material on dried leaf (Figure 2) and dried powder leaf of guava (Figure 3).



**Figure 2.** The model properties of the dried guava leaf herbal material training set in (a) partial least square (PLS) with R-Square values and (b) Linear Discriminant Analysis (LDA) with accuracy values.



**Figure 3.** The model properties of the dried guava leaf powder herbal material training set in (a) partial least square (PLS) with R-Square values and (b) Linear Discriminant Analysis (LDA) with accuracy values.

Based on the data of the Linear Discriminant Analysis, the %recognition and %prediction values of each model formed based on the shape of the consumed guava leaves are obtained, namely the dried simplicia form (Data Set 1) and the powder form (Data Set 2) as shown in Table 2.

**Table 2.** Assessment of the dataset based on the results of chemometric analysis in Linear Discriminant Analysis.

Data Set	LDA	
	% Recognition	% Prediction
1	80%	90%
2	100%	100%

Subsequently, an application was conducted on guava leaf samples obtained from the ethnopharmacological study of medicinal plants for anti-diarrheal purposes in the Village of Banjar, Licin District, Banyuwangi Regency. Based on the results obtained using chemometric techniques applied to guava leaf simplicia collected from the area of Banjar Village, Licin District, Banyuwangi Regency, the samples indicated a mixed herbal material showing the presence of quercetin flavonoids.

### 3. DISCUSSION

This ethnopharmacy research was conducted among the Osing tribe in the village of Banjar, Licin Subdistrict, Banyuwangi Regency, utilizing a questionnaire distributed to respondents who met the inclusion criteria. The selection of this location was based on the substantial Osing population residing in Banjar village, Licin Subdistrict, where the majority still adheres to cultural practices passed down through generations [14]. This choice aimed to deepen information concerning the traditional treatment of diarrhea using medicinal plants. The ethnopharmacy study was chosen because it includes several methods for determining which medical plants are widely used and their properties in society. Ethnopharmacy does not only focus on medicinal plants themselves, but also on the cultural and social aspects associated with their use. This allows researchers to understand in more depth how medicinal plants are integrated into people’s live. The ethnopharmacy results revealed that guava leaf is the most frequently consumed plant by the Osing tribe to cure diarrhea [15,16]. Subsequently, the methods used in ethnopharmacy studies include laboratory analysis or phytochemical screening. Phytochemical screening was performed to investigate the compound composition present in guava leaves using various methods tailored to identify specific compounds. The phytochemical screening results indicated that guava leaves contain several bioactive compounds with potential therapeutic benefits for the community [17].

The next stage involves the preparation of simulated guava leaf samples followed by analysis using Near Infrared (NIR) spectroscopy and chemometrics. This preparation entails creating a dataset using medicinal plants identified through ethnopharmacy, specifically guava leaves [18]. The simulated guava leaf samples include pure guava leaf, quercetin flavonoid standard, and a mixture of guava leaf herbal material with the flavonoid standard. Flavonoids have been shown to have potential benefits in treating diarrhea with potential mechanisms as antibacterial activity so potentially helping fight off diarrhea-causing bacteria. This aligns with the practices of the Osing tribe in Banjar Village, who commonly consume guava leaves in the form of dried or powdered leaves to heal diarrhea [16]. Both forms have advantages and disadvantages.

Dried simplicias (whole or chopped plant parts) are easier to handle and store without clumping, but it requires grinding or crushing before use, which can lead to loss of volatile oils and other active components. Whereas powder simplicia (ground plant parts) are easier to measure doses and more convenient for more applications (e.g., capsules or hot drink) but they have higher risk of clumping due to increased surface area contact.

Afterwards, the IR spectrum was utilized to form a chemometric classification model and its application to the guava leaf samples used by the Osing Tribe community in Banjar Village, Licin Subdistrict, Banyuwangi Regency. This modeling was conducted to ensure which model was appropriate for application to samples found in the Osing tribe community, thereby determining the proper dataset model for the classification of anti-diarrheal compound content [19]. The powder samples, totaling twenty for the training set and five for the test set, were also measured using a Near Infrared (NIR) spectrophotometer. The similarity in NIR spectra patterns may arise due to the presence of similar compound contents in the tested herbal material dataset. Meanwhile, variations in absorbance can be attributed to differing quantities of compounds in each tested herbal material [20].

Subsequently, the establishment of a classification model was undertaken utilizing the training set data. The spectra derived from each dataset within the training set were employed to formulate a chemometric classification model, specifically employing Linear Discriminant Analysis (LDA). LDA hinges on the dataset's capacity to effectively categorize or discriminate one class from another, with a higher degree of precision indicative of superior dataset performance [21,22]. Two distinct datasets were delineated based on the structural forms of guava leaf herbal material, encompassing both the dried simplicia and powdered forms [23]. Consequently, a systematic evaluation was conducted to discern the most optimal model that establishes a meaningful association between categories and discriminants. The results, in the form of mapping, demonstrate a distinct segregation between the categories of pure and mixed guava leaf herbal material. Meanwhile, the prediction table presents the classification of samples into the designated categories, as depicted in Figures 2 and 3.

Subsequently, testing was conducted to assess the model's ability to distinguish between the two types of guava leaf herbal material, as evident in the recognition capability values for samples in the training set and the predictive ability of LDA for samples in the test set. It was found that the LDA model of dried guava leaf herbal material resulted RMSEC and RMSEP values of 0.0257 ( $R^2_{\text{cal}} = 0.9821$ ) and 0.0277 ( $R^2_{\text{val}} = 0.9798$ ), respectively. Whereas, the LDA model of dried guava leaf powder herbal material resulted RMSEC and RMSEP values of 0.0109 ( $R^2_{\text{cal}} = 0.9931$ ) and 0.0124 ( $R^2_{\text{val}} = 0.9919$ ), respectively. Table 2 demonstrates that the optimal chemometric classification model for pure guava leaf herbal material and mixed guava leaf simplicia is the LDA model based on the IR spectrum in training set 2, in powder form, where the recognition and prediction values are both 100%. Hence, the guava leaf samples obtained from the ethnopharmacy study need to be processed into dry powder form [24]. This is inline with the theory that dry powder simplicia have a form that easier for interaction of flavonoid compounds with *near infra red*, so that the results of the analysis process can be identified more quickly and accurately.

Next, an application was carried out on guava leaf samples obtained from the ethnopharmacy study of medicinal plants for antidiarrheal purposes in Banjar Village, Licin Subdistrict, Banyuwangi Regency. Based on the results in Table 2, utilizing the chemometric techniques conducted on guava leaf simplicia collected in the area of Banjar Village, Licin Subdistrict, Banyuwangi Regency, it indicates that the samples are categorized as mixed simplicia, demonstrating the presence of quercetin flavonoid compound content.

#### 4. CONCLUSION

From the results of the ethnopharmacy study on medicinal plants for anti-diarrheal purposes, the most widely consumed by the Osing Tribe in Licin Subdistrict, Banyuwangi Regency, is guava leaf. However, there are other plants also used as anti-diarrheals. Subsequently, the identification of anti-diarrheal compounds in guava leaves in Banjar Village, Licin Subdistrict, Banyuwangi Regency, using Near Infrared and Chemometrics indicates the presence of quercetin flavonoid content. A recommendation for further research is to conduct anti-diarrheal activity tests on medicinal plants used for this purpose by the Osing Tribe in Banjar Village or other villages within the Licin Subdistrict of Banyuwangi Regency.

#### 5. MATERIALS AND METHODS

##### 5.1. Material

The materials in this study utilized samples of medicinal plants (comprising dried herbal material

and reference compounds); quercetin flavonoid standard with ChromaDex analysis certificate from PT. Dexa Medica; and samples of medicinal plants from the Osing Tribe.

## 5.2. Tools

A questionnaire attached with ethical approval was used the survey stage. The instruments used in this study were analytical balance, oven, blender, sieve with size B-30. A set of near-infrared spectrometer (Brimrose Corporation Luminar 3070) was exploited for data acquisition process coupled with BRIMROSE software and The Unscrambler X 10.2 software for data analysis.

## 5.3. Ethnopharmacy study

This research commenced with a preliminary survey, followed by data collection through an ethnopharmacy approach conducted in the Licin Subdistrict, Banyuwangi Regency. The selection of the village was based on the predominant presence of the Osing Tribe in the community. Subsequently, chemical content testing on the collected samples will be carried out in the Chemistry Laboratory of the Faculty of Pharmacy at the University of Jember.

The ethnopharmacy study took place in Banjar Village, Licin Subdistrict, Banyuwangi Regency (Latitude: -8.193077160373061; Longitude: 114.26683913916348; R748+QPM). Ethical approval for the ethnopharmacy study was obtained, submitted to Universitas dr. Soebandi Jember, with the reference number: 441/KEPK/UDS/IX/2023. Following this, data collection was conducted using interview and questionnaire methods involving 96 samples from the Osing Tribe in Banjar Village. The ethnopharmacy results identified guava leaf as a frequently used plant to address diarrhea-related complaints among the Osing Tribe in Banjar Village, Licin Subdistrict, Banyuwangi Regency. The guava leaf samples were then determined at the Herbal Materia Medica Laboratory in Batu, with herbarium number: 000.9.3/2911/102.20/2023.

## 5.4. Sampel preparation

The sample preparation was carried out by combining simplicia and compounds. The prepared simulated samples consisted of plant simplicia and compound mixtures. The simulated samples were divided into two groups, namely the training set and test set. Guava leaf simplicia samples mixed with quercetin flavonoid were prepared at various concentrations within the quercetin flavonoid concentration range of 1-100% of the total ( $\pm 10$  milligrams), as presented in Table 3.

## 5.5. Chemometrics techniques

The sample to be measured was placed on the sample holder located in the Integrating Sphere unit. Light from the halogen lamp would pass through several sets of equipment and be filtered as specified. After the light hits the sample, the reflection of near-infrared light would be captured by the sensor and undergo digitization. Measurements on the NIR Spectrophotometer employed a filter with a wavelength range of 1400-2000 nm and a data collection interval of 5 nm, resulting in obtaining reflection data for a total of 120 points.

In chemometric analysis using The Unscrambler X 10.2 software, the selection of the classification model was based on the best recognition ability and prediction ability. Recognition ability is defined as the percentage of correct classification by the model against the training set samples, while prediction ability is defined as the percentage of correct classification by the model against the test set samples. Quality of the chemometrics models were assessed according to the values of the calibration determination coefficient ( $R^2_{cal}$ ), validation determination coefficient ( $R^2_{val}$ ), root mean square error of calibration (RMSEC) and root mean square error of prediction (RMSEP).

**Table 3.** The Training Set data was created with a composition of guava leaf simplicia and quercetin flavonoid.

Quercetin Flavonoid (gram)	Guava Leave Herbal Material (gram)	Concentration (%)	Category
0	10	0%	Pure
0.011	10	1.1%	Mixture
0.021	10	2.1%	Mixture
0.032	10	3.2%	Mixture
0.040	10	4%	Mixture
0.051	10	5.1%	Mixture
0.060	10	6%	Mixture
0.073	10	7.3%	Mixture
0.082	10	8.2%	Mixture
0.092	10	9.2%	Mixture
0.1015	10	10.05%	Mixture
0.2040	10	20.14%	Mixture
0.3028	10	30.18%	Mixture
0.4004	10	40.03%	Mixture
0.5014	10	50.10%	Mixture
0.6005	10	60.00%	Mixture
0.7038	10	70.00%	Mixture
0.8022	10	80.01%	Mixture
0.9033	10	90.93%	Mixture
1.0011	0	100.00%	Mixture

The test set consists of five sets of simulated guava leaf herbal material data, namely pure guava leaf raw material, quercetin flavonoid material, and guava leaf raw material mixture with concentrations of 20%, 50%, and 70%, as shown in Table 4. Pure guava leaf herbal material is categorized within the mixed guava leaf herbal material to evaluate the formed model.

**Table 4.** Test set data with a composition of guava leaf herbal material and quercetin flavonoid.

Quercetin Flavonoid (gram)	Guava Leave Herbal material (gram)	Concentration (%)	Category
0	10	0%	Pure
0.2040	10	20.00%	Mixture
0.5014	10	50.05%	Mixture
0.703	10	70%	Mixture
1.0011	0	100%	Mixture

This is an open access article which is publicly available on our journal's website under Institutional Repository at <http://dspace.marmara.edu.tr>.

**Acknowledgements:** The researcher expresses gratitude to Dexa Medica for their collaboration in obtaining the quercetin standard, to the Village Government of Banjar in the Licin Subdistrict of Banyuwangi Regency for facilitating the collection of ethnopharmacy data, to the Faculty of Pharmacy at the University of Jember for their partnership in processing chemometric data, and to Universitas dr. Soebandi Jember for their collaboration in obtaining ethical clearance.

**Author contributions:** Concept – K.A.C, F.D.O.R.; Design – K.A.C.; Supervision – F.D.O.R.; Resources – K.A.C., D.A.S.; Materials – D.A.S.; Data Collection and/or Processing – K.A.C., N.W., D.A.S; Analysis and/or Interpretation – K.A.C., F.D.O.R., D.A.S., N.W.; Literature Search – K.A.C., D.A.S., N.W.; Writing – K.A.C.; Critical Reviews – K.A.C., F.D.O.R., N.W., D.A.S.

**Conflict of interest statement:** The authors declared no conflict of interest.

## REFERENCES

- [1] Reyes-García V. The Relevance of Traditional Knowledge Systems for Ethnopharmacological Research: Theoretical

- and Methodological Contributions. *J Ethnobiol Ethnomed.* 2010; 6: 1-12. <https://doi.org/10.1186/1746-4269-6-32/METRICS>.
- [2] Prasetyo B, Chikmawati T, Walujo E.B, Amzu E. Ethnoecology: The traditional landscape of Osing Tribe in Banyuwangi, Indonesia. *Biodiversitas.* 2018; 19(6): 2003–2009. <https://doi.org/10.13057/BIODIV/D190604>.
- [3] Kusumo DW, Sulistiyowati EL, Rohmah H, Melinda NA. Ethno-pharmaceutical study of medicinal plants for care and treatment of wounds in Indonesia: Systematic data search and preclinical review. *J Jamu Indones.* 2023; 8: 1-9. <https://doi.org/10.29244/JJI.V8I1.232>.
- [4] Sumarni W, Sudarmin S, Sumarti SS. The scientification of Jamu: A study of Indonesian's traditional medicine. *J Phys Conf Ser.* 2019; 1321(3): 1-7. <https://doi.org/10.1088/1742-6596/1321/3/032057>.
- [5] Bhagawan WS, Suproborini A, Prastya Putri DL, Nurfatma A, Putra RT. Ethnomedicinal study, phytochemical characterization, and pharmacological confirmation of selected medicinal plant on the northern slope of mount Wilis, East Java, Indonesia. *Biodiversitas.* 2022; 23(8): 4303–4313. <https://doi.org/10.13057/BIODIV/D230855>.
- [6] Puspitasari E, Damayanti AE, Zikrina INS, Dianasari D. Exploring the ethnopharmaceutical plants of Osing Tribe in Banyuwangi Regency: Potential application for COVID-19 Therapy. *Ulum Islam.* 2021; 33: 113-120. <https://doi.org/10.33102/uij.vol33no3.330>.
- [7] Fakayode SO, Baker GA, Bwambok DK, Bhawawet N, Elzey B, Siraj N, Macchi S, Pollard DA, Perez RL, Duncan AV. Molecular (Raman, NIR, and FTIR) Spectroscopy and multivariate analysis in consumable products analysis. *Appl Spectros Rev.* 2020; 55(8): 647–723. <https://doi.org/10.1080/05704928.2019.1631176>.
- [8] Hair JF, Hult GTM, Ringle CM, Sarstedt M, Danks NP, Ray S. Partial Least Squares Structural Equation Modeling (PLS-SEM) Using R. Springer. 2021: 197 <https://doi.org/10.1007/978-3-030-80519-7>.
- [9] Irnawati, Riyanto S, Martono S, Rohman A. Analysis of palm oil as oil adulterant in olive and pumpkin seed oils in ternary mixture systems using FTIR Spectroscopy and Chemometrics. *Int J Appl Pharm.* 2019; 11(5): 210-215. <https://doi.org/10.22159/ijap.2019v11i5.34274>.
- [10] Windarsih A, Rohman A, Irnawati, Riyanto S. The combination of vibrational spectroscopy and chemometrics for analysis of milk products adulteration. *Int J Food Sci.* 2021; 2021: 8853358. <https://doi.org/10.1155/2021/8853358>.
- [11] Gad HA, El-Ahmady SH, Abou-Shoer MI, Al-Azizi MM. Application of chemometrics in authentication of herbal medicines: A review. *Phytochem Anal.* 2013; 24: 1–24. <https://doi.org/10.1002/pca.2378>.
- [12] Farag RS, Abdel-Latif MS, Abd El Baky HH, Tawfeek LS. Phytochemical screening and antioxidant activity of some medicinal plants' crude juices. *Biotechnol Rep.* 2020; 28: e00536. <https://doi.org/10.1016/J.BTRE.2020.E00536>.
- [13] Luo D, Shao Y. Classification of Chinese herbal medicine based on improved LDA algorithm using machine olfaction. *Appl Mech Mater.* 2013; 239-240:1532–1536. <https://doi.org/10.4028/www.scientific.net/AMM.239-240.1532>.
- [14] Singh YD, Panda MK, Satapathy KB. Ethnomedicine for drug discovery. *Adv Pharm Biotechnol Recent Prog Futur Appl.* 2020; 15-28. [https://doi.org/10.1007/978-981-15-2195-9\\_2/COVER](https://doi.org/10.1007/978-981-15-2195-9_2/COVER).
- [15] Suryana S, Junaedi EC, Choerunisa Y, Prasatiawati R, Lubis N. Analysis of flavonoid in extract of rose Guava (*Syzygium jambos* (L.) Alston) leaves using Infrared Spectroscopy and chemometrics. *J Phys Conf Ser.* 2019; 1402(5): 1-4. <https://doi.org/10.1088/1742-6596/1402/5/055044>.
- [16] Laily N, Kusumaningtyas RW, Sukarti I, Rini MRDK. The potency of *Guava psidium* Guajava (L.) leaves as a functional immunostimulatory ingredient. *Procedia Chem.* 2015; 14: 301-307. <https://doi.org/10.1016/J.PROCHE.2015.03.042>.
- [17] Zaman W, Ye J, Saqib S, Liu Y, Shan Z, Hao D, Chen Z, Xiao P. Predicting potential medicinal plants with phylogenetic topology: Inspiration from the research of traditional Chinese Medicine. *J Ethnopharmacol.* 2021; 281: 114515. <https://doi.org/10.1016/J.JEP.2021.114515>.
- [18] Rinnan Å, van den Berg F, Engelsen SB. Review of the most common pre-processing techniques for Near-Infrared Spectra. *TrAC - Trends Anal Chem.* 2009; 28: 1201-1222. <https://doi.org/10.1016/j.trac.2009.07.007>.
- [19] Bevilacqua M, Bro R, Marini F, Rinnan Å, Rasmussen MA, Skov T. Recent chemometrics advances for foodomics. *TrAC - Trends Anal Chem.* 2017; 96: 42-51. <https://doi.org/10.1016/J.TRAC.2017.08.011>.
- [20] Brereton RG, Jansen J, Lopes J, Marini F, Pomerantsev A, Rodionova O, Roger JM, Walczak B, Tauler R. Chemometrics in analytical chemistry—Part II: Modeling, validation, and applications. *Anal Bioanal Chem.* 2018; 410: 6691-6704. <https://doi.org/10.1007/s00216-018-1283-4>.
- [21] Sima IA, Andrásí M, Sârbu C. Chemometric assessment of chromatographic methods for herbal medicines authentication and fingerprinting. *J Chromatogr Sci.* 2018; 56: 49-55. <https://doi.org/10.1093/chromsci/bmx080>.
- [22] Nurani LH, Rohman A, Windarsih A, Guntarti A, Riswanto FDO, Lukitaningsih E, Fadzillah NA, Rafi M. Metabolite fingerprinting using 1H-NMR Spectroscopy and chemometrics for classification of three *Curcuma* species from different origins. *Molecules.* 2021; 26: 7626. <https://doi.org/10.3390/molecules26247626>.
- [23] Kumar M, Tomar M, Amarowicz R, Saurabh V, Sneha Nair M, Maheshwari C, Sasi M, Prajapati U, Hasan M, Singh S. Guava (*Psidium Guajava* L.) leaves: Nutritional composition, phytochemical profile, and health-promoting bioactivities. *Foods.* 2021; 10(4): 752. <https://doi.org/10.3390/FOODS10040752>.
- [24] Amat-Ur-rasool H, Symes F, Tooth D, Schaffert LN, Elmorsy E, Ahmed M, Hasnain S, Carter WG. Potential nutraceutical properties of leaves from several commonly cultivated plants. *Biomolecules* 2020; 10(11) : 1556. <https://doi.org/10.3390/BIOM10111556>.