


ISSN: 1411-9420 (print); 2460-1578 (online)

Indonesian Journal of Chemistry

Vol. 24, No. 6, December 2024



Approved by Ministry of Education, Culture,
Research, and Technology
No. 177/E/KPT/2024

Indonesian Journal of Chemistry

ISSN: 1411-9420 (print); 2460-1578 (online)

Published by Chemistry Department Universitas Gadjah Mada



Menu

[Home](#)
[About](#)
[Login](#)
[Register](#)
[Search](#)
[Current](#)
[Archives](#)
[Announcements](#)
[Statistics](#)
[Indexing & Abstracting](#)
[Journal History](#)
[Contact](#)

Home > **Vol 24, No 6 (2024)**

Indonesian Journal of Chemistry

Indonesian Journal of Chemistry is an International, peer-reviewed, open-access journal that publishes original research articles, review articles, and short communication in all chemistry areas, including applied chemistry. The journal is accredited by The Ministry of Education, Culture, Research, and Technology **No.: 177/E/KPT/2024** (in First Rank), indexed by Scopus since 2012 and has a journal impact factor (JIF) score since 2022. In 2018 and 2019 (Volumes 18 and 19, respectively) Indonesian Journal of Chemistry published four issues (numbers) annually (February, May, August, and November). Due to the large number of qualified papers accepted, to speed up the publication of the scientific and valuable research results, since 2020 (Volume 20) Indonesian Journal of Chemistry has increased the frequency of issues to six numbers annually (February, April, June, August, October, and December).



Subscribing on:



ARTICLE IN PRESS

List of the accepted articles for future issues

[Focus & Scope](#)

[Author Guidelines](#)

[Author Fees](#)

[Online Submission](#)

[Publication Ethics](#)

[Plagiarism Policy](#)

[Editorial Board](#)

[Open Access Policy](#)

[Peer Reviewers](#)

Announcements

Publication Frequency and call for paper

After publishing four issues since volume 18 (2018), the number of submitted papers increases significantly. Therefore, to speed up qualified articles to be published internationally, Indonesian Journal of Chemistry publishes six issues (numbers) annually (February, April, June, August, October, and December) since 2020 (Volume 20). Therefore, we invite all authors to submit your qualified manuscripts of original research articles, reviews, short communication in our Journal. Within two months (longest) from submission, the decision of acceptance or rejection has been made. Submission is only via online.

Posted: 2019-12-17

Most Cited Articles

1. WD Fitriana, T Ersam, K Shimizu, S Fatmawati, 2016, Antioxidant Activity of Moringa oleifera Extracts, *Indones. J. Chem.*, 16 (3), 297–301
2. A Rahman, A Ismail, D Jumbianti, S Magdalena, H Sudrajat, 2009, Synthesis of copper oxide nano particles by using *Phormidium cyanobacterium*, *Indones. J. Chem.*, 9 (3), 355–360
3. MD Hidayati, T Ersam, K Shimizu, S Fatmawati, 2017, Antioxidant Activity of *Syzygium polyanthum* Extracts, *Indones. J. Chem.*, 17 (1), 49–53
4. N Syafni, DP Putra, D Arbain, 2012, 3,4-Dihydroxybenzoic Acid and 3,4-Dihydroxybenzaldehyde from the Fern *Trichomanes chinense* L.; Isolation, Antimicrobial and Antioxidant Properties, *Indones. J. Chem.*, 12 (3), 273–278
5. H Winarno, 2009, Benzophenone Glucoside Isolated from the Ethyl Acetate Extract of the Bark of Mahkota Dewa [*Phaleria macrocarpa* (Scheff.) Boerl.] and Its Inhibitory Activity on leukemia L121 Cell Line, *Indones. J. Chem.*, 9 (1), 142–145
6. MT Ekaprasada, H Nurdin, S Ibrahim, D Dachriyanus, 2009, Antioxidant Activity of Methyl Gallate Isolated from the Leaves of *Toona sureni*, *Indones. J. Chem.*, 9 (3), 457–460
7. CS Tang, MH Norziah, 2007, Stability of Betacyanin Pigments from Red Purple Pitaya Fruit (*Hylocereus polyrhizus*): Influence of pH, Temperature, Metal Ions and Ascorbic Acid, *Indones. J. Chem.*, 7 (3), 327–331
8. N Herdyastuti, TJ Raharjo, M Mudasir, S Matsjeh, 2009, Chitinase and Chitinolytic Microorganism: Isolation, Characterization and Potential, *Indones. J. Chem.*, 9 (1), 37–47
9. S Suryati, H Nurdin, D Dachriyanus, MNH Lajis, 2011, Structure Elucidation of Antibacterial Compound from *Ficus deltoidea* Jack Leaves, *Indones. J. Chem.*, 11 (1), 67–70
10. A Hakim, L Liliarsari, A Kadarohman, YM Syah, 2016, Improvement of Student Critical Thinking Skills with the Natural Product

Order Journal

Visitor Statistics

USER

Username

Password

Remember me

Login

JOURNAL CONTENT

Search

Search Scope

All 

Search

Browse

- ▶ By Issue
- ▶ By Author
- ▶ By Title
- ▶ Other Journals

INFORMATION

- Mini Project Laboratory Learning, *Indones. J. Chem.*, 16 (3), 322–328
11. A Ridhay, A Noor, NH Soekamto, T Harlim, I van Altena, 2012, A Stigmasterol Glycoside from the Root Wood of *Melochia umbellata* (Houtt) Stapf var. *degrabrata* K., *Indones. J. Chem.*, 12 (1), 100–103
 12. M Basyuni, N Amri, LAP Putri, I Syahputra, D Arifiyanto, 2017, Characteristics of Fresh Fruit Bunch Yield and the Physicochemical Qualities of Palm Oil during Storage in North Sumatra, Indonesia, *Indones. J. Chem.*, 17 (2), 182–190
 13. A Lesbani, P Tamba, R Mohadi, F Riyanti, 2013, Preparation of Calcium Oxide from *Achatina fulica* as Catalyst for Production of Biodiesel from Waste Cooking Oil, *Indones. J. Chem.*, 13 (2), 176–180
 14. I Fatimah, E Sugiharto, K Wijaya, I Tahir, K Kamalia, 2006, Titanium Oxide Dispersed on Natural Zeolite (TiO₂/Zeolite) and Its Application for Congo Red Photodegradation, *Indones. J. Chem.*, 6 (1), 38–42
 15. D Balarak, FK Mostafapour, 2019, Photocatalytic Degradation of Amoxicillin Using UV/Synthesized NiO from Pharmaceutical Wastewater, *Indones. J. Chem.*, 19 (1), 211–218
 16. K Wahyuningsih, ES Iriani, F Fahma, 2016, Utilization of Cellulose from Pineapple Leaf Fibers as Nanofiller in Polyvinyl Alcohol-Based Film, *Indones. J. Chem.*, 16 (2), 181–189
 17. H Natsir, AR Patong, MT Suhartono, A Ahmad, 2010, Production and Characterization of Chitinase Enzymes from Sulili Hot Spring in South Sulawesi, *Bacillus* sp. HSA,3-1a, *Indones. J. Chem.*, 10 (2), 256–260
 18. E Hermiati, J Azuma, D Manguwidjaja, TC Sunarti, O Suparno, 2011, Hydrolysis of Carbohydrates in Cassava Pulp and Tapioca Flour under Microwave Irradiation, *Indones. J. Chem.*, 11 (3), 238–245
 19. S Sembiring, 2011, Synthesis and Characterisation of Rice Husk Silica Based Borosilicate (B₂SiO₅) Ceramic by Sol-Gel Routes, *Indones. J. Chem.*, 11 (1), 85–89
 20. N Kasanah, T Triyanto, DS Seto, W Amelia, A Isnansetyo, 2015, Antibacterial Compounds from Red Seaweeds (Rhodophyta), *Indones. J. Chem.*, 15 (2), 201–209

(Based on Sinta database)

Posted: 2022-12-04

More Announcements...

Vol 24, No 6 (2024)

Accredited by Ministry of Education, Culture, Research, and Technology No.: 177/E/KPT/2024 (October 15, 2024)

- ▶ For Readers
- ▶ For Authors
- ▶ For Librarians

KEYWORDS

FTIR HPLC Schiff base
adsorption
antibacterial antioxidant
biodiesel catalyst chitosan
eugenol extraction heavy
metals immobilization
kinetics methylene blue
molecular docking
photocatalyst silica silver
nanoparticles synthesis
transesterification

Indones. J. Chem.
indexed by:

Scopus®

Clarivate
Web of Science™





Table of Contents

Articles

- Synthesis of Printed Hollow Fiber Membranes Urea as a Membrane Candidate Hemodialysis** 1583-1601
Muhammad Cholid Djunaidi, Denandha Putri Ayuningrum, Nesti Dwi Maharani, Khabibi Khabibi, Pardoyo Pardoyo, Yanuardi Raharjo, Heru Susanto, Abdullah Malik Islam Filardli
- [10.22146/ijc.83068](https://doi.org/10.22146/ijc.83068) Abstract views : 4187 | views : 1522 | views : 1295
- Synthesis of Activated Carbon from Dragon Fruit Peel for Adsorption of Methyl Blue** 1602-1614
Thi Ngoc Diem Tran, Thi Tuu Tran, Bich Ngoc Hoang, Van Tan Lam, Thi Cam Quyen Ngo
- [10.22146/ijc.85429](https://doi.org/10.22146/ijc.85429) Abstract views : 3103 | views : 1164
- Synthesis of Antimicrobial and Antioxidant Zinc Oxide Hydrogel for Drug Delivery Applications** 1615-1627
Krithika Ramesh, Jeyanthi Ponnusamy, Pazhanisamy Periasamy



 [10.22146/ijc.85663](https://doi.org/10.22146/ijc.85663)  Abstract views : 2990 |  views : 1063

Removal of *para*-Phenylenediamine (PPD) Dye from Its Aqueous Solution by Adsorption Using the Activated Carbon Nanoparticles

1628-1637

Shabaa Fayyad Bdewi, Hanaa Hassan Hussein, Shireen Abdulmohsin Azeez

 [10.22146/ijc.87184](https://doi.org/10.22146/ijc.87184)  Abstract views : 2844 |  views : 1186 |  views : 766

Preparation and Characterization of Encapsulated *Cymbopogon citratus* Essential Oils in Alginate/Chitosan Complexes Using Ion-Gel Technique

1638-1649

Huynh Mai Pham, Thuong Nhan Phu Nguyen, Chi Khang Van, Huynh Cang Mai

 [10.22146/ijc.91251](https://doi.org/10.22146/ijc.91251)  Abstract views : 2979 |  views : 951

Synthesis of Aragonite from Precipitated Calcium Carbonate: A Pilot Scale Study

1650-1660

Ellyta Sari, Reni Desmiarti, Zulhadjri Zulhadjri, Matlal Fajri Alif, Maulana Yusup Rosadi, Syukri Arief

 [10.22146/ijc.92169](https://doi.org/10.22146/ijc.92169)  Abstract views : 3935 |  views : 1301 |  views : 821

Antibacterial Activity and Molecular Docking of Compounds from *Avicennia marina* Leaves Extracts: Obtained by Natural Deep Eutectic Solvents

1661-1674

Hartati Kartikaningsih, Heder Djamaludin, Nanda Audina, Jihan Nur Fauziyah

 [10.22146/ijc.92444](https://doi.org/10.22146/ijc.92444)  Abstract views : 3173 |  views : 1252

Optimizing Dye-Sensitized Solar Cell (DSSC) Performance through Synergistic Natural Dye Combinations from *Beta vulgaris* L., *Curcuma longa* L., and *Pandanus amaryllifolius*

1675-1687

Nita Kusumawati, Pirim Setiarso, Supari Muslim, Nafisatus Zakiyah, Khofifatul Rahmawati, Fadlurachman Faizal Fachrirakarsie

 [10.22146/ijc.93830](https://doi.org/10.22146/ijc.93830)  Abstract views : 3158 |  views : 1259

Immobilization of Cerium(IV) Oxide onto Reduced Graphene Oxide in Epoxy Resin Matrix as Radar Absorbing Composite for X-band Region

1688-1700

Patricya Inggrid Wilhelmina Bolilanga, Rahmat Basuki, Yusuf Bramastya Apriliyanto, Agus Eko Prasojo, Ardyan Lazuardy, Reza Anitasari, Riyanti Putri, Nugroho Adi Sasongko, Arief Budi Santiko



CURRENT ISSUE

ATOM	1.0
RSS	2.0
RSS	1.0

 [10.22146/ijc.94404](https://doi.org/10.22146/ijc.94404)  Abstract views : 3504 |  views : 1258

Synthesis of Water-Soluble Menthol Derivatives Using Response Surface Methodology

1701-1716

Shafaq Nisar, Muhammad Asif Hanif, Umer Rashid, Muhammad Idrees Jilani, Ijaz Ahmad Bhatti, Imtiaz Ali, Bartłomiej Zieniuk

 [10.22146/ijc.94761](https://doi.org/10.22146/ijc.94761)  Abstract views : 3989 |  views : 1240

One-Pot Synthesis and *In Vitro* Studies of Calix[4]-2-methylresorcinarene Derivatives as Antimalarial Agents Against *Plasmodium falciparum* Chloroquine-Resistant Strain FCR-3

1717-1729

Baiq Ike Nursafia, Yehezkiel Steven Kurniawan, Jumina Jumina, Harno Dwi Pranowo, Eti Nurwening Sholikhah, Jeffry Julianus, Susalit Setya Wibowo, Hana Anisa Fatimi, Yoga Priastomo, Krisfian Tata Aneka Priyangga

 [10.22146/ijc.94885](https://doi.org/10.22146/ijc.94885)  Abstract views : 3116 |  views : 1119

Preparation, Structural Identification, and Biomedical Evaluation of Some New Complexes

1730-1742

Alyaa Khider Abbas, Asmaa Edrees Fadhil

 [10.22146/ijc.95759](https://doi.org/10.22146/ijc.95759)  Abstract views : 2767 |  views : 1000 |  views : 716

New Insights of Response Surface Methodology Approach in Optimizing Total Phenolic Content of *Zanthoxylum acanthopodium* DC. Fruit Extracted Using Microwave-Assisted Extraction and the Impact to Antioxidant Activity

1743-1759

Sumaiyah Sumaiyah, Retno Murwanti, Didi Nurhadi Illian, Muhammad Fauzan Lubis, Keshia Tampubolon

 [10.22146/ijc.95922](https://doi.org/10.22146/ijc.95922)  Abstract views : 3221 |  views : 1171

Synthesis and Characterization of Aluminosilicate Catalysts from Volcano Mud for Biofuel Production with Different Feedstocks

1760-1771

Hartati Hartati, Qurrota A'yuni, Nita Safira Dewi, Putri Bintang Dea Firda, Adiba Naila Izzah, Didik Prasetyoko, Harmami Harmami, Shahrul Nizam Ahmad

 [10.22146/ijc.96149](https://doi.org/10.22146/ijc.96149)  Abstract views : 3468 |  views : 1301

Volatilomics Profiling of Counterfeit Perfume by Gas Chromatography Hyphenated to Mass Spectrometry and Fourier-Transformed Infrared Spectroscopy


1772-1783

and Fourier Transformed Infrared Spectroscopy*Siti Nurul Hidayah, Artania Adnin Tri Suma, Endang Lukitaningsih*doi [10.22146/ijc.96313](https://doi.org/10.22146/ijc.96313)  Abstract views : 3466 |  views : 1495**Magnetically Active GO-Fe₃O₄ Nanocomposite for Enhanced Rhodamine B Removal Efficiency**

1784-1796

*Alexander Souhuat, Henry Fonda Aritonang, Harry Steven Julius Koleangan*doi [10.22146/ijc.96383](https://doi.org/10.22146/ijc.96383)  Abstract views : 3035 |  views : 1098**Forensic Discrimination of Lipstick Stains Using Attenuated Total Reflectance-Fourier Transform Infrared Spectroscopy and Pattern Recognition Techniques**


1797-1810

*Mohd Afiq Mohd Azis, Naji Arafat Mahat, Hasmerya Maarof, Sarah Aina Azman*doi [10.22146/ijc.97097](https://doi.org/10.22146/ijc.97097)  Abstract views : 2901 |  views : 1193**Physicochemical Characterization of Chitosan-Multiwalled Carbon Nanotubes Hybrid Material for Ofloxacin Delivery**

1811-1823

*Zainab Jasim Khudair, Zeina Mohammad Kadam*doi [10.22146/ijc.97160](https://doi.org/10.22146/ijc.97160)  Abstract views : 2853 |  views : 1056**Analysis of Hg(II) Metal by Voltammetric Method Using a Carbon Electrode Modified with EDTA from Used Tire Waste**

1824-1835

*Sukoma Sukoma, Sagir Alva, Eka Safitri, Julinawati Julinawati, Mustanir Mustanir, Syafrizal Fonna, Muhammad Ihsan, Fitri Dara, Meri Dayanti, Ahmad Kamal Arifin, Khairi Suhud*doi [10.22146/ijc.97561](https://doi.org/10.22146/ijc.97561)  Abstract views : 3339 |  views : 1113**A Comparative Study of LiNCA Cathode Recycled from Spent Lithium-Ion Batteries and Synthesized from Metal Precursor**

1836-1850

*Arif Jumari, Enni Apriliani, Cornelius Satria Yudha, Agus Purwanto, Anne Zulfia Syahrial, Wara Dyah Pita Rengga*doi [10.22146/ijc.98276](https://doi.org/10.22146/ijc.98276)  Abstract views : 2991 |  views : 1062**Synthesis, Characterization, Biological, and Antioxidant Activity of New Metal Ion Complexes with Schiff Base Derived from 2-Hydroxybenzohydrazide**

1851-1860


Ayyat Jawad Kadhim, Naser Shaalan

 [10.22146/ijc.98292](https://doi.org/10.22146/ijc.98292)  Abstract views : 3225 |  views : 1229 |  views : 770

Short Communication

LC-HRMS-Based Metabolomics Approach Reveals Antioxidant Compounds from *Centella asiatica* Leaves Extracts 1861-1869


Riva Silvia, Wulan Tri Wahyuni, Eti Rohaeti, Siti Aisyah, Dewi Anggraini Septaningsih, Alfi Hudatul Karomah, Mohamad Rafi

 [10.22146/ijc.90782](https://doi.org/10.22146/ijc.90782)  Abstract views : 4227 |  views : 3417

Review

Determination of Chlorpromazine Using Molecular Imprinting Polymers in Different Sample Matrices 1870-1882

Eman Wajeh Ammen, Yehya Kamal Al-Bayati

 [10.22146/ijc.94705](https://doi.org/10.22146/ijc.94705)  Abstract views : 3001 |  views : 1087

Phytochemistry and Biological Activities of *Amomum* Species 1883-1905

Deden Indra Dinata, Rani Maharani, Fauzan Zein Muttaqin, Unang Supratman

 [10.22146/ijc.95402](https://doi.org/10.22146/ijc.95402)  Abstract views : 4182 |  views : 1464

Review on Melanin Application as an Antibacterial and Antioxidant Agent in Food Packaging 1906-1920

Zainal Mustakim, Agus Prasetya, Joko Wintoko, Fitri Nur Kayati, Chandra Wahyu Purnomo

 [10.22146/ijc.95550](https://doi.org/10.22146/ijc.95550)  Abstract views : 3249 |  views : 1203

03308679 View The Statistics of Indones. J. Chem.

Indonesian Journal of Chemistry

ISSN: 1411-9420 (print); 2460-1578 (online)

Published by Chemistry Department Universitas Gadjah Mada



Menu

[Home](#) [About](#) [Login](#) [Register](#) [Search](#) [Current](#) [Archives](#) [Announcements](#) [Statistics](#) [Indexing & Abstracting](#) [Journal History](#) [Contact](#)

[Home](#) > [About the Journal](#) > [Editorial Team](#)

Editorial Team

Editor-in-Chief

Prof. Nuryono Nuryono, Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Gadjah Mada, Indonesia

Managing Editor

Dr. Niko Prasetyo, Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Gadjah Mada, Indonesia

Editorial Board

Dr. Minh-Vien Le, Ho Chi Minh city University of Technology (HCMUT), VNU-HCM, Viet Nam

Dr. Sadhan Jana, Cornell University, New York, USA, United States

Dr. Nurul Izza binti Taib, Faculty of Applied Sciences, Universiti Teknologi MARA, Perak Branch, Tapah, Malaysia

Dr. Faisal Mukhtar, Institute of Physics, The Islamia University of Bahawalpur, Pakistan

Prof. Indriana Kartini, Laboratory of Inorganic Chemistry, Department of Chemistry, Universitas Gadjah Mada, Indonesia

Hisao Kokusen, Department of Chemistry, Tokyo Gakugei University, Japan

Mr. Arif Fadlan, Department of Chemistry Institut Teknologi Sepuluh Nopember, Indonesia

Prof. Endang T. Wahyuni, Universitas Gadjah Mada, Indonesia

Subscribing on:



ARTICLE IN PRESS

List of the accepted articles for future issues

[Focus & Scope](#)

[Author Guidelines](#)

[Author Fees](#)

[Online Submission](#)

[Publication Ethics](#)

[Plagiarism Policy](#)

[Editorial Board](#)

[Open Access Policy](#)

[Peer Reviewers](#)

Prof. Bambang Purwono, Department of Chemistry, Universitas Gadjah Mada, Indonesia
 Dr. Wahyu Satpriyo Putro, National Institute of Advanced of Industrial Science and Technology, Japan
 Dr. Ramesh Katla, Research Scientist, Department of Hematology/Oncology, at the University of Virginia, Charlottesville, VA, United States, Zip: 22903., United States
 Dr. Satya Candra Wibawa Sakti, Laboratory of Inorganic Chemistry, Department of Chemistry, Universitas Airlangga, Indonesia
 Dr. Stalis Norma Ethica, Universitas Muhammadiyah Semarang, Indonesia
 Dr. Wahyu Dita Saputri, National Research and Innovation Agency, Indonesia
 Dr. Maulidan Firdaus, Department of Chemistry, Universitas Sebelas Maret, Indonesia
 Prof. Belkheir Hammouti, Laboratory of Applied Analytical Chemistry, Materials and Environment (L2ACME), Mohamed First University, Faculty of Science, B.P. 717, 60000 Oujda, Morocco
 Asst. Prof. Dr. Vy Anh Tran, Department of Chemical and Biochemical Engineering, Gachon University, Korea, Republic of
 Asst. Prof. Dr. Mahmood Ahmed, Department of Chemistry, Division of Science and Technology, University of Education, College Road, Lahore, Pakistan
 Dr. Charles Edwin Raja Gabriel, University of Notre Dame, Notre Dame, Indiana, United States
 Assoc. Prof. Dr. Roswanira Abdul Wahab, Department of Chemistry, Universiti Teknologi Malaysia, Malaysia
 Prof. Joe da Costa, Department of Chemical Engineering, University of Queensland, Australia
 Dr. Maurizio Barbieri, Department of Earth Science, Sapienza University of Rome, Italy
 Ming Cai, Tongji University, China
 Muhammad Idham Darussalam Mardjan, Laboratory of Organic Chemistry, Department of Chemistry, Universitas Gadjah Mada, Indonesia
 Prof. Pornthep Sompornpisut, Department of Chemistry, Chulalongkorn University, Thailand
 Prof. Dr. Taghreed Hashim Al-Noor, Ibn-Al-Haitham Education College /University of Baghdad, Iraq
 Assoc. Prof. Dr. Praveen Kumar Sharma, Department of Chemistry, Lovely Professional University, Punjab, India
 Dr. Saprizal Hadisaputra, Chemistry Education Division, Faculty of Science and Education, University of Mataram, Indonesia
 Dr. Winifred Uduak Anake, Department of Chemistry, College of Science and Technology, Covenant University, Nigeria

Editing staffs

Djoko Prihandono, Department of Chemistry, Universitas Gadjah Mada, Indonesia
 Tantiana Indriani, Indonesia
 Ervan Yudha, Department of Chemistry, Universitas Gadjah Mada, Indonesia, Indonesia
 Selma Wulandari, Universitas Gadjah Mada, Indonesia

Treasurer

Nurzanah Hidayanti, Department of Chemistry, Universitas Gadjah Mada, Indonesia

Order Journal

Visitor Statistics

USER

Username

Password

Remember me

Login

JOURNAL CONTENT

Search

Search Scope

All 

Search

Browse

- ▶ By Issue
- ▶ By Author
- ▶ By Title
- ▶ Other Journals

INFORMATION

Indonesian Journal of Chemistry (ISSN 1411-9420 /e-ISSN 2460-1578) - Chemistry Department, Universitas Gadjah Mada, Indonesia.

03308683 [View The Statistics of Indones. J. Chem.](#)

- ▶ For Readers
- ▶ For Authors
- ▶ For Librarians

KEYWORDS

FTIR HPLC Schiff base
adsorption
antibacterial antioxidant
biodiesel catalyst **chitosan**
eugenol extraction heavy
metals immobilization
kinetics methylene blue
molecular docking
photocatalyst silica silver
nanoparticles synthesis
transesterification

Indones. J. Chem.
indexed by:

Scopus®

Clarivate
Web of Science™





Indonesian Journal of Chemistry

Q3

Chemistry (miscellaneous)

best quartile

SJR 2023

0.26



powered by scimagojr.com



CURRENT ISSUE

ATOM	1.0
RSS	2.0
RSS	1.0

One-Pot Synthesis and *In Vitro* Studies of Calix[4]-2-methylresorcinarene Derivatives as Antimalarial Agents Against *Plasmodium falciparum* Chloroquine-Resistant Strain FCR-3

Baiq Ike Nursafia¹, Yehezkiel Steven Kurniawan¹, Jumina Jumina^{1*}, Harno Dwi Pranowo¹, Eti Nurwening Sholikhah², Jeffry Julianus³, Susalit Setya Wibowo⁴, Hana Anisa Fatimi⁵, Yoga Priastomo⁶, and Krisfian Tata Aneka Priyangga¹

¹Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Gadjah Mada, Sekip Utara, Yogyakarta 55281, Indonesia

²Department of Pharmacology and Therapeutics, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Sekip Utara, Yogyakarta 55281, Indonesia

³Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Universitas Sanata Dharma, Yogyakarta 55282, Indonesia

⁴Research Center for Process and Manufacturing Industry Technology, National Research and Innovation Agency (BRIN), KST BJ Habibie, Banten 15314, Indonesia

⁵Pharmacy Study Program, Faculty of Mathematics and Natural Science, Universitas Sebelas Maret, Jl. Ir. Sutami 36A, Surakarta 57126, Indonesia

⁶Department of Chemical Education, Faculty of Mathematics and Natural Science, Universitas Negeri Yogyakarta, Jl. Colombo No. 1, Yogyakarta 55281, Indonesia

* **Corresponding author:**

email: jumina@ugm.ac.id

Received: April 2, 2024

Accepted: August 5, 2024

DOI: 10.22146/ijc.94885

Abstract: Malaria is an endemic disease in Indonesia caused by infection from the *Plasmodium* parasite. Recently, antimalarial resistance significantly contributed to the decline in the cure rate of malaria sufferers. In this work, three calix[4]resorcinarenes have been synthesized from 2-methylresorcinol and different benzaldehyde derivatives, i.e., 4-chlorobenzaldehyde, 4-methoxybenzaldehyde, and 4-dimethylaminobenzaldehyde through the one-pot synthesis procedure. The calix[4]resorcinarenes synthesis was done through a cyclo-condensation reaction by using HCl 37% as the catalyst and ethanol as the solvent in an one-pot reaction. The structures of the synthesized products were confirmed using Fourier transform infrared, proton-nuclear magnetic resonance, and liquid chromatography-mass spectrometry techniques. The antimalarial activity assay was evaluated against the *Plasmodium falciparum* FCR-3 strain through an in vitro study. Three synthesized compounds, i.e., C-4-chlorophenylcalix[4]-2-methylresorcinarene, C-4-methoxyphenylcalix[4]-2-methylresorcinarene and C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene have been successfully synthesized in up to 97% yield. The C-4-chlorophenylcalix[4]-2-methylresorcinarene exhibited the most potent antimalarial activity with a half-maximal inhibitory concentration (IC_{50}) value of 2.66 μ M against *P. falciparum* FCR-3 while the C-4-methoxyphenylcalix[4]-2-methylresorcinarene and C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene gave the IC_{50} values of 23.63 and 13.82 μ M, respectively. From the results, it could be concluded that the antimalarial activity of calix[4]-2-methylresorcinarenes was influenced by the type of substituent of aromatic rings at the para position.

Keywords: Antimalarial; calix[4]-2-methylresorcinarene; in vitro assay; one-pot synthesis; *Plasmodium falciparum*

■ INTRODUCTION

Malaria is one of the most fatal diseases, which is an easily spread parasite disease globally. In 2022, as many as 241 million global active malaria cases and 627 thousand deaths due to malaria have been reported. Compared to the reported data in 2019, there is an increase of 22 million active cases and 50 thousand deaths. Southeast Asia had nine countries with malaria endemic areas with 5 million cases or equal to 2.1% of global cases. Among these countries, Indonesia is the only one that reported an increase in malarial deaths in 2020. In 2019, Indonesia reported 32 deaths; however, this number increased to 49 deaths in 2020. Additionally, Indonesia is the second country in South and Southeast Asia regions with the highest number of malaria active cases after India in 2020. Indonesia reported 800 thousand malaria cases and 49 deaths in 2020, accounting for 15.6% of active cases and 22% of malaria deaths in the Southeast Asia region [1].

It is reported that malaria disease is caused by the Plasmodium infection through the mosquito (*Anopheles quadrimaculatus* Say) host. Four Plasmodium species, i.e., *P. falciparum*, *P. vivax*, *P. ovale*, and *P. malariae*, are responsible for the malaria diseases. The *P. falciparum* and *P. vivax* are the predominant species, accounting for 60 and 40% of global malaria cases. On the other hand, *P. ovale* is mostly found in the Sub-Saharan Africa region, while *P. malariae* is widely found in the South American region. Among them, *P. falciparum* has been reported as the most predominant species in global cases with serious symptoms, including acute respiratory dysfunction, anemia, renal failure, central nervous system damage, fever, headache, and chills [2]. Some antimalarial drugs are commercially available nowadays, but most of them are resistant to malaria parasites, including chloroquine diphosphate [3] and artemisinin [4]. Artemisinin-based combination treatments (ACTs) are currently used for malaria therapy in treating malarial disease caused by the *P. falciparum* parasite [5]. The ACT treatment consists of artemisinin and other effective malaria drugs, which can suppress artemisinin resistance. However, the effectiveness of this treatment continues to decline due to the domino effects of drug resistance and undesired side effects on human health [6]. Therefore, there is an urgent

need to discover new antimalarial drugs as soon as possible to overcome these issues.

Hundreds of antimalarial agents have been isolated from natural sources as well as synthesized by organic chemists. In general, natural antimalarial agents are superior because of their safety to the human body. However, these natural compounds often suffer from poor antimalarial activity with complicated isolation and/or purification procedures. On the other hand, synthetic antimalarial agents offer higher antimalarial activity due to their rational design and higher purity than natural compounds. Recently, interest in macrocycle-based compounds has increased in medicinal chemistry, especially for calixarene. Calix[4]resorcinarene, a calixarene derivative, has gained significant interest in supramolecular chemistry as the basic framework for synthesizing cations, anions, biomolecules, gas, and neutral host molecules [7]. Calix[4]resorcinarenes have been employed in various applications, such as adsorbents [8], surfactants [9], complexation agents [10], liquid membranes [11], and chemical sensors [12]. In addition, calix[4]resorcinarenes are well known to exhibit several biological activities and uses in analytical and pharmaceutical fields [13], such as anticancer and antioxidant agents [14-15]. However, the exploration of calix[4]resorcinarene derivatives has yet to be thoroughly carried out as an active drug ingredient, especially as an antimalarial agent.

According to the published literature, the presence of aromatic pharmacophoric groups is critical to enhance the antimalarial activity of organic compounds. Chloro substituent in the para position of the quinoxaline's aromatic ring gave high antimalarial activity with a half-maximal inhibitory concentration (IC_{50}) value of 0.20 $\mu\text{g/mL}$, which is more active than the chloroquine diphosphate (IC_{50} value = 0.43 $\mu\text{g/mL}$) [16]. The methoxy group is also pivotal for the antimalarial activity of hydroxy-substituted chalcones with IC_{50} values less than 10 $\mu\text{g/mL}$ [17]. On the other hand, the alkylamino group in chalcone derivatives exhibits high antimalarial activity with IC_{50} values below 2 $\mu\text{g/mL}$ [18]. Our previous work evaluated the antimalarial activity of

calix[4]resorcinarenes with methyl, heptyl, and nitrophenyl functional groups through an *in vitro* heme polymerization inhibition assay. The experimental data revealed that calix[4]resorcinarenes with methyl, heptyl, and nitrophenyl functional groups yielded lower IC₅₀ values of 0.282, 0.814, and 0.198 µg/mL, respectively, which were lower than of chloroquine diphosphate (IC₅₀ value = 1.157 µg/mL) [19]. This result demonstrated that calix[4]resorcinarene with aromatic moiety gave a higher antimalarial activity than the calix[4]resorcinarene with alkyl groups.

In another work, we also evaluated the antimalarial activity of calix[4]pyrogallolarenes through an *in vitro* heme polymerization inhibition assay. The experimental data revealed that calix[4]pyrogallolarenes with phenyl, 4-hydroxy-3-methoxyphenyl, and chlorophenyl functional groups yielded lower IC₅₀ values of 1.268, 1.029, and 0.238 µg/mL, respectively, which were lower than of chloroquine diphosphate as the positive control [20]. This result demonstrated that calix[4]pyrogallolarenes yielded a weaker antimalaria activity than calix[4]resorcinarenes. This phenomenon might be caused by the stronger intramolecular hydrogen bonds in calix[4]pyrogallolarenes. The stronger intramolecular hydrogen bonds yield a weaker capacity of electron donating and/or withdrawing groups to attack the Plasmodium. Furthermore, the ability of methoxy and chloro groups to yield a stronger antimalaria has been confirmed compared with the unsubstituted aromatic ring, i.e., the phenyl group. To approve our hypothesis, we have examined the antimalarial activity of calix[4]-2-methylresorcinarenes with the nitro group at ortho and para positions at the aromatic rings through *in vitro* antiplasmodial method. The experimental data showed that the calix[4]-2-methylresorcinarenes with the nitro group at the para position gave a lower IC₅₀ value (1.79 µM) than the calix[4]-2-methylresorcinarenes with the nitro group at ortho position (IC₅₀ value = 2.35 µM) against *P. falciparum* [21]. These reports suggested that chloro, methoxy, and alkylamino functional groups in the para position of calix[4]-2-methylresorcinarene framework shall be considered to have potent antimalarial activity.

Therefore, this study aims to obtain potential antimalarial candidates from calix[4]-2-methylresorcinarene derivatives with various functional groups of chloro, methoxy, and dimethylamino groups at the para position. Three calix[4]-2-methylresorcinarenes were synthesized from aromatic precursors, namely 2-methylresorcinol with 4-chlorobenzaldehyde, 4-methoxybenzaldehyde, and 4-dimethylaminobenzaldehyde in a one-pot synthesis method under acidic conditions. The chemical structure of all synthetic products was elucidated using Fourier transform infrared (FTIR), nuclear magnetic resonance (NMR), and liquid chromatography-mass spectrometry (LC-MS) techniques. Furthermore, the synthetic compounds were examined for their antimalarial activity against *P. falciparum* chloroquine-resistant strain (FCR-3) through an *in vitro* assay. The effect of the various functional groups on the antimalarial activity was also discussed from a structure-activity relationship point of view.

■ EXPERIMENTAL SECTION

Materials

Materials employed for the synthesis were 2-methylresorcinol (C₇H₈O₂), 4-chlorobenzaldehyde (C₇H₅ClO), 4-methoxybenzaldehyde (C₈H₈O₂), 4-(dimethylamino)benzaldehyde (C₉H₁₁NO), hydrochloric acid (HCl) 37%, and ethanol (C₂H₅OH). These reagents were purchased from Merck in pro-analytical grade and used without further purification. The materials used in the antimalarial evaluation were *P. falciparum* strain FCR-3, dimethyl sulfoxide (DMSO, C₃H₆SO, Merck), methanol (CH₃OH, Merck), serum blood, RPMI-1640 (Sigma-Aldrich), red blood cells (RBC), Giemsa 10% solution (Merck), and distilled water.

Instrumentation

The melting point of the synthesized calix[4]resorcinarene was determined by a Electrothermal 9100 apparatus. Structure elucidation of the synthesized products was performed using an FTIR spectrometer (Shimadzu Prestige 21), proton NMR spectrometer (¹H-

NMR, JEOL JNMECA 500 MHz), and LC-MS instrument (ACQUITY UPLC®H-Class System waters, USA). Pieces of equipment used for the antimalarial assay were glass slides, 96-well microplate, micropipette (Gilson and Thermo Scientific), candle jar, microcentrifuge (Thermo Sorvall Legend Micro 17R), and microscope (Nikon eclipse E100).

Procedure

One-pot synthesis of calix[4]-2-methylresorcinarene

The synthesis procedure of calix[4]-2-methylresorcinarene was carried out using an adapted method from the previously published literature [22]. The 2-methylresorcinol (5 mmol) was completely dissolved in 50 mL of ethanol and then separately mixed with benzaldehyde derivatives, i.e., 4-chlorobenzaldehyde, 4-methoxybenzaldehyde, 4-dimethylaminobenzaldehyde (5 mmol). Afterward, as much as 1 mL of HCl 37% was slowly poured into the mixture and the mixture was heated at 78 °C. After 24 h, the mixture was cooled and added with 50 mL of distilled water to precipitate the desired products. Next, the solid residue was filtered and washed with cold ethanol and water to remove the unreacted precursors. Then, the solid product was recrystallized from ethanol to obtain the title compound as a pure chemical and finally characterized by FTIR, ¹H-NMR, and LC-MS analyses.

C-4-chlorophenylcalix[4]-2-methylresorcinarene.

The desired compound was obtained in 97% yield as an orange solid. melting point (m.p.) 263 °C; FTIR ν_{\max} (KBr, cm^{-1}): 3441 (O–H stretching), 2924 and 2862 (C_{sp^3} –H stretching), 1604 and 1481 (C=C aromatic stretching), 1435 (C–H methine stretching), 1288 and 1188 (C–O stretching), 725 (C–Cl stretching); ¹H-NMR (DMSO-*d*₆; 500 MHz) δ (ppm): 1.95 and 1.97 (s, 12H, ArCH₃), 5.71, 5.76 and 5.96 (s, 4H, ArCH), 5.99 and 6.23 (s, 8H, ArH meta to OH), 6.51 and 6.62 (d, coupling constant (*J*) = 8.6 Hz, 8H, ArH), 6.91 and 7.05 (d, *J* = 8.5 Hz, 8H, ArH), 7.79, 7.88, 7.99, and 8.17 (s, 8H, ArOH); LC-MS: two signals at 12.81 and 13.14 min with *m/z* of $[\text{M}+\text{K}+\text{H}]^+ = 1004$.

C-4-methoxyphenylcalix[4]-2-methylresorcinarene.

The title molecule was obtained in 95% yield as an orange solid. m.p. > 367 °C (decomposition); FTIR ν_{\max} (KBr, cm^{-1}): 3487 (O–H stretching), 2931 and 2839 (C_{sp^3} –H

stretching), 1604 and 1512 (C=C aromatic stretching), 1473 (C–H methine stretching), 1242 and 1180 and 1033 (C–O stretching); ¹H-NMR (CDCl₃; 500 MHz) δ (ppm): 2.06 (s, 12H, ArCH₃), 3.77 (s, 12H, OCH₃), 4.54 (s, 8H, ArOH), 5.37 (s, 4H, ArCH), 5.58 (s, 4H, ArH meta to OH), 6.62 (d, 8H, *J* = 8.5 Hz, ArH), 6.70 (d, 8H, *J* = 9.0 Hz, ArH); LC-MS: a single signal at 10.81 min with *m/z* of $[\text{M}+\text{NH}_4]^+ = 986$.

C-4-dimethylaminophenylcalix[4]-2-

methylresorcinarene. The product was obtained in 35% yield as a purple solid. m.p. > 289 °C (decomposition); FTIR ν_{\max} (KBr, cm^{-1}): 3394 (O–H stretching), 2924 (C_{sp^3} –H stretching), 1604 and 1473 (C=C aromatic stretching), 1435 (C–H methine stretching), 1180 (C–O stretching), 1342 (C–N stretching); ¹H-NMR (D₂O; 500 MHz) δ (ppm): 1.92 and 1.97 (s, 12H, ArCH₃), 3.09 and 3.16 (s, 24H, –N(CH₃)₂), 5.62, 5.71 and 5.86 (s, 4H, ArCH), 5.77, 5.89 and 5.92 (s, 4H, meta to OH), 6.88–6.79 (d, 8H, *J* = 8.5 Hz, ArH), 7.11–7.37 (d, 8H, *J* = 8.5 Hz, ArH); LCMS: two signals at 8.55 and 9.35 min with *m/z* of $[\text{M}]^+ = 1021$.

Antimalarial activity assay

The synthesized calix[4]-2-methylresorcinarene derivatives were evaluated for their antimalarial activity against *P. falciparum* FCR-3 as a chloroquine-resistant strain. In this study, the antimalarial activity evaluation was conducted using an *in vitro* hybrid microtechnical method as reported by Congpoung et al. [23]. As much as 100 μL of inoculum solution containing the *P. falciparum* FCR-3 parasite in serum blood and red blood cells was added to each well of the 96-well assay plate. Several concentrations of calix[4]-2-methylresorcinarene compounds (1, 2, 10, 20, and 25 $\mu\text{g}/\text{mL}$) in DMSO were prepared. Then, as much as 100 μL of each solution concentration was added to the 96-well assay plate, in which each concentration was subjected to triplicates in a row. The incubation was carried out for 72 h in a 5% CO₂ incubator at 37 °C, using the previously reported procedure [24]. Afterward, the blood suspension was subjected to a vortex separation to obtain supernatant and blood precipitate. The blood precipitate was placed in the glass slide to make a thin blood smear. The blood smear was fixed through

methanol washing and then stained by using a 10% Giemsa solution containing methylene blue, eosin, and azure B dyes. These dyes were added for better visualization of health and infected red blood cells by *P. falciparum* parasites. The growth of *P. falciparum* was observed during the schizont phase. The number of schizonts was counted by using a microscope and then compared for at least 1,000 normal red blood cells. The inhibition percentage is calculated by using Eq. (1);

$$\text{Inhibition (\%)} = \frac{A - B}{A} \times 100\% \quad (1)$$


where A and B represent the negative control and sample parasitemia percentage, respectively. The IC_{50} value of each synthesized compound was calculated through a probit analysis using IBM SPSS Statistics 26 software. Chloroquine diphosphate and DMSO were employed as the positive and negative controls, respectively.

RESULTS AND DISCUSSION

One-pot Synthesis of Calix[4]-2-methylresorcinarenes Derivatives

All of the synthesized compounds are produced through a cyclo-condensation reaction between 2-methylresorcinol and benzaldehyde derivatives. This cyclo-condensation reaction produced a cyclic tetramer of calix[4]-2-methylresorcinarene, as shown in Scheme 1. The reaction mechanism for the production of calix[4]-2-methylresorcinarene has been investigated in our previous work for calix[4]resorcinarenes and calix[4]pyrogallolarenes [19]. Due to the similar structure of calix[4]resorcinarenes, calix[4]pyrogallolarenes, and calix[4]-2-methylresorcinarenes, the reaction mechanism

shall not be significantly different. Briefly, the carbonyl group of benzaldehyde derivative was protonated by hydrochloric acid in ethanol media, thus increasing the electrophilicity properties as reported before. The 2-methylresorcinol has three electron-donating groups, i.e., two hydroxyl groups and one methyl group. Therefore, 2-methylresorcinol could act as a nucleophile for an electrophilic aromatic substitution reaction with activated benzaldehyde ones. This coupling reaction happens until a cyclic tetramer structure is formed through the elimination of water molecules. Through this reaction mechanism, three calix[4]-2-methylresorcinarene derivatives, i.e., C-4-chlorophenylcalix[4]-2-methylresorcinarene, C-4-methoxyphenylcalix[4]-2-methylresorcinarene, and C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene have been successfully obtained in 97, 95, and 35% yield, respectively. The C-4-chlorophenylcalix[4]-2-methylresorcinarene compound was obtained in the highest yield due to the presence of chloro substituent as an electron-withdrawing group, thus contributing to the higher electrophilicity properties of benzaldehyde. Meanwhile, C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene compound was obtained in the lowest yield due to the presence of the polar dimethylamino group. This C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene compound has eight hydroxyl groups and four dimethylamino groups; thus, it is highly soluble in water and ethanol media. Consequently, this high solubility might lead to the lowest isolation yield during the recrystallization process.



Scheme 1. The one-pot synthesis route of calix[4]-2-methylresorcinarene derivatives

All of the synthetic calix[4]-2-methylresorcinarene products were elucidated using FTIR, $^1\text{H-NMR}$, and LC-MS analyses. The FTIR spectra of C-4-chlorophenylcalix[4]-2-methylresorcinarene, C-4-methoxyphenylcalix[4]-2-methylresorcinarene, and C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene are shown in Fig. 1. In the FTIR spectrum of each calix[4]-2-methylresorcinarene, the formation of calix[4]-2-methylresorcinarene structure was indicated by the loss of strong absorption of the carbonyl group (C=O) stretching and C-H stretching of benzaldehyde at 1700–1660 and 2700–2800 cm^{-1} , respectively [25]. The appearance of the methine bridge (C-H) absorption at 1435–1437 cm^{-1} strengthened the evidence of the formation of calix[4]-2-methylresorcinarene as this absorption signal is quite characteristic of calix[4]resorcinarene framework [21]. The other functional groups such as hydroxyl (O-H), $\text{C}_{\text{sp}^3}\text{-H}$ of methylresorcinol structure, C=C aromatic and C-O phenolic were also observed in 3487–3394, 2931–2839, 1604–1473, and 1288–1180 cm^{-1} , respectively, giving a stronger evidence that the calix[4]-2-methylresorcinarene structure has been successfully generated. The presence of these functional groups has also been reported for a successful synthesis of calixarenes, calix[4]resorcinarenes, and calix[4]pyrogallolarenes in other published works [7,10-11,15,22]. Additionally, the presence of the C-Cl functional group in C-4-chlorophenylcalix[4]-2-methylresorcinarene was confirmed by an absorption signal at 725 cm^{-1} on its FTIR spectrum. The C-O methoxy functional group of C-4-methoxyphenylcalix[4]-2-methylresorcinarene was also observed at 1033 cm^{-1} , while the C-N functional group of C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene was detected as a signal at 1342 cm^{-1} . These FTIR data indicated the successful production of calix[4]-2-methylresorcinarene derivatives.

The $^1\text{H-NMR}$ spectra of C-4-chlorophenylcalix[4]-2-methylresorcinarene, C-4-methoxyphenylcalix[4]-2-methylresorcinarene, and C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene are shown in Fig. 2. In $^1\text{H-NMR}$ analysis, the C-H aldehyde signal of benzaldehyde at around 10 ppm was not found due to the cyclo-condensation reaction. Instead, new singlet




Fig 1. The FTIR spectra of (a) C-4-chlorophenylcalix[4]-2-methylresorcinarene, (b) C-4-methoxyphenylcalix[4]-2-methylresorcinarene, and (c) C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene

peaks from the methine proton at 5.37–5.76 ppm gave solid evidence for the successful formation of the calix[4]-2-methylresorcinarene [26]. It was well-known that the $^1\text{H-NMR}$ characterization data are the key to elucidating the supramolecular structures, including calix[4]resorcinarenes, calixarenes, and calix[4]pyrogallolarenes. Ohto [8] reported that there are four possible conformations of calix[4]resorcinarene, i.e., crown (C_{4v}), boat (C_{2v}), chair (C_{2h}), diamond (C_s), and saddle (S_4) conformations. Crown conformation of calix[4]resorcinarene is defined when a singlet signal of methine proton is observed in the $^1\text{H-NMR}$ spectrum. Boat and chair conformations of calix[4]resorcinarene are interpreted when there are two methine proton signals with the same integral value. On the other hand, diamond and saddle conformation of calix[4]resorcinarene are clarified when there are four methine proton signals with the same integral value [27-29]. This spectroscopic evidence aligns with the crystallographic data of each conformation, demonstrating the meaningful $^1\text{H-NMR}$ characterization data as proved by previous works [30-34]. With only the difference of a methyl group between the hydroxyl groups of calix[4]resorcinarene, the existing conformations of calix[4]-2-methylresorcinarene and calix[4]resorcinarene are not significantly different.




Fig 2. The $^1\text{H-NMR}$ spectra of (a) C-4-chlorophenylcalix[4]-2-methylresorcinarene, (b) C-4-methoxyphenylcalix[4]-2-methylresorcinarene, and (c) C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene

The $^1\text{H-NMR}$ spectrum analysis showed that C-4-chlorophenylcalix[4]-2-methylresorcinarene and C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene compounds form two conformations due to variable possible intramolecular and intermolecular hydrogen bonds. The C-4-chlorophenylcalix[4]-2-methylresorcinarene compound existed in boat and chair conformations, while C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene compound observed in the crown and chair conformations. Boat and chair conformations of C-4-chlorophenylcalix[4]-2-methylresorcinarene compound were strongly indicated by four singlet proton signals of resorcinol OH group at a chemical shift value of 7.79, 7.88, 7.99, and 8.17 ppm. The boat and chair conformations have OH groups divided in two directions of different chemical environments, axial and equatorial; thus, these hydroxyl groups appeared as two proton signals [35-36]. The C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene compound existed in crown and

chair conformations, as indicated by the presence of peaks for the same proton type, but appeared with different proton integrations, similar to the trend of C-4-chlorophenylcalix[4]-2-methylresorcinarene. The crown conformation shows a singlet signal with one proton integral because all methine protons are symmetrical with each other. In contrast, the chair will show two proton peaks of the same type with the proton integration value half of the crown proton integral. The observed crown and chair conformations in this study have similarities with the results of research by Castillo-Aguirre et al. [26]. On the other hand, C-4-methoxyphenylcalix[4]-2-methylresorcinarene was found in a single crown conformation, as confirmed by a singlet signal of C-H methine at a chemical shift value of 5.37 ppm. This prediction was also confirmed by the appearance of two peaks in the LC chromatogram of both C-4-chlorophenylcalix[4]-2-methylresorcinarene and C-4-dimethylaminophenylcalix[4]-2-

methylresorcinarene compounds. On the other hand, the single conformation of C-4-methoxyphenylcalix[4]-2-methylresorcinarene, i.e., in a crown conformation, agreed with a single peak in the LC chromatogram of C-4-methoxyphenylcalix[4]-2-methylresorcinarene.

The other proton signals of each calix[4]-2-methylresorcinarene were also observed in their $^1\text{H-NMR}$ spectra. The methyl protons of 2-methylresorcinol were observed as a singlet signal at a chemical shift range of 1.95–2.06 ppm. The aromatic protons of 2-methylresorcinol moiety were also detected as a singlet signal at a chemical shift range of 5.58–6.23 ppm. Meanwhile, the aromatic protons of the benzaldehyde structure were found as two doublet signals with a J value of 8.5–9.0 Hz at a chemical shift range of 6.51–7.37 ppm. The OH protons of calix[4]-2-methylresorcinarene were observed as a singlet signal at a chemical shift range of 4.54 to 8.17 ppm. It is well-known that hydroxyl protons can be found in a wide range of chemical shifts in the $^1\text{H-NMR}$ spectra. The methoxy protons of C-4-methoxyphenylcalix[4]-2-methylresorcinarene were observed as a singlet signal at 3.77 ppm, while the dimethylamino protons of C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene were detected as singlet signals at a chemical shift range of 3.09–3.16 ppm.

Another characterization that proves the formation of calix[4]-2-methylresorcinarene compounds is done through LC-MS analysis. Characterization of C-4-chlorophenylcalix[4]-2-methylresorcinarene compound showed two signals with the same m/z value of 1005 for $[\text{M}+\text{H}+\text{K}]^+$ at the retention time of 12.81 and 13.14 min. Both peaks had the same m/z value, indicating that two conformations existed as suggested by the $^1\text{H-NMR}$ analysis, i.e., boat and chair conformations. The C-4-methoxyphenylcalix[4]-2-methylresorcinarene compound showed an m/z value of 986 $[\text{M}+\text{NH}_4]^+$ as a single peak at the retention time of 10.81 min, confirming that C-4-methoxyphenylcalix[4]-2-methylresorcinarene existed in one conformation only (crown conformation). Meanwhile, the C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene compound was observed as two signals with the same m/z value of 1021 $[\text{M}]^+$ at 8.55 and

9.35 min, confirming the presence of crown and chair conformations as revealed by the $^1\text{H-NMR}$ analysis. Based on the FTIR, $^1\text{H-NMR}$, and LC-MS data, it can be concluded that C-4-chlorophenylcalix[4]-2-methylresorcinarene, C-4-methoxyphenylcalix[4]-2-methylresorcinarene, and C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene have been successfully synthesized in the present work.

Antimalarial Activity of Calix[4]-2-methylresorcinarenes

The antimalarial activity of the synthesized calix[4]-2-methylresorcinarene derivatives was examined through an *in vitro* assay against chloroquine-resistant *P. falciparum* FCR-3. This resistant strain (FCR-3) was selected due to serious concern about malarial resistance against chloroquine diphosphate as a commercial drug during the last decade. This *in vitro* assay observes the schizont growth in the red blood cells in RPMI-1640 media. Table 1 shows the inhibition percentage of *P. falciparum* FCR-3 in the presence of various calix[4]-2-methylresorcinarene concentrations. Increasing the calix[4]-2-methylresorcinarene concentrations generally increased the inhibition percentage. At the highest concentration (12.5 $\mu\text{g/mL}$), C-4-chlorophenylcalix[4]-2-methylresorcinarene gave a higher inhibition percentage (82.17%) than C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene (53.77%) and much higher than C-4-methoxyphenylcalix[4]-2-methylresorcinarene (41.12%). A similar trend was also observed for the lowest concentration (0.5 $\mu\text{g/mL}$), whereas the inhibition percentage of C-4-chlorophenylcalix[4]-2-methylresorcinarene (22.72%) > C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene (17.87%) > C-4-methoxyphenylcalix[4]-2-methylresorcinarene (14.31%). This result shows that C-4-chlorophenylcalix[4]-2-methylresorcinarene was the best antimalarial agent against *P. falciparum* FCR-3 parasites among the synthesized calix[4]-2-methylresorcinarenes in this work.

The antimalarial activity is classified based on the IC_{50} values by Batista et al. [36]. These classifications are

Table 1. *Plasmodium falciparum* FCR-3 parasites inhibition percentages in a various concentration of calix[4]-2-methylresorcinarenes

Compound	Concentration (µg/mL)	Inhibition percentage (%)
C-4-chlorophenylcalix[4]-2-methylresorcinarene	12.5	82.17
	10.0	72.55
	5.0	57.99
	1.0	31.33
	0.5	22.72
C-4-methoxyphenylcalix[4]-2-methylresorcinarene	12.5	43.56
	10.0	43.11
	5.0	35.42
	1.0	22.94
	0.5	14.31
C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene	12.5	53.77
	10.0	41.12
	5.0	35.62
	1.0	23.98
	0.5	17.87

Table 2. The IC₅₀ value of antimalarial compounds based on calix-structure

Compounds	IC ₅₀ value (µM)
C-4-chlorophenylcalix[4]-2-methylresorcinarene	2.66
C-4-methoxyphenylcalix[4]-2-methylresorcinarene	23.63
C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene	13.82
C-4-nitrophenylcalix[4]-2-methylresorcinarene [21]	1.79
C-8-hydroxyquinolinecalix[4]arene [39]	0.07
C-2-aminopyrimidinecalix[4]arene [39]	0.04
Chloroquine	0.62
1,6-dihydroxyxanthone [24]	81.77
1,4,5-trihydroxyxanthone [40]	14.54
1,3,5-trihydroxyxanthone [40]	102.33

very active (IC₅₀ value < 1 µM), active (IC₅₀ value = 1–20 µM), moderate (IC₅₀ value = 20–100 µM), and inactive (IC₅₀ value > 100 µM) antimalarial agents. Table 2 shows that C-4-chlorophenylcalix[4]-2-methylresorcinarene, C-4-methoxyphenylcalix[4]-2-methylresorcinarene, and C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene compounds gave IC₅₀ values of 2.66, 23.63, and 13.82 µM, respectively, against *P. falciparum*. The C-4-chlorophenylcalix[4]-2-methylresorcinarene compound was found as an active antimalarial agent with the lowest IC₅₀ value of 2.88 µM, demonstrating the strongest antimalarial activity. The C-4-methoxyphenylcalix[4]-2-

methylresorcinarene compound was found as a moderate antimalarial agent with an IC₅₀ value of 23.63 µM while the C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene compound was an active antimalarial agent with an IC₅₀ value of 13.82 µM. These *in vitro* results showed that the antimalarial activities of calix[4]-2-methylresorcinarene derivatives were influenced by the various functional groups at the para position of the aromatic rings. The chloro functional group exhibited a higher antimalarial activity than the dimethylamino and methoxy functional groups. Both polar functional groups, i.e., chloro and dimethylamino,

exhibited higher antimalarial activity than the non-polar methoxy group. This phenomenon was in agreement with other reported publications [19-21,28,37]. More polar calix derivatives, *i.e.*, C-4-nitrophenylcalix[4]-2-methylresorcinarene (IC₅₀ value = 1.79 μM), C-8-hydroxyquinolinecalix[4]arene (IC₅₀ value = 0.07 μM), and C-2-aminopyrimidinecalix[4]arene (IC₅₀ value = 0.04 μM) yield much lower IC₅₀ values than C-4-chlorophenylcalix[4]-2-methylresorcinarene (IC₅₀ value = 2.66 μM). However, the antimalarial activity of calix[4]-2-methylresorcinarenes in this work was much stronger than hydroxyxanthone derivatives, *i.e.*, 1,6-dihydroxyxanthone (IC₅₀ value = 81.77 μM), 1,5,6-trihydroxyxanthone (IC₅₀ value = 27.64 μM), 1,6,8-trihydroxyxanthone (IC₅₀ value = 6.76 μM), 1,4,5-trihydroxyxanthone (IC₅₀ value = 14.54 μM), and 1,3,5-trihydroxyxanthone (IC₅₀ value = 102.33 μM), as a representative of polyphenolic compounds, which was remarkable. This meant that the cyclization reaction of the polyphenolic compound significantly enhanced the antimalarial activity. Even though the calix[4]-2-methylresorcinarenes exhibit lower activity than chloroquine diphosphate, the C-4-chlorocalix[4]-2-methylresorcinarenes may have great potential to overcome chloroquine resistance as they have a different geometry from chloroquine diphosphate [38]. Thus, C-4-chlorocalix[4]-2-methylresorcinarenes could be further functionalized to have a better and stronger antimalarial agent in the future.

■ CONCLUSION

Three calix[4]-2-methylresorcinarene have been successfully synthesized from 2-methylresorcinol through a one-pot reaction method under acidic conditions. Their chemical structures were confirmed through spectroscopic elucidation analyses. The C-4-chlorophenylcalix[4]-2-methylresorcinarene product was obtained in boat and chair conformations in 97% yield. The C-4-methoxyphenylcalix[4]-2-methylresorcinarene compound was produced in a single crown conformation in 95% yield. Meanwhile, the C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene compound was constructed in crown and chair conformations in 35% yield. The *in vitro*

antimalarial activity assay against *P. falciparum* FCR-3 showed that the C-4-chlorophenylcalix[4]-2-methylresorcinarene compound exhibited the most potent antimalarial activity with an IC₅₀ value of 2.66 μM, followed by C-4-dimethylaminophenylcalix[4]-2-methylresorcinarene compound (IC₅₀ value = 13.82 μM) and C-4-methoxyphenylcalix[4]-2-methylresorcinarene compound (IC₅₀ value = 23.63 μM). Therefore, the antimalarial activity of calix[4]-2-methylresorcinarenes depends on the type of functional group attached to the aromatic ring at the para position.

■ ACKNOWLEDGMENTS

The authors thank for the financial support from the National Research and Innovation Agency (BRIN) and Indonesia Endowment Fund for Education (LPDP) through *Riset dan Inovasi untuk Indonesia Maju* (RIIM) program for the budget year 2023-2024 with a contract number of 172/IV/KS/11/2023 and 6815/UN1/DITLIT/Dit-Lit/KP.01.03/2023.

■ CONFLICT OF INTEREST

The authors have no competing interests to be declared.

■ AUTHOR CONTRIBUTIONS

Jumina did conceptualization of this work. Jumina, Harno Dwi Pranowo, Eti Nurwening Sholikhah, Jeffry Julianus, and Susalit Setya Wibowo did supervision. Baiq Ike Nursofia conducted the experiment. Baiq Ike Nursofia, Krisfian Tata Aneka Priyanga, Yehezkiel Steven Kurniawan, Jeffry Julianus, Eti Nurwening Sholikhah, Harno Dwi Pranowo, and Hana Anisa Fatimi did formal analysis. Baiq Ike Nursofia, Krisfian Tata Aneka Priyanga, Yehezkiel Steven Kurniawan, and Yoga Priastomo wrote and revised the manuscript. All authors agreed to the final version of this manuscript.

■ REFERENCES

- [1] World Health Organization, 2023, *World Malaria Report 2021*, World Health Organization, Geneva, Switzerland, <https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2023/>, accessed on 2 February 2024.

- [2] Oluwafemi, T., and Azuaba, E., 2022, Impact of hygiene on malaria transmission dynamics: A mathematical model, *J. Multidiscip. Appl. Nat. Sci.*, 2 (1), 1–9.
- [3] Shibeshi, M.A., Kifle, Z.D., and Atnafie, S.A., 2020, Antimalarial drug resistance and novel targets for antimalarial drug discovery, *Infect. Drug Resist.*, 13, 4047–4060.
- [4] Balikagala, B., Fukuda, N., Ikeda, M., Katuru, O.T., Tachibana, S., Yamauchi, M., Opio, W., Emoto, S., Anywar, D.A., Kimura, E., Odongo-Aginya, E., Ogwang, M., Horii, T., and Mita, T., 2021, Evidence of artemisinin-resistant malaria in Africa, *N. Engl. J. Med.*, 385 (13), 1163–1171.
- [5] Chen, I., and Hsiang, M.S., 2022, Triple artemisinin-based combination therapies for malaria: A timely solution to counter antimalarial drug resistance, *Lancet Infect. Dis.*, 22 (6), 751–753.
- [6] Saunders, D.L., Vanachayangkul, P., and Lon, C., 2014, Dihydroartemisinin–piperazine failure in Cambodia, *N. Engl. J. Med.*, 371 (5), 484–485.
- [7] Pei, W.Y., Yang, J., Wu, H., Zhou, W., Yang, Y.W., and Ma, J.F., 2020, A calix[4]resorcinarene-based giant coordination cage: Controlled assembly and iodine uptake, *Chem. Commun.*, 56 (16), 2491–2494.
- [8] Ohto, K., 2021, Review of adsorbents incorporating calixarene derivatives used for metals recovery and hazardous ions removal: The concept of adsorbent design and classification of adsorbents, *J. Inclusion Phenom. Macrocyclic Chem.*, 101 (3), 175–194.
- [9] Basílio, N., Garcia-Rio, L., and Martín-Pastor, M., 2012, Calixarene-based surfactants: Evidence of structural reorganization upon micellization, *Langmuir*, 28 (5), 2404–2414.
- [10] Kurniawan, Y.S., Ryu, M., Sathuluri, R.R., Iwasaki, W., Morisada, S., Kawakita, H., Ohto, K., Maeki, M., Miyazaki, M., and Jumina, J., 2019, Separation of Pb(II) ion with tetraacetic acid derivative of calix[4]arene by using droplet-based microreactor system, *Indones. J. Chem.*, 19 (2), 368–375.
- [11] Memon, F.N., Memon, S., and Minhas, F.T., 2016, Calix[4]arene-mediated uphill transport of methyl red through bulk liquid membrane: Kinetics of operational variables, *Desalin. Water Treat.*, 57 (18), 8358–8371.
- [12] Priyanga, K.T.A., Kurniawan, Y.S., Ohto, K., and Jumina, J., 2022, A review on calixarene fluorescent chemosensor agents for various analytes, *J. Multidiscip. Appl. Nat. Sci.*, 2 (1), 23–40.
- [13] Naseer, M.M., Ahmed, M., and Hameed, S., 2017, Functionalized calix[4]arenes as potential therapeutic agents, *Chem. Biol. Drug Des.*, 89 (2), 243–256.
- [14] Du, D., Liu, Y., Lan, J., Hou, X., Liu, J., Shi, Q., Huang, Q., Xue, Y., Yan, C., and An, L., 2023, Novel biotin-linked amphiphilic calix[4]arene-based supramolecular micelles as doxorubicin carriers for boosted anticancer activity, *Chem. Commun.*, 59 (83), 12487–12490.
- [15] Ni, J., Lu, L., and Liu, Y., 2019, Antiradical and antioxidative activity of azocalix[4]arene derivatives: Combined experimental and theoretical study, *Molecules*, 24 (3), 485.
- [16] Zarranz, B., Jaso, A., Aldana, I., Monge, A., Maurel, S., Deharo, E., Jullian, V., and Sauvain, M., 2005, Synthesis and antimalarial activity of new 3-arylquinoxaline-2-carbonitrile derivatives, *Arzneimittelforschung*, 55 (12), 754–761.
- [17] Liu, M., Wilairat, P., Croft, S.L., Tan, A.L.C., and Go, M.L., 2003, Structure-activity relationships of antileishmanial and antimalarial chalcones, *Bioorg. Med. Chem.*, 11 (13), 2729–2738.
- [18] Syahri, J., Nasution, H., Nurohmah, B.A., Purwono, B., Yuanita, E., Zakaria, N.H., and Hassan, N.I., 2020, Design, synthesis and biological evaluation of aminoalkylated chalcones as antimalarial agent, *Sains Malays.*, 49 (11), 2667–2677.
- [19] Putri, R.R., Pranowo, H.D., Kurniawan, Y.S., Fatimi, H.A., and Jumina, J., 2023, Synthesis of calix[4]resorcinarene derivatives as antimalarial agents through heme polymerization inhibition assay, *Indones. J. Chem.*, 23 (4), 1032–1041.
- [20] Sari, D.K., Hidayat, D.N.W., Fatmawati, D.R., Triono, S., Kurniawan, Y.S., and Jumina, J., 2022, Synthesis and antimalarial activity assay of C-arylcalix[4]pyrogallolarenes using heme polymerization inhibition activity (HPIA) method,

- Mater. Sci. Forum*, 1061, 187–193.
- [21] Nisa, S.A., Jumina, J., Mardjan, M.I.D., and Kurniawan, Y.S., 2023, Synthesis, activity test and molecular docking of novel nitrophenylcalix[4]-2-methylresorcinarene derivatives as antimalarial agent, *Molekul*, 18 (3), 404–413.
- [22] Jumina, J., Siswanta, D., Zulkarnaian, A.K., Triono, S., Priatmoko, P., Yuanita, E., Imawan, A.C., Fatmasari, N., and Nursalim, I., 2019, Development of C-arylcalix[4]resorcinarenes and C-arylcalix[4]pyrogallolarenes as antioxidant and UV-B protector, *Indones. J. Chem.*, 19 (2), 273–284.
- [23] Congpuong, K., Sirtichaisinthop, J., Tippawangkosol, P., Suprakrob, K., Na-Bangchang, K., Tan-ariya, P., and Karbwang, J., 1998, Incidence of antimalarial pretreatment and drug sensitivity *in vitro* in multidrug-resistant *Plasmodium falciparum* infection in Thailand, *Trans. R. Soc. Trop. Med. Hyg.*, 92 (1), 84–86.
- [24] Zakiah, M., Syarif, R.A., Mustofa, M., Jumina, J., Fatmasari, N., and Sholikhah, E.N., 2021, *In vitro* antiplasmodial, heme polymerization, and cytotoxicity of hydroxyxanthone derivatives, *J. Trop. Med.*, 2021 (1), 8866681.
- [25] Rajkumar, P., Buvanewari, N., Vaheith, Z.A., Ahamed, A.F., Saraswathy, G., and Dayanandhan, R., 2021, Kinetic analysis of oxidation of α -hydroxy acids by Caro's acid in micellar medium, *Rasayan J. Chem.*, 14 (2), 785–793.
- [26] Castillo-Aguirre, A., Rivera-Monroy, Z., and Maldonado, M., 2017, Selective O-alkylation of the crown conformer of tetra(4-hydroxyphenyl)calix[4]resorcinarene to the corresponding tetraalkyl ether, *Molecules*, 22 (10), 1660.
- [27] Elidrisi, I., Bhatt, P.V., Govender, T., Kruger, H.G., and Maguire, G.E.M., 2015, Synthesis and NMR elucidation of novel octa-amino acid resorcin[4]arenes derivatives, *S. Afr. J. Chem.*, 68, 27–38.
- [28] Pineda-Castañeda, H., Maldonado-Villamil, M., Parra-Giraldo, C.M., Leal-Castro, A.L., Fierro-Medina, R., Rivera-Monroy, Z.J., and García-Castañeda, J.E., 2023, Peptide-resorcinarene conjugates obtained via click chemistry: Synthesis and antimicrobial activity, *Antibiotics*, 12 (4), 773.
- [29] Galindres, D.M., Cifuentes, D., Tinoco, L.E., Murillo-Acevedo, Y., Rodrigo, M.M., Ribeiro, A.C.F., and Estes, M.A., 2022, A review of the application of resorcinarenes and SBA-15 in drug delivery, *Processes*, 10 (4), 684.
- [30] Pineda-Castañeda, H.M., Maldonado, M., and Rivera-Monroy, Z.J., 2023, Efficient separation of C-tetramethylcalix[4]resorcinarene conformers by means of reversed-phase solid-phase extraction, *ACS Omega*, 8 (1), 231–237.
- [31] Shebitha, A.M., Sreejith, S.S., Sherly Mole, P.B., Mohan, N., Avudaiappan, G., Hiba, K., Priya, K.S., and Sreekumar, K., 2020, Facile synthesis, X-ray diffraction studies, photophysical properties and DFT-D based conformational analysis of octa and dodecacyanomethoxycalix[4]resorcinarenes, *J. Mol. Struct.*, 1214, 128215.
- [32] Taylor, D., Ling, I., Vilela, F., and Dalgarno, S.J., 2022, Intermolecular interactions drive the unusual co-crystallization of different calix[4]arene conformations, *Crystals*, 12 (2), 250.
- [33] Liu, J.L., Zhang, P.Z., Jia, A.Q., Shi, H.T., and Zhang, Q.F., 2022, Supramolecular assemblies of sulfonatomethylated calix[4]resorcinarenes with aquated sodium(I), cesium(I), and aluminum(III) ions, *ChemistrySelect*, 7 (1), e202104118.
- [34] Liu, J.L., Liu, X.L., Jia, A.Q., Shi, H.T., and Zhang, Q.F., 2020, Supramolecular structures and crystal stability of diisobutylaminomethylated calix[4]resorcinarenes, *J. Inclusion Phenom. Macrocyclic Chem.*, 98 (1), 49–56.
- [35] Eddaif, L., Trif, L., Telegdi, J., Egyed, O., and Shaban, A., 2019, Calix[4]resorcinarene macrocycles, *J. Therm. Anal. Calorim.*, 137 (2), 529–541.
- [36] Batista, R., De Jesus Silva Júnior, A., and De Oliveira, A.B., 2009, Plant-derived antimalarial agents: New leads and efficient phytomedicines. Part II. Non-alkaloidal natural products, *Molecules*, 14 (8), 3037–3072.
- [37] Park, G.M., Park, H., Oh, S., and Lee, S., 2017, Antimalarial activity of C-10 substituted triazolyl artemisinin, *Korean J. Parasitol.*, 55 (6), 661–665.

- [38] Wicht, K.J., Mok, S., and Fidock, D.A., 2020, Molecular mechanisms of drug resistance in *Plasmodium falciparum* malaria, *Annu. Rev. Microbiol.*, 74, 431–454.
- [39] Shah, R.B., Valand, N.N., Sutariya, P.G., and Menon, S.K., 2016, Design, synthesis and characterization of quinoline-pyrimidine linked calix[4]arene scaffolds as anti-malarial agents, *J. Inclusion Phenom. Macrocyclic Chem.*, 84 (1), 173–178.
- [40] Kpotin, G.A., Bédé, A.L., Houngue-Kpota, A., Anatovi, W., Kuevi, U.A., Atohou, G.S., Mensah, J.B., Gómez-Jeria, J.S., and Badawi, M., 2019, Relationship between electronic structures and antiplasmodial activities of xanthone derivatives: A 2D-QSAR approach, *Struct. Chem.*, 30 (6), 2301–2310.