

SERTIFIKAT

Direktorat Jendral Pendidikan Tinggi, Riset dan Teknologi
Kementerian Pendidikan, Kebudayaan, Riset dan Teknologi Republik Indonesia



Kutipan dari Keputusan Direktorat Jendral Pendidikan Tinggi, Riset, dan Teknologi
Kementerian Pendidikan, Kebudayaan, Riset dan Teknologi Republik Indonesia

Nomor: 79/E/KPT/2023

Peringkat Akreditasi Jurnal Ilmiah Periode I Tahun 2023

Nama Jurnal Ilmiah:

Komputasi: Jurnal Ilmiah Ilmu Komputer dan Matematika

E-ISSN: 26543990

Universitas Pakuan

Ditetapkan Sebagai Jurnal Ilmiah:

TERAKREDITASI PERINGKAT 4

Akreditasi Berlaku selama 5 (lima) Tahun, yaitu:
volume 19 Nomor 2 Tahun 2022 Sampai Volume 24 Nomor 1 Tahun 2027

Jakarta, 11 May 2023

Plt. Direktur Jendral Pendidikan Tinggi, Riset, dan Teknologi



Prof. Ir. Nizam, M.Sc., DIC, Ph.D., IPU, ASEAN Eng
NIP. 196107061987101001

TERAKREDITASI





Numerical Verification of the Order of Accuracy of a Runge-Kutta-Fehlberg Method in Solving an SEIR Model

Lintang Kinanthi Widi

Sanata Dharma University

Sudi Mungkasi

Sanata Dharma University

ABSTRACT

A Runge-Kutta-Fehlberg method (RKF45) is considered to solve an SEIR (Susceptible-Exposed-Infected-Recovered) mathematical model for infectious disease transmission. This numerical method is of the Runge-Kutta class, and has theoretical order of accuracy four and five depending on the involved formulations. We limit our problem to the RKF45 formula having theoretical order five. It is important to understand the behavior of RKF45 in solving an initial value problem, especially when implemented for real problems. In this paper, we take an SEIR mathematical model in our case, because this model is applicable in the prediction of infectious disease transmission in real life. Our research method is quantitative. We record the errors of numerical simulations. From the error values, we determine the numerical order of accuracy. Our research results show that RKF45 produces numerical order of accuracy five for the time step is sufficiently small. Therefore, the numerical order of accuracy matches with the theoretical one.

Vol. 22, No. 2 July 2025

KOMPUTASI

Jurnal Ilmiah Ilmu Komputer dan Matematika

IOT-based Soil Moisture
Monitoring and Water
Pump Control System for
Ornamental Plants

Numerical Verification of the
Order of Accuracy of a Runge-
Kutta-Fehlberg Method in Solving
an SEIR Model

Shallot Growth Stage Monitoring
with Multispectral Imagery using
MobileNetV2

Cyberscrime Awareness Phishing
Email Using Secure Socket Layer
as an Interpretation of Information
Security

The similarity of COVID-19-
Related Profiles and Pandemic
Conditions in ASEAN Countries

Implementation of Fuzzy Search
and Audio Player for Web-Based
Quranic Verse Retrieval

Improvement Market Basket n
Transaction Sales Using Apriori
Algorithm

Application of Proportional
Integral Derivative System
as Speed Control on Mobile
Robot Line Follower

Recent Advances in
Deep Learning for MRI-Based
Brain Tumor Identification:
A Systematic Review (2020-2025)

Application of Principal
Componen Analysis on Factors
Causing Inflation in West

Interpreting Sign
Language Images into
Text-Based using YOLOv7

Smart Waste Management
System Using Computer Vision

KOMPUTASI

Vol.22

No.2

Hal 1 - 107

P-ISSN 1693 - 7554

E-ISSN 2654 - 3990

 PDF

PUBLISHED

31-07-2025

HOW TO CITE

Widi, L. K., & Mungkasi, S. (2025). Numerical Verification of the Order of Accuracy of a Runge-Kutta-Fehlberg Method in Solving an SEIR Model. *Komputasi: Jurnal Ilmiah Ilmu Komputer Dan Matematika*, 22(2), 10–19. Retrieved from <https://komputasi-fmipa.unpak.ac.id/index.php/komputasi/article/view/30>

More Citation Formats ▼

ISSUE

[Vol. 22 No. 2 \(2025\): Komputasi: Jurnal Ilmiah Ilmu Komputer dan Matematika](#)

SECTION

Articles

LICENSE

Copyright (c) 2025 Komputasi: Jurnal Ilmiah Ilmu Komputer dan Matematika



This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License](#).

ABOUT JOURNAL

Editorial Team

Peer-Reviewers

Peer-Reviewers Process

ACCREDITATION STATUS



TEMPLATE



INDEX BY

Google Scholar



SCREENED BY



ROAD



ISSN

E-ISSN :1538725485

P-ISSN : 1693-7554

TOOLS



MENDELEY



grammarly

VISITOR



[View My Stats](#)

MAKE A SUBMISSION

LANGUAGE

[English](#)

[Bahasa Indonesia](#)

INFORMATION

[For Readers](#)

[For Authors](#)

[For Librarians](#)

Komputasi: Jurnal Ilmiah Ilmu Komputer dan Matematika (P-ISSN: [1693-7554](#) | e-ISSN: [2654-3990](#) (Online))
by Department of Computer Science, Faculty of Natural Science, Universitas Pakuan
Powered by [Public Knowledge Project OJS](#).

Komputasi: Jurnal Ilmiah Ilmu Komputer dan Matematika
Jalan Raya Pakuan PO. BOX 452, Bogor, Indonesia
Telp: +62251-8363419, Fax: +62251-8356927
Email: komputasi@unpak.ac.id



[HOME](#) / [Editorial Team](#)

Editorial Team

EDITOR IN CHIEF

- [Asep Denih](#), (Scopus ID:57220039564) Department of Computer Science, Pakuan University, Indonesia

EDITORIAL BOARD

- [Herry Pribawanto Suryawan](#), (Scopus ID: 56201983000) Faculty of Science and Technology, Universitas Sanata Dharma, Yogyakarta, Indonesia
- [Eneng Tita Tosida](#), (Scopus ID: 57189053683) Department of Computer Science, Pakuan University, Indonesia
- [Prihastuti Harsani](#), (Scopus ID: 57201548846) Department of Computer Science, Pakuan University, Indonesia
- [Asep Saepulrohman](#), (Scopus ID: 57192938246) Department of Computer Science, Universitas Pakuan, Indonesia, Indonesia
- [Lita Karlitasari](#), (Scopus ID: 57193497111) Department of Computer Science, Pakuan University, Indonesia, Indonesia
- [Rahmat Hidayat](#), (Scopus ID: 57205744812) Department of Mathematics, Universitas Cokroaminoto Palopo, Indonesia
- [Andi Chairunnas](#), (Scopus ID: 57201556112) Department of Computer Science, Pakuan University, Indonesia
- [Teguh Puja Negara](#), (Scopus ID: 36004810100) Department of Computer Science, Pakuan University, Indonesia
- [Dinar Munggaran Akhmad](#), (Scopus ID: 57202585094) Department of Computer Science, Pakuan University, Indonesia, Indonesia

ABOUT JOURNAL

[Editorial Team](#)

[Peer-Reviewers](#)

[Peer-Reviewers Process](#)

[Focus & Scope](#)

[Author Guidelines](#)



Focus and Scope

Komputasi: Jurnal Ilmiah Ilmu Komputer dan Matematika is initiated and organized by Department of Computer Science, Faculty of Mathematics and Science, Pakuan University (Unpak), Bogor, Indonesia to accommodate the writing of research results for the academics and institutions other. Komputasi journal was originally launched in 1992, and published online since 2007 with ISSN version [p-ISSN: 1693-7554](#) and version of the daring of [e-ISSN: 2654-3990](#) in 2018 (SK No. 0005.26543990/JI.3.1/SK.ISSN/2018.10-15 October 2018 (starting Vol. 16, No. 1, January 2019). The journal is a publication media for original manuscripts related information technology development and science written in Bahasa Indonesia which is published twice times a year (January and July).

Manuscripts accepted for publication in KOMPUTASI may be results of the latest scientific research (maximum during the past 10 years). The scope of the articles published in this journal deal with a broad range of computer and mathematics topics, in general, this journal consists of 4 main themes, including:

1. Software engineering
 - Information systems
 - Software testing
2. Hardware engineering
 - Network
 - Sensor network
 - Robotics, IoT
3. Computing Intelligence
 - Data minning
 - Big data
 - Data intelligence
 - Numerical analysis
 - Bioinformatics
 - Operations research
4. Mathematics
 - Discrete Mathematics and Combinatorics
 - Modelling and Simulation
 - Applied Mathematics
 - Computational Mathematics

ABOUT JOURNAL

Editorial Team

Peer-Reviewers

Peer-Reviewers Process

Focus & Scope

Author Guidelines

[HOME](#) / [ARCHIVES](#) / Vol. 22 No. 2 (2025): Komputasi: Jurnal Ilmiah Ilmu Komputer dan Matematika

Vol. 22 No. 2 (2025): Komputasi: Jurnal Ilmiah Ilmu Komputer dan Matematika



DOI: <https://doi.org/10.33751/komputasi.v22i2>

PUBLISHED: 31-07-2025

ARTICLES



IOT-based Soil Moisture Monitoring and Water Pump Control System for Ornamental Plants

Sistem Monitoring Kelembapan Tanah Dan Kendali Pompa Air Tanaman Hias Berbasis Iot

Attalarik Zidan, Yunita Rahma , Sufiatul Maryana

1-9





Numerical Verification of the Order of Accuracy of a Runge-Kutta-Fehlberg Method in Solving an SEIR Model

Lintang Kinanthi Widi, Sudi Mungkasi

10-19



PDF



Shallot Growth Stage Monitoring with Multispectral Imagery using MobileNetV2

Yanuar Firmansyah, Nugraha Priya Utama

20-27



PDF



Cybercrime Awareness Cybercrime Awareness Phishing Email Using Secure Socket Layer as an Interpretation of Information Security

Cybercrime Awareness Phishing Email Using Secure Socket Layer as an Interpretation of Information Security

Kotim Subandi, Dinar Munggaran A, Victor Ilyas Sugara, Adriana Sari Aryani, Hermawan T

28-34



PDF



COVID-19-Related The Similarity of COVID-19-Related Profiles and Pandemic Conditions in ASEAN Countries

The Similarity of COVID-19-Related Profiles and Pandemic Conditions in ASEAN Countries

Elis Khatizah

35-44



PDF



Implementation of Fuzzy Search and Audio Player for Web-Based Quranic Verse Retrieval

Dewi Primasari, Firman Hakim, Muhammad Danise Raditya Saneistha, Ihsan Wahid Cahya

45-52



PDF



IMPROVEMENT MARKET BASKET ON TRANSACTION SALES USING APRIORI ALGORITHM (Case study : Workshop The Sari (Mulya)

Nurul Aziz, Sri Setyaningsih, Dini Suhartini

53-63



PDF



Application of Proportional Integral Derivative system as speed control on Mobile Robot Line Follower

submission of Journal Thesis

Raka Pramudiya Ramadani, Arie Qur'ania, Agus Ismangil

64-74



PDF



Recent Advances in Deep Learning for MRI-Based Brain Tumor Identification: A Systematic Review (2020 - 2025)

adly maulana, Suherman

75-82



PDF



Application of Principal Component Analysis on Factors Causing Inflation in West Kalimantan

Asri Rahmawati, Yuyun Eka Pratiwi, Onelia Rochmah

83-89



PDF



Interpreting sign language images into Text-Based using YOLOv7

Julius Sabilala, Rv Jan Zaldo Bautista, Lea Mae Gambol Tortogo, Lerry Joy Ga Juntarciego, Charles Vincent Gayoma, Adrian Forca

90-98



PDF



Smart Waste Management System Using Computer Vision

Mj Gagnao, Mark Leonel De Isidro, Ronelo Galaura , Jyrillen Segobre, Jamaica Mae Gamala, Reiner Jun Alminaza

99-107



PDF

ABOUT JOURNAL

Editorial Team

Peer-Reviewers

Peer-Reviewers Process

Focus & Scope

Author Guidelines

ACCREDITATION STATUS



TEMPLATE



INDEX BY

Google Scholar



SCREENED BY



ROAD



ISSN

E-ISSN :1538725485

P-ISSN : 1693-7554

TOOLS



MENDELEY



grammarly

VISITOR



[View My Stats](#)

[MAKE A SUBMISSION](#)

LANGUAGE

[English](#)

Bahasa Indonesia

INFORMATION

[For Readers](#)

[For Authors](#)

[For Librarians](#)

Komputasi: Jurnal Ilmiah Ilmu Komputer dan Matematika (P-ISSN: [1693-7554](#) | e-ISSN: [2654-3990](#) (Online))

by Department of Computer Science, Faculty of Natural Science, Universitas Pakuan

Powered by [Public Knowledge Project OJS](#).

Komputasi: Jurnal Ilmiah Ilmu Komputer dan Matematika

Jalan Raya Pakuan PO. BOX 452, Bogor, Indonesia

Telp: +62251-8363419, Fax: +62251-8356927

Email: komputasi@unpak.ac.id

Numerical Verification of the Order of Accuracy of a Runge-Kutta-Fehlberg Method in Solving an SEIR Model

Lintang Kinanthi Widi^{1*}, Sudi Mungkasi²

^{1,2} Department of Mathematics, Faculty of Science and Technology, Sanata Dharma University, Yogyakarta, 55281, Indonesia

*Corresponding author. E-mail address: lintangkinanthiw@gmail.com

Abstract

A Runge-Kutta-Fehlberg method (RKF45) is considered to solve an SEIR (Susceptible-Exposed-Infected-Recovered) mathematical model for infectious disease transmission. This numerical method is of the Runge-Kutta class, and has theoretical order of accuracy four and five depending on the involved formulations. We limit our problem to the RKF45 formula having theoretical order five. It is important to understand the behavior of RKF45 in solving an initial value problem, especially when implemented for real problems. In this paper, we take an SEIR mathematical model in our case, because this model is applicable in the prediction of infectious disease transmission in real life. Our research method is quantitative. We record the errors of numerical simulations. From the error values, we determine the numerical order of accuracy. Our research results show that RKF45 produces numerical order of accuracy five for the time step is sufficiently small. Therefore, the numerical order of accuracy matches with the theoretical one.

Keywords: Numerical method; order of accuracy; Runge-Kutta-Fehlberg method; SEIR model; verification

1. Introduction

Numerical methods have been important tools for solving various mathematical problems [1]. A mathematical problem itself is usually occurred from a real-world problem. This means that numerical methods are useful for solving real-world problems [2].

In this paper we consider a Runge-Kutta-Fehlberg method (RKF45) [3], which is applicable for solving an initial value problem of ordinary differential equations, such as an SEIR (Susceptible-Exposed-Infected-Recovered) mathematical model [4]. RKF45 method and SEIR model are of our interest, because RKF45 is simple to program (as it is a one-step method) and it has the order of accuracy four or five, and SEIR model is useful in modelling the real-world infectious disease transmission. In this paper, we limit our RKF45 method to be the one having the fifth order of accuracy.

The theoretical order of accuracy of a numerical method needs to be verified numerically. This is important, because modelers must know the behavior of the numerical method when implemented. Understanding the numerical behavior of the method is useful, so that the analysis of numerical results will be more accurate [5]-[7]. This will aid in understanding the solutions of the real-world problem through mathematical modeling and its solutions [8]-[13]. Based on these grounds, this paper aims to verify numerically the fifth order RKF45 method in solving a SEIR model.

The rest of this paper consists of Section 2 recalling the mathematical model, Section 3 writing the numerical method, Section 4 containing results and discussion, and Section 5 remarking some conclusions.

2. Mathematical Model

The SEIR model for infectious disease transmission consists of four compartments of the population, namely, S (Susceptible) subpopulation, E (Exposed) subpopulation, I (Infected) subpopulation, and R (Recovered) subpopulation. In this paper we consider the improved SEIR model proposed by Jiao et al. [4]. The schematic diagram of the model is illustrated in Figure 1.

*Corresponding author. E-mail address: lintangkinanthiw@gmail.com




Figure 1. Schematic diagram for the SEIR model.

The SEIR model is expressed as a system of ordinary differential equation as follows [4]:

$$\begin{cases} \frac{dS}{dt} = \Lambda - \beta(1 - \theta_1)S(t)[I(t) + \theta_2 E(t)] - \mu S(t), \\ \frac{dE}{dt} = \beta(1 - \theta_1)S(t)[I(t) + \theta_2 E(t)] - (\delta + \mu)E(t), \\ \frac{dI}{dt} = \delta E(t) - (\gamma + \sigma + \mu)I(t), \\ \frac{dR}{dt} = (\gamma + \theta_3 \sigma)I - \mu R(t). \end{cases} \quad (1)$$

In this model, t is the time variable and $\Lambda, \beta, \theta_1, \theta_2, \theta_3, \mu, \delta, \gamma, \sigma$ are parameters. In addition, S, E, I, R are dependent on t . More explanation and properties of this model are available in the work of Jiao et al. [4].

3. Methods

The numerical method for solving the problem is the fifth-order RKF45. Suppose we are given the initial value problem:

$$\begin{aligned} \frac{dy}{dt} &= f(t, y) \\ y(t_0) &= y_0 \end{aligned}$$

then the fifth-order RKF45 method takes the following scheme [3]:

$$y_{i+1} = y_i + \frac{16}{135}k_1 + \frac{6656}{12825}k_3 + \frac{28561}{56430}k_4 - \frac{9}{50}k_5 + \frac{2}{55}k_6 \quad (2)$$

where:

$$\begin{aligned} k_1 &= h f(t_i, y_i), \\ k_2 &= h f\left(t_i + \frac{1}{4}h, y_i + \frac{1}{4}k_1\right), \\ k_3 &= h f\left(t_i + \frac{3}{8}h, y_i + \frac{3}{32}k_1 + \frac{9}{32}k_2\right), \\ k_4 &= h f\left(t_i + \frac{12}{13}h, y_i + \frac{1932}{2197}k_1 - \frac{7200}{2197}k_2 + \frac{7296}{2197}k_3\right), \\ k_5 &= h f\left(t_i + h, y_i + \frac{439}{216}k_1 - 8k_2 + \frac{3680}{513}k_3 - \frac{845}{4104}k_4\right), \\ k_6 &= h f\left(t_i + \frac{1}{2}h, y_i - \frac{8}{27}k_1 + 2k_2 - \frac{3544}{2565}k_3 + \frac{1859}{4104}k_4 - \frac{11}{40}k_5\right). \end{aligned} \quad (3)$$

Therefore, the RKF45 scheme for solving the model is:

$$\begin{aligned} S_{i+1} &= S_i + \left(\frac{16}{135}k_{1,S} + \frac{6656}{12825}k_{3,S} + \frac{28561}{56430}k_{4,S} - \frac{9}{50}k_{5,S} + \frac{2}{55}k_{6,S}\right) \\ E_{i+1} &= E_i + \left(\frac{16}{135}k_{1,E} + \frac{6656}{12825}k_{3,E} + \frac{28561}{56430}k_{4,E} - \frac{9}{50}k_{5,E} + \frac{2}{55}k_{6,E}\right) \\ I_{i+1} &= I_i + \left(\frac{16}{135}k_{1,I} + \frac{6656}{12825}k_{3,I} + \frac{28561}{56430}k_{4,I} - \frac{9}{50}k_{5,I} + \frac{2}{55}k_{6,I}\right) \\ R_{i+1} &= R_i + \left(\frac{16}{135}k_{1,R} + \frac{6656}{12825}k_{3,R} + \frac{28561}{56430}k_{4,R} - \frac{9}{50}k_{5,R} + \frac{2}{55}k_{6,R}\right) \end{aligned} \quad (4)$$

with the $k_1, k_2, k_3, k_4, k_5, k_6$ values are as follows:

1. The k_1 values are given by:

$$\begin{aligned} k_{1,S} &= h * \{\Lambda - \beta(1 - \theta_1)S_i[I_i + \theta_2 E_i] - \mu S_i\} \\ k_{1,E} &= h * \{\beta(1 - \theta_1)S_i[I_i + \theta_2 E_i] - (\delta + \mu)E_i\} \\ k_{1,I} &= h * \{\delta E_i - (\gamma + \sigma + \mu)I_i\} \\ k_{1,R} &= h * \{(\gamma + \theta_3 \sigma)I_i - \mu R_i\} \end{aligned} \quad (5)$$

2. The k_2 values are given by:

$$\begin{aligned} k_{2,S} &= h * \left\{ \Lambda - \beta(1 - \theta_1) \left(S_i + \frac{1}{4} k_{1,S} \right) \left[\left(I_i + \frac{1}{4} k_{1,I} \right) + \theta_2 \left(E_i + \frac{1}{4} k_{1,E} \right) \right] \right. \\ &\quad \left. - \mu \left(S_i + \frac{1}{4} k_{1,S} \right) \right\} \\ k_{2,E} &= h * \left\{ \beta(1 - \theta_1) \left(S_i + \frac{1}{4} k_{1,S} \right) \left[\left(I_i + \frac{1}{4} k_{1,I} \right) + \theta_2 \left(E_i + \frac{1}{4} k_{1,E} \right) \right] \right. \\ &\quad \left. - (\delta + \mu) \left(E_i + \frac{1}{4} k_{1,E} \right) \right\} \\ k_{2,I} &= h * \left\{ \delta \left(E_i + \frac{1}{4} k_{1,E} \right) - (\gamma + \sigma + \mu) \left(I_i + \frac{1}{4} k_{1,I} \right) \right\} \\ k_{2,R} &= h * \left\{ (\gamma + \theta_3 \sigma) \left(I_i + \frac{1}{4} k_{1,I} \right) - \mu \left(R_i + \frac{1}{4} k_{1,R} \right) \right\} \end{aligned} \quad (6)$$

3. The k_3 values are given by:

$$\begin{aligned} k_{3,S} &= h * \left\{ \Lambda - \beta(1 - \theta_1) \left(S_i + \frac{3}{32} k_{1,S} + \frac{9}{32} k_{2,S} \right) \left[\left(I_i + \frac{3}{32} k_{1,I} + \frac{9}{32} k_{2,I} \right) \right. \right. \\ &\quad \left. \left. + \theta_2 \left(E_i + \frac{3}{32} k_{1,E} + \frac{9}{32} k_{2,E} \right) \right] - \mu \left(S_i + \frac{3}{32} k_{1,S} + \frac{9}{32} k_{2,S} \right) \right\} \\ k_{3,E} &= h * \left\{ \beta(1 - \theta_1) \left(S_i + \frac{3}{32} k_{1,S} + \frac{9}{32} k_{2,S} \right) \left[\left(I_i + \frac{3}{32} k_{1,I} + \frac{9}{32} k_{2,I} \right) \right. \right. \\ &\quad \left. \left. + \theta_2 \left(E_i + \frac{3}{32} k_{1,E} + \frac{9}{32} k_{2,E} \right) \right] - (\delta + \mu) \left(E_i + \frac{3}{32} k_{1,E} + \frac{9}{32} k_{2,E} \right) \right\} \\ k_{3,I} &= h * \left\{ \delta \left(E_i + \frac{3}{32} k_{1,E} + \frac{9}{32} k_{2,E} \right) - (\gamma + \sigma + \mu) \left(I_i + \frac{3}{32} k_{1,I} + \frac{9}{32} k_{2,I} \right) \right\} \\ k_{3,R} &= h * \left\{ (\gamma + \theta_3 \sigma) \left(I_i + \frac{3}{32} k_{1,I} + \frac{9}{32} k_{2,I} \right) - \mu \left(R_i + \frac{3}{32} k_{1,R} + \frac{9}{32} k_{2,R} \right) \right\} \end{aligned} \quad (7)$$

4. The k_4 values are given by:

$$\begin{aligned} k_{4,S} &= h * \left\{ \Lambda - \beta(1 - \theta_1) \left(S_i + \frac{1932}{2197} k_{1,S} - \frac{7200}{2197} k_{2,S} \right. \right. \\ &\quad \left. \left. + \frac{7296}{2197} k_{3,S} \right) \left[\left(I_i + \frac{1932}{2197} k_{1,I} - \frac{7200}{2197} k_{2,I} + \frac{7296}{2197} k_{3,I} \right) \right. \right. \\ &\quad \left. \left. + \theta_2 \left(E_i + \frac{1932}{2197} k_{1,E} - \frac{7200}{2197} k_{2,E} + \frac{7296}{2197} k_{3,E} \right) \right] \right. \\ &\quad \left. - \mu \left(S_i + \frac{1932}{2197} k_{1,S} - \frac{7200}{2197} k_{2,S} + \frac{7296}{2197} k_{3,S} \right) \right\} \\ k_{4,E} &= h * \left\{ \beta(1 - \theta_1) \left(S_i + \frac{1932}{2197} k_{1,S} - \frac{7200}{2197} k_{2,S} \right. \right. \\ &\quad \left. \left. + \frac{7296}{2197} k_{3,S} \right) \left[\left(I_i + \frac{1932}{2197} k_{1,I} - \frac{7200}{2197} k_{2,I} + \frac{7296}{2197} k_{3,I} \right) \right. \right. \\ &\quad \left. \left. + \theta_2 \left(E_i + \frac{1932}{2197} k_{1,E} - \frac{7200}{2197} k_{2,E} + \frac{7296}{2197} k_{3,E} \right) \right] \right. \\ &\quad \left. - (\delta + \mu) \left(E_i + \frac{1932}{2197} k_{1,E} - \frac{7200}{2197} k_{2,E} + \frac{7296}{2197} k_{3,E} \right) \right\} \\ k_{4,I} &= h * \left\{ \delta \left(E_i + \frac{1932}{2197} k_{1,E} - \frac{7200}{2197} k_{2,E} + \frac{7296}{2197} k_{3,E} \right) \right. \\ &\quad \left. - (\gamma + \sigma + \mu) \left(I_i + \frac{1932}{2197} k_{1,I} - \frac{7200}{2197} k_{2,I} + \frac{7296}{2197} k_{3,I} \right) \right\} \\ k_{4,R} &= h * \left\{ (\gamma + \theta_3 \sigma) \left(I_i + \frac{1932}{2197} k_{1,I} - \frac{7200}{2197} k_{2,I} + \frac{7296}{2197} k_{3,I} \right) \right. \\ &\quad \left. - \mu \left(R_i + \frac{1932}{2197} k_{1,R} - \frac{7200}{2197} k_{2,R} + \frac{7296}{2197} k_{3,R} \right) \right\} \end{aligned} \quad (8)$$

5. The k_5 values are given by:

$$\begin{aligned}
 k_{5,S} &= h * \left\{ \Lambda - \beta(1 - \theta_1) \left(S_i + \frac{439}{216} k_{1,S} - 8k_{2,S} + \frac{3680}{513} k_{3,S} - \frac{845}{4104} k_{4,S} \right) \right. \\
 &\quad \left[\left(I_i + \frac{439}{216} k_{1,I} - 8k_{2,I} + \frac{3680}{513} k_{3,I} - \frac{845}{4104} k_{4,I} \right) \right. \\
 &\quad \left. + \theta_2 \left(E_i + \frac{439}{216} k_{1,E} - 8k_{2,E} + \frac{3680}{513} k_{3,E} - \frac{845}{4104} k_{4,E} \right) \right] \\
 &\quad \left. - \mu \left(S_i + \frac{439}{216} k_{1,S} - 8k_{2,S} + \frac{3680}{513} k_{3,S} - \frac{845}{4104} k_{4,S} \right) \right\} \\
 k_{5,E} &= h * \left\{ \beta(1 - \theta_1) \left(S_i + \frac{439}{216} k_{1,S} - 8k_{2,S} + \frac{3680}{513} k_{3,S} - \frac{845}{4104} k_{4,S} \right) \right. \\
 &\quad \left[\left(I_i + \frac{439}{216} k_{1,I} - 8k_{2,I} + \frac{3680}{513} k_{3,I} - \frac{845}{4104} k_{4,I} \right) \right. \\
 &\quad \left. + \theta_2 \left(E_i + \frac{439}{216} k_{1,E} - 8k_{2,E} + \frac{3680}{513} k_{3,E} - \frac{845}{4104} k_{4,E} \right) \right] \\
 &\quad \left. - (\delta + \mu) \left(E_i + \frac{439}{216} k_{1,E} - 8k_{2,E} + \frac{3680}{513} k_{3,E} - \frac{845}{4104} k_{4,E} \right) \right\} \\
 k_{5,I} &= h * \left\{ \delta \left(E_i + \frac{439}{216} k_{1,E} - 8k_{2,E} + \frac{3680}{513} k_{3,E} - \frac{845}{4104} k_{4,E} \right) \right. \\
 &\quad \left. - (\gamma + \sigma + \mu) \left(I_i + \frac{439}{216} k_{1,I} - 8k_{2,I} + \frac{3680}{513} k_{3,I} - \frac{845}{4104} k_{4,I} \right) \right\} \\
 k_{5,R} &= h * \left\{ (\gamma + \theta_3 \sigma) \left(I_i + \frac{439}{216} k_{1,I} - 8k_{2,I} + \frac{3680}{513} k_{3,I} - \frac{845}{4104} k_{4,I} \right) \right. \\
 &\quad \left. - \mu \left(R_i + \frac{439}{216} k_{1,R} - 8k_{2,R} + \frac{3680}{513} k_{3,R} - \frac{845}{4104} k_{4,R} \right) \right\}
 \end{aligned} \tag{9}$$

6. The k_6 values are given by:

$$\begin{aligned}
 k_{6,S} &= h * \left\{ \Lambda - \beta(1 - \theta_1) \left(S_i - \frac{8}{27} k_{1,S} + 2k_{2,S} - \frac{3544}{2565} k_{3,S} + \frac{1859}{4104} k_{4,S} - \frac{11}{40} k_{5,S} \right) \right. \\
 &\quad \left[\left(I_i - \frac{8}{27} k_{1,I} + 2k_{2,I} - \frac{3544}{2565} k_{3,I} + \frac{1859}{4104} k_{4,I} - \frac{11}{40} k_{5,I} \right) \right. \\
 &\quad \left. + \theta_2 \left(E_i - \frac{8}{27} k_{1,E} + 2k_{2,E} - \frac{3544}{2565} k_{3,E} + \frac{1859}{4104} k_{4,E} - \frac{11}{40} k_{5,E} \right) \right] \\
 &\quad \left. - \mu \left(S_i - \frac{8}{27} k_{1,S} + 2k_{2,S} - \frac{3544}{2565} k_{3,S} + \frac{1859}{4104} k_{4,S} - \frac{11}{40} k_{5,S} \right) \right\} \\
 k_{6,E} &= h * \left\{ \beta(1 - \theta_1) \left(S_i - \frac{8}{27} k_{1,S} + 2k_{2,S} - \frac{3544}{2565} k_{3,S} + \frac{1859}{4104} k_{4,S} - \frac{11}{40} k_{5,S} \right) \right. \\
 &\quad \left[\left(I_i - \frac{8}{27} k_{1,I} + 2k_{2,I} - \frac{3544}{2565} k_{3,I} + \frac{1859}{4104} k_{4,I} - \frac{11}{40} k_{5,I} \right) \right. \\
 &\quad \left. + \theta_2 \left(E_i - \frac{8}{27} k_{1,E} + 2k_{2,E} - \frac{3544}{2565} k_{3,E} + \frac{1859}{4104} k_{4,E} - \frac{11}{40} k_{5,E} \right) \right] \\
 &\quad \left. - (\delta + \mu) \left(E_i - \frac{8}{27} k_{1,E} + 2k_{2,E} - \frac{3544}{2565} k_{3,E} + \frac{1859}{4104} k_{4,E} - \frac{11}{40} k_{5,E} \right) \right\} \\
 k_{6,I} &= h * \left\{ \delta \left(E_i - \frac{8}{27} k_{1,E} + 2k_{2,E} - \frac{3544}{2565} k_{3,E} + \frac{1859}{4104} k_{4,E} - \frac{11}{40} k_{5,E} \right) \right. \\
 &\quad \left. - (\gamma + \sigma + \mu) \left(I_i - \frac{8}{27} k_{1,I} + 2k_{2,I} - \frac{3544}{2565} k_{3,I} + \frac{1859}{4104} k_{4,I} - \frac{11}{40} k_{5,I} \right) \right\} \\
 k_{6,R} &= h * \left\{ (\gamma + \theta_3 \sigma) \left(I_i - \frac{8}{27} k_{1,I} + 2k_{2,I} - \frac{3544}{2565} k_{3,I} + \frac{1859}{4104} k_{4,I} - \frac{11}{40} k_{5,I} \right) \right. \\
 &\quad \left. - \mu \left(R_i - \frac{8}{27} k_{1,R} + 2k_{2,R} - \frac{3544}{2565} k_{3,R} + \frac{1859}{4104} k_{4,R} - \frac{11}{40} k_{5,R} \right) \right\}
 \end{aligned} \tag{10}$$

4. Results and Discussion

For simulations, we take initial values of the variables listed in Table 1 and parameters in Table 2. We take two different parameters θ_1 which are $\theta_{1_s} = 0.9$ and $\theta_{1_e} = 0.7$. Parameter $\theta_{1_s} = 0.9$ leads to a reproduction number R_0 smaller than 1, whereas $\theta_{1_e} = 0.7$ results in a reproduction number R_0 greater than 1.

Table 1. Variables and their initial values

Variables	Initial values
$S(0)$	100
$E(0)$	15
$I(0)$	20
$R(0)$	4

Table 2. Parameters and their values

Parameters	Values
Λ	10
β	0.2
θ_{1_s}	0.9
θ_{1_e}	0.7
θ_2	0.1
θ_3	0.3
μ	0.3
δ	0.3
γ	0.2
σ	0.2

1. For $R_0 < 1$, it can be obtained using $\theta_1 = \theta_{1_s} = 0.9$, that is,

$$R_0 = \frac{\Lambda\beta(1 - \theta_{1_s})[\delta + \theta_2(\gamma + \sigma + \mu)]}{\mu(\gamma + \sigma + \mu)(\delta + \mu)} = 0.5873 < 1$$

Because reproduction number $R_0 < 1$, the system converges to a disease free equilibrium:

$$\mathcal{E}^0 = (S^0, E^0, I^0, R^0) = \left(\frac{\Lambda}{\mu}, 0, 0, 0\right) = \left(\frac{10}{0.3}, 0, 0, 0\right) = (33.33, 0, 0, 0)$$

Simulation results for this case $R_0 < 1$ using the RKF45 method is shown in Figure 2. We observe that for this case, as time t gets large, the system contains only the Susceptible subpopulation S .

2. For $R_0 > 1$, it can be obtained using $\theta_1 = \theta_{1_e} = 0.7$, that is,

$$R_0 = \frac{\Lambda\beta(1 - \theta_{1_e})[\delta + \theta_2(\gamma + \sigma + \mu)]}{\mu(\gamma + \sigma + \mu)(\delta + \mu)} = 1.761 > 1$$

Because reproduction number $R_0 > 1$, the system converges to an endemic equilibrium:

$$\mathcal{E}^* = (S^*, E^*, I^*, R^*)$$

where

$$\begin{aligned} S^* &= \frac{(\gamma + \sigma + \mu)(\delta + \mu)}{\beta(1 - \theta_{1_e})[\delta + \theta_2(\gamma + \sigma + \mu)]} = 18.918 \\ E^* &= \frac{\Lambda}{(\delta + \mu)} - \frac{\mu(\gamma + \sigma + \mu)}{\beta(1 - \theta_{1_e})(\delta + \theta_2(\gamma + \sigma + \mu))} = 7.207 \\ I^* &= \delta \left(\frac{\Lambda}{(\delta + \mu)(\gamma + \sigma + \mu)} - \frac{\mu}{\beta(1 - \theta_{1_e})(\delta + \theta_2(\gamma + \sigma + \mu))} \right) = 3.088 \\ R^* &= \left(\frac{\Lambda\delta(\gamma + \theta_3\sigma)}{\mu(\delta + \mu)(\gamma + \sigma + \mu)} - \frac{\delta(\gamma + \theta_3\sigma)}{\beta(1 - \theta_{1_e})(\delta + \theta_2(\gamma + \sigma + \mu))} \right) = 2.676 \end{aligned}$$

Simulation results for this case $R_0 > 1$ using the RKF45 method is shown in Figure 3. We observe from this figure that as time evolves, the Infected subpopulation I does not vanish. This means the disease exists in the system all the time.




Figure 2. Simulation results using the RKF45 method for the case of disease-free equilibrium




Figure 3. Simulation results using the RKF45 method for the case of endemic equilibrium

Now we shall report the main results of our research. To do so, we recall the formula of the order of accuracy for the numerical method [14]:

$$R_i = \frac{\log\left(\frac{\text{Error}_i}{\text{Error}_{i+1}}\right)}{\log\left(\frac{\Delta t_i}{\Delta t_{i+1}}\right)}$$

where Error_i is the average error, Δt_i is the time step $h = \Delta t_i$ and $i = 0, 1, 2, \dots, n$. Note that the exact solution to the SEIR model is not known until when this paper is written, so we use ODE45 of the MATLAB software as our reference solution. When ODE45 function is used, we set the AbsTol and RelTol both to be 10^{-13} .

1. For $R_0 < 1$

For the case of disease-free equilibrium, we record the average absolute and average relative errors and report the orders of accuracy based on these errors. Tables 3-6 provide the errors and orders of accuracy for Susceptible, Exposed, Infected, and Recovered subpopulations. We observe that the orders of accuracy converge to five as the time step gets small.

Table 3. Order of Accuracy of RKF45 for *Susceptible S* with $R_0 < 1$.

h	Absolute error	Relative error	Order of accuracy based on absolute error	Order of accuracy based on relative error
0.8	6.601E-03	2.078E-04	-	-
0.4	3.581E-05	1.242E-06	7.526	7.387
0.2	3.794E-07	1.176E-08	6.561	6.721
0.1	1.058E-08	2.415E-10	5.164	5.606
0.05	3.395E-10	6.938E-12	4.962	5.122
0.025	1.088E-11	2.160E-13	4.963	5.005

Table 4. Order of Accuracy of RKF45 for *Exposed E* with $R_0 < 1$.

h	Absolute error	Relative error	Order of accuracy based on absolute error	Order of accuracy based on relative error
0.8	2.998E-03	1.569E-04		
0.4	4.618E-06	7.021E-07	9.343	7.805
0.2	5.063E-07	2.779E-08	3.189	4.659
0.1	1.874E-08	9.079E-10	4.755	4.936
0.05	6.084E-10	2.834E-11	4.945	5.001
0.025	1.921E-11	8.826E-13	4.985	5.005

Table 5. Order of Accuracy of RKF45 for *Infected I* with $R_0 < 1$.

h	Absolute error	Relative error	Order of accuracy based on absolute error	Order of accuracy based on relative error
0.8	2.208E-03	2.239E-04	-	-
0.4	3.003E-05	2.684E-06	6.200	6.383
0.2	6.169E-07	5.041E-08	5.605	5.735
0.1	1.581E-08	1.238E-09	5.286	5.348
0.05	4.500E-10	3.460E-11	5.135	5.161
0.025	1.344E-11	1.025E-12	5.064	5.076

Table 6. Order of Accuracy of RKF45 for *Recovered R* with $R_0 < 1$.

h	Absolute error	Relative error	Order of accuracy based on absolute error	Order of accuracy based on relative error
0.8	1.267E-03	2.195E-04	-	-
0.4	1.205E-05	1.999E-06	6.716	6.779
0.2	1.985E-07	3.207E-08	5.924	5.962
0.1	4.640E-09	7.366E-10	5.419	5.444
0.05	1.274E-10	2.008E-11	5.187	5.197
0.025	3.756E-12	5.942E-13	5.084	5.079

2. For $R_0 > 1$

For the case of endemic equilibrium, again we record the average absolute and average relative errors and report the orders of accuracy based on these errors. Tables 7-10 contain the error values and orders of accuracy for Susceptible, Exposed, Infected, and Recovered subpopulations. Once again, we observe that the orders of accuracy tend to five as the time step approaches small values. However, the order of accuracy can be less than five if the time step is too small.

Table 7. Order of Accuracy of RKF45 for *Susceptible S* with $R_0 > 1$.

h	Absolute error	Relative error	Order of accuracy based on absolute error	Order of accuracy based on relative error
0.8	1.642E-01	9.527E-03	-	-
0.4	3.719E-04	1.407E-05	8.786	9.403
0.2	7.132E-06	4.746E-07	5.705	4.891
0.1	3.072E-07	1.802E-08	4.537	4.719
0.05	1.012E-08	5.778E-10	4.923	4.963
0.025	3.204E-10	1.810E-11	4.982	4.996

Table 8. Order of Accuracy of RKF45 for *Exposed E* with $R_0 > 1$.

h	Absolute error	Relative error	Order of accuracy based on absolute error	Order of accuracy based on relative error
0.8	1.536E-01	3.741E-03	-	-
0.4	4.914E-04	1.547E-05	8.288	7.918
0.2	1.211E-05	2.878E-07	5.342	5.748
0.1	3.944E-07	8.512E-09	4.941	5.079
0.05	1.210E-08	2.540E-10	5.027	5.066
0.025	3.728E-10	7.800E-12	5.020	5.026

Table 9. Order of Accuracy of RKF45 for *Infected I* with $R_0 > 1$.

h	Absolute error	Relative error	Order of accuracy based on absolute error	Order of accuracy based on relative error
0.8	3.679E-02	2.448E-03	-	-
0.4	3.303E-04	1.984E-05	6.799	6.947
0.2	4.927E-06	2.720E-07	6.067	6.189
0.1	1.042E-07	5.409E-09	5.563	5.652
0.05	2.688E-09	1.356E-10	5.277	5.318
0.025	7.700E-11	3.918E-12	5.126	5.113

Table 10. Order of Accuracy of RKF45 for *Recovered R* with $R_0 > 1$.

h	Absolute error	Relative error	Order of accuracy based on absolute error	Order of accuracy based on relative error
0.8	2.027E-02	1.954E-03	-	-
0.4	1.220E-04	1.225E-05	7.376	7.317
0.2	1.149E-06	1.318E-07	6.730	6.539
0.1	1.784E-08	2.425E-09	6.009	5.764
0.05	4.454E-10	6.525E-11	5.324	5.216
0.025	1.544E-11	2.166E-12	4.850	4.913

We obtain by simulations that the fifth-order RKF45 method is numerically verified that the numerical order of accuracy matches with the theoretical order of accuracy. Some variations may occur, that is, the numerical order of accuracy can be larger and smaller than five, as recorded in Tables 3-10. It can be larger than five for relatively large time-step variations, and smaller than five for relatively too small time-step variations.

5. Conclusion

We have verified numerically that the theoretically fifth-order RKF45 method achieved the order five when time step is sufficiently small. Larger numerical order of accuracy occurred in our simulations when time step is quite large. We note that large numerical order of accuracy is advantageous for solving initial value problems numerically if accuracy is more concerned than the speed of computation. This research is limited to numerical experiments. Future research direction could take numerical analysis of the errors of RKF45 when implemented to solve an SEIR model.

6. Acknowledgement

The authors thank Sanata Dharma University for the opportunity of this research. This paper is an improvement of a part of the first author's Bachelor of Mathematics (S.Mat.) thesis [15] written at Sanata Dharma University in 2024 under the supervision of the second author.

References

- [1] Jana, A. K., *Numerical Methods: Theory and Engineering Applications*, Cambridge: Cambridge University Press, 2024.
- [2] Mungkasi, S., *Matematika: Jalan Kesejahteraan*, Yogyakarta: Sanata Dharma University Press, 2022.
- [3] Mathews, J. H., Fink, K. D., *Numerical Methods using MATLAB*, (Fourth Edition), Upper Saddle River: Prentice Hall, 2004.
- [4] Jiao, J., Liu, Z., Cai, S., "Dynamics of an SEIR model with infectivity in incubation period and homestead-isolation on the susceptible," *Applied Mathematics Letters*, vol. 107, art. 106442, 2020.
- [5] Palma, D. I., Mungkasi, S., "Fourth and fifth-order Runge-Kutta methods for solving a susceptible-exposed-infected-recovered mathematical model of the spread of COVID-19," *AIP Conference Proceedings*, vol. 3074, art. 020006, 2024.
- [6] Rosari, F. I., Mungkasi, S., "Co-dynamics model of the spread of malaria and COVID-19 with numerical solutions using the third-order and the fourth-order Runge-Kutta methods," *AIP Conference Proceedings*, vol. 3074, art. 020008, 2024.
- [7] Setiawan, L. I., Mungkasi, S., "Penyelesaian model epidemi SIR menggunakan metode Runge-Kutta orde empat dan metode Adams-Bashforth-Moulton," *Komputasi: Jurnal Ilmiah Ilmu Komputer dan Matematika*, vol. 18, no. 2, pp. 55-61, 2021.
- [8] Alkhazzan, A., Wang, J., Nie, Y., Khan, H., Alzabut, J., "A novel SIRS epidemic model for two diseases incorporating treatment functions, media coverage, and three types of noise," *Chaos, Solitons & Fractals*, vol. 181, art. 114631, 2024.
- [9] Batiha, I. M., Ogilat, O., Bendib, I., Ouannas, A., Jebril, I. H., Anakira, N., "Finite-time dynamics of the fractional-order epidemic model: Stability, synchronization, and simulations," *Chaos, Solitons & Fractals: X*, vol. 13, art. 100118, 2024.
- [10] Chen, B., Chen, R., Zhao, L., Ren, Y., Zhang, L., Zhao, Y., Lian, X., Yan, W., Gao, S., "High-resolution short-term prediction of the COVID-19 epidemic based on spatial-temporal model modified by historical meteorological data," *Fundamental Research*, vol. 4, no. 3, pp. 527-539, 2024.
- [11] Esterhuizen, W., Sauerteig, P., Streif, S., Worthmann, K., "MPC without terminal ingredients tailored to the SEIR compartmental epidemic model," *Systems & Control Letters*, vol. 193, art. 105908, 2024.

- [12] Guan, G., Guo, Z., Xiao, Y., “Dynamical behaviors of a network-based SIR epidemic model with saturated incidence and pulse vaccination,” *Communications in Nonlinear Science and Numerical Simulation*, vol. 137, art. 108097, 2024.
- [13] Turkyilmazoglu, M., “Solutions to SIR/SEIR epidemic models with exponential series: Numerical and non numerical approaches,” *Computers in Biology and Medicine*, vol. 183, art. 109294, 2024.
- [14] Hidayat, N., Suhariningsih, Suryanto, A., Mungkasi, S., “The significance of spatial reconstruction in finite volume methods for the shallow water equations,” *Applied Mathematical Sciences*, vol. 8, no. 29, pp. 1411-1420, 2014.
- [15] Widi, L. K., *Model Matematis SEIR Pengaruh Isolasi Mandiri pada Kelompok Rentan dalam Penyebaran Penyakit Menular dan Penyelesaiannya dengan Metode RKF45 Orde Lima*, Skripsi (Bachelor’s Thesis), Yogyakarta: Sanata Dharma University, 2024.