



HOME / Editorial Team

Editorial Team

Editor in Chief

- [Dr. Florentinus Dika Octa Riswanto](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia

Vice Editor in Chief

- [Dr. apt. Dita Maria Virginia](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia

Advisory Editorial Board

- [apt. Dina Christin Ayuning Putri, M.Sc.](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia
- [Dr. apt. Dewi Setyaningsih](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia
- [dr. Fenty M.Kes., Sp.PK.](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia
- [apt. Phebe Hendra M.Si., Ph.D.](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia
- [Prof. Dr. apt. Sri Hartati Yuliani](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia
- [Dr. apt. Yosef Wijoyo](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia

Managing Editor

- [apt. Handika Immanuel S.Farm., M.Si.](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia
- [apt. Michael Raharja Gani, M.Farm.](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia
- [apt. Zita Dhirani Pramono S.Farm., M.Clin.Pharm.](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia
- Anjar Bayu Saputra, Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia

Editorial Board

- [Dr. Yohanes Dwiatmaka](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia
- [apt. T.B. Titien Siwi Hartayu, M.Kes., Ph.D.](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia
- [Dr. apt. Rini Dwiastuti](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia
- [apt. Christofori Maria Ratna Rini Nastiti Ph.D.](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia
- [Dr. Jeffry Julianus M.Si.](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia
- [apt. Aris Widayati Ph.D.](#), Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Indonesia
- [Prof. Dr. apt. Aty Widawaruyanti](#), Faculty of Pharmacy, Universitas Airlangga, Indonesia
- [Prof. Dr. apt. Auliya A. Suwantika](#), Faculty of Pharmacy, Universitas Padjadjaran, Indonesia
- [Prof. Dr. Bandana Saini](#), Faculty of Pharmacy, University of Sydney, Australia
- [Dr. apt. Christine Patramurti](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia
- [Prof. apt. Irma Melyani Puspitasari Ph.D.](#), Faculty of Pharmacy, Universitas Padjadjaran, Indonesia
- [Prof. Dr. Monet M. Loquias](#), College of Pharmacy, University of the Philippines, Philippines
- [Dr. Phayom Sookaneknun](#), Faculty of Pharmacy, Mahasarakham University, Thailand
- [Prof. Dr. P.T. Thomas](#), Faculty of Health and Medical Sciences, Taylor's University, Malaysia
- [apt. Rano Kurnia Sinuraya, MKM, Ph.D.](#), Faculty of Pharmacy, Universitas Padjadjaran, Indonesia
- [Prof. Dr. apt. Rizky Abdulah](#), Faculty of Pharmacy, Universitas Padjadjaran, Indonesia
- [Prof. Dr. apt. Sri Hartati Yuliani](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia
- [Prof. Dr. Yashwant Pathak](#), College of Pharmacy, University of South Florida, United States
- [Dr. apt. Yustina Sri Hartini](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia
- [Prof. Dr. apt. Enade Perdana Istyastono](#), Department of Pharmacy, School of Medicine and Health Sciences, Atma Jaya Catholic University of Indonesia, Indonesia

JOURNAL MENU

Focus and Scope

Editorial Team

Publication Ethics

Author Guidelines

Reviewers

Peer Review Process

Journal History

Indexing & Archiving

Accreditation Document

Article Processing Charge

Journal Subscription

Generative AI Usage Policy

MAKE A SUBMISSION

ARTICLE TEMPLATE



NATIONAL ACCREDITATION



Editorial Office:

FACULTY OF PHARMACY
UNIVERSITAS SANATA DHARMA
Kampus III, Paingan, Maguwoharjo,
Depok, Sleman, Yogyakarta 55282



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Jurnal Farmasi Sains dan Komunitas (Journal of
Pharmaceutical Sciences and Community)

Published by [Faculty of Pharmacy, Universitas Sanata
Dharma Yogyakarta](#)





About the Journal



Jurnal Farmasi Sains dan Komunitas (Journal of Pharmaceutical Sciences and Community) or J. Pharm. Sci. Community firstly published in 2003, is a peer-reviewed, open access journal that publishes research articles, review articles, as well as short communication in various pharmaceutical fields, including Pharmaceutical Technology and Pharmaceutics, Pharmaceutical Analysis, Medicinal Chemistry, Pharmacology and Toxicology, Pharmaceutical Biology, Community Pharmacy, and Clinical Pharmacy. *J. Pharm. Sci. Community* is managed and published in Yogyakarta, Indonesia by Faculty of Pharmacy, Sanata Dharma University Yogyakarta

Journal Title	: Jurnal Farmasi Sains dan Komunitas (Journal of Pharmaceutical Sciences and Community)
Frequency	: 2 issues per year May and November
DOI	: https://doi.org/10.24071/jpsc
ISSN	: 2527-7146 (Online) 1693-5683 (Print)
Editor in Chief	: Florentinus Dika Octa Riswanto
Publisher	: Faculty of Pharmacy, Universitas Sanata Dharma
Indexing	: Garuda , DOAJ , Sinta , Dimensions

Since October 2018, *J. Pharm. Sci. Community* has been "**Accredited with Grade 2**" by the Indonesian government. The recognition has been extended by DIKTISAINTEK and published in the [Director Decree No. 1295/C/C3/KPT/2026](#). This accreditation status has effective from Volume 21 No. 2 (2024) until Volume 26 No. 1 (2029).

Official Migration to OJS 3

2026-05-18

Dear Authors, Reviewers, and Readers,

We would also like to inform you that *Jurnal Farmasi Sains dan Komunitas* (Journal of Pharmaceutical Sciences and Community) or *J. Pharm. Sci. Community* currently extends its national accreditation ranking of "SINTA 2" (Decree of the Director General of Research and Development No. 295/C/C3/KPT/2026, dated January 2, 2026). We believe this achievement is due to the contributions and trust of our authors.

On behalf of *J. Pharm. Sci. Community*, we officially inform you regarding the official Open Journal System (OJS) migration from OJS 2 to OJS 3. This migration has been undertaken to enhance the security and quality of article management within our journal.

[READ MORE >](#)

JOURNAL MENU

- Focus and Scope
- Editorial Team
- Publication Ethics
- Author Guidelines
- Reviewers
- Peer Review Process
- Journal History
- Indexing & Archiving
- Accreditation Document
- Article Processing Charge
- Journal Subscription
- Generative AI Usage Policy

MAKE A SUBMISSION

ARTICLE TEMPLATE



NATIONAL ACCREDITATION



CURRENT ISSUE

Vol. 23 No. 1 (2026)

Articles

Antibacterial Activity Test Against *Cutibacterium acne* and Contaminant Testing of Spirulina (*Arthrospira platensis*) Anti-Acne Serum

DOI: <https://doi.org/10.24071/jpsc.v23i1.1069>

Hilma Putri, Phebe Hendra (Author)

1-10



Stability of Orally Dissolving Film of *Zingiber officinale* Roscoe Extract With HPMC And Maltodextrin as Polymer

DOI: <https://doi.org/10.24071/jpsc.v23i1.1075>

Srie Rezeki Nur Endah, Iis Nur Asiah, Susanti Susanti, Nitya Nurul Fadilah, Ali Nofriyaldi (Author)

11-17



Relationship of Knowledge and Attitudes on Vitamin D Status in Active Smokers

DOI: <https://doi.org/10.24071/jpsc.v23i1.1077>

Amelia Lorensia, Marthy Meliana Ariyanti Jalmav, Eva Amilia, Amirah Dyandra Sybil (Author)

18-34



Medication Adherence and Antihypertensive Treatment Intensification on Blood Pressure Control at Bali Mandara Hospital

DOI: <https://doi.org/10.24071/jpsc.v23i1.1067>

Desak Putu Meilinda Asri Swantari, Dita Maria Virginia (Author); Phebe Hendra (Translator)

35-44



Anticancer Activity Studies of Self-nanoemulsifying Drug Delivery Systems (SNEDDS) *Hibiscus sabdariffa* L. as Cisplatin Co-Chemotherapy in T47D Breast Cancer Cells

DOI: <https://doi.org/10.24071/jpsc.v23i1.1079>

Nurma Wahyuningtyas , Laela Hayu Nurani, Nuri Ari Efiana, Nurkhasanah Nurkhasanah, Fara Azzahra, Any Guntarti, Septiana Indratmoko, Citra Ariani Edityaningrum (Author)

45-52



The Analysis of Health Professional's Knowledge and Attitudes Related to The Implementation of Pharmacovigilance in Pharmaceutical Service Facilities

DOI: <https://doi.org/10.24071/jpsc.v23i1.1080>

Ema Nillafita Kusuma, Aas Sa'adah, Dian Farida Ismyama (Author)

53-64



Employing Ensemble Protein-Ligand Interaction Fingerprints to Mimic Induced-Fit Theory in Structure-Based Virtual Screening Targeting Dipeptidyl Peptidase IV

DOI: <https://doi.org/10.24071/jpsc.v23i1.1070>

Enade Perdana Istyastono, Bonifacius Ivan Wiranata, Florentinus D.O. Riswanto, Fransiska Kurniawan, Tasia Amelia, Nunung Yuniarti, Eko Adi Prasetyanto (Author)

65-74



Beyond Use Date of Rifampicin Pulverized and Its Influencing Factors

DOI: <https://doi.org/10.24071/jpsc.v23i1.1083>

Zaldy Rusli, Septia Andini, Nisa Najwa Rokhmah, Idul Suci Fitriana, Mutiara Apriliana Suvitri (Author)

75-83



Formulation and Physical Evaluation of Antihypertensive Sustained-Release Tablet From Matoa Leaf Ethanolic Extract (*Pometia pinnata* J.R & G.Forst)

DOI: <https://doi.org/10.24071/jpsc.v23i1.1084>

Arifuddin Yunus, Siti Qurrataayun, Lutfiah Lutfiah (Author)

84-91



Evaluation of Antibiotic Use in Paediatric Inpatients Diagnosed with Urinary Tract Infection at a Private Hospital in Yogyakarta

DOI: <https://doi.org/10.24071/jpsc.v23i1.1085>

Christina Ika Sindudisastra, Titien Siwi Hartayu (Author)

92-104



The Effect of Medication Adherence and Controlled HbA1c to Blood Pressure and ASCVD Risk in Hypertensive and Type 2 Diabetes Mellitus Patients: A multicenter study in Primary Health Care

DOI: <https://doi.org/10.24071/jpsc.v23i1.1071>

C. Vera Dwi Pratiwi, Phebe Hendra, Agustina Rimanita Fortinasari, Dita Maria Virginia (Author)

105-113



Optimization and Validation of Caffeine in Robusta Coffee (*Coffea canephora*), Black Tea (*Camellia sinensis*), and Cocoa (*Theobroma cacao* L.) Using RP-HPLC Liquid-liquid Extraction

DOI: <https://doi.org/10.24071/jpsc.v23i1.1086>

Lindawati Setyaningrum, Aliyah Purwanti, Dina Trianggaluh Fauziah, Retno Eka Lestari, Dyah Purwaningtyas, Sindi Anggraiyani, Muhammad Yuwono (Author)

114-123



Evaluation of the Analgesic Activity of Earthworm Gel (*Lumbricus rubellus*) in Male Mice with Formalin-Induced Pain Model

DOI: <https://doi.org/10.24071/jpsc.v23i1.1087>

Yovita Mercya, Ayu Amisa Fajrianti (Author)

124-132



Optimization of temperature and drying time in the process of making paracetamol tablets using PVP K-25 as a Binder

DOI: <https://doi.org/10.24071/jpsc.v23i1.1072>

Yanita Yomberlin Sinta Maria Barangan, Agatha Budi Susiana Lestari (Author)

133-143



Drug Information Services at Primary Healthcare, A Study of Patient Satisfaction Analysis among Diabetes Mellitus Patients in Kapanewon Depok Yogyakarta Indonesia

DOI: <https://doi.org/10.24071/jpsc.v23i1.1074>

Geraldi Fredrik Daud Tumimbang, Maria Wisnu Donowati (Author)

144-150



Phytochemical Screening and *In Vitro* Anti-Inflammatory Activity through Protein Denaturation Inhibition of Tigarun (*Crataeva nurvala* Buch. -Ham.) Stem Bark

DOI: <https://doi.org/10.24071/jpsc.v23i1.1089>

Fadlilaturrahmah Fadlilaturrahmah, Mia Fitriana, Alya Hanifa, Resa Amalia Syabana (Author)

151-159



Assessment of the Quality of Life of Family Caregivers of Hemodialysis Patients with Chronic Kidney Disease in Yogyakarta

DOI: <https://doi.org/10.24071/jpsc.v23i1.1088>

Pinasti Utami, Binar Sukma Cahyarni, Miranda Miranda, Bangunawati Rahajeng, Triana Arum Kusumaningtyas (Author)

160-166



Antibacterial Activity and Wound Healing of Excision Gel of Ethanol Extract of Sintrong (*Crassochevallum*

crepidioides)(Benth.) S. Moore.) Leaves in Mice Infected *Staphylococcus aureus*

DOI: <https://doi.org/10.24071/jpsc.v23i1.1081>

Candrika Candrika, Aminah Dalimunthe, Hakim Bangun (Author)

167-179



The Effect of Glycerin and Squalane on Skin Moisture in Green Tea (*Camellia sinensis* L.) Nanostructure Lipid Carrier Preparation

DOI: <https://doi.org/10.24071/jpsc.v23i1.1073>

Cynthia Gracella Gamas, Rini Dwiastuti, Lutfi Chabib (Author)

180-192



Improving Adherence to ARV Therapy in person Living with HIV/AIDS (PLWHA) by The Community-Based Interactive Approach (CBIA) in Temanggung District, Indonesia

DOI: <https://doi.org/10.24071/jpsc.v23i1.1076>

Sisilia Esti Purwaningsih, Titien Siwi Hartayu (Author)

193-203



Cognitive Levels, Perceptions, and Attitudes of Pharmacists at Pharmacies toward Herbal Medicine

DOI: <https://doi.org/10.24071/jpsc.v23i1.1082>

Diah Ari Purwaningsih, Yosef Wijoyo, Nunung Priyatni Waluyatiningsih (Author)

204-222



[VIEW ALL ISSUES](#) ➤



Editorial Office:

FACULTY OF PHARMACY
UNIVERSITAS SANATA DHARMA
Kampus III, Paingan, Maguwaharjo,
Depok, Sleman, Yogyakarta 55282



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Jurnal Farmasi Sains dan Komunitas (Journal of Pharmaceutical Sciences and Community)

Published by [Faculty of Pharmacy, Universitas Sanata Dharma Yogyakarta](#)

Optimization of Temperature and Drying Time in the Process of Making Paracetamol Tablets Using PVP K-25 as a Binder

Yanita Yomberlin Sinta Maria Barangan¹, Agatha Budi Susiana Lestari^{2*}

¹Pharmacist Profession Education, Faculty of Pharmacy, Sanata Dharma University, Yogyakarta 55282, Indonesia

²Department of Pharmaceutical, Faculty of Pharmacy, Sanata Dharma University, Yogyakarta 55282, Indonesia

 <https://doi.org/10.24071/jpsc.v23i1.1072>

 J. Pharm. Sci. Community, 2026, 23(1), 133-143

Article Info

Received: 2025-06-22

Revised: 2026-04-09

Accepted: 2026-04-11

***Corresponding author:**

Agatha Budi Susiana Lestari

email:

a_budi@usd.ac.id

Keywords:

drying time; factorial design; paracetamol tablet; temperature of drying; PVP K-25

ABSTRACT

Wet granulation is a crucial technique in tablet manufacturing since it improves the physical properties of the formulation, such as the flowability and compressibility of paracetamol. One of the critical stages in this process is drying the granules using oven heating. This study aims to optimize the drying temperature and time and evaluate their effects on the physical properties of paracetamol granules and tablets. PVP K-25 was selected as a binder due to its ability to enhance tablet compactness. A full factorial experimental design was used with two factors and two levels, optimizing drying temperature (°C) and drying time (hours) based on the physical properties of granules and tablets. Data were analyzed using ANOVA and Design Expert 13 software to determine the optimal formulation. The results showed that drying temperature, drying time, and their interaction significantly affected the physical characteristics of granules and tablets. The optimal formulation was achieved at 55°C for 4 hours. Formula A (40°C, 6 hours) and Formula B (60°C, 3 hours) demonstrated the best results in producing tablets with desirable physical properties. Thus, optimizing drying temperature and time can enhance the quality of paracetamol granules and tablets.

INTRODUCTION

Oral drug administration is the most widely used method in clinical practice, primarily in the form of tablets. As a solid dosage form, tablets must remain visually and functionally intact. However, despite technological advancements, tablet defect during production remains a challenge and may even worsen due to the complexity of the manufacturing process and the stringent quality standards required (Ardiansyah *et al.*, 2022; Zaman & Sopyan, 2020). Tablet manufacturing can be done using three main methods: wet granulation, dry granulation, and direct compression. In this study, the wet granulation method was used to produce paracetamol tablets because it enhances compactibility and flow properties with the help of a binder (Abimanyu *et al.*, 2023; Sudarsono *et al.*, 2021). The tablet formulation must consider

excipients such as fillers, disintegrants, binders, lubricants, and glidants. One commonly used binder is polyvinylpyrrolidone (PVP), specifically PVP K-25, due to its good flow properties, ability to produce compact tablets, and inert and stable nature. PVP K-25 also has high solubility in various solvents and good compatibility with many types of drugs (Direktorat Jenderal Kefarmasian dan Alat Kesehatan, 2020; Cahyani *et al.*, 2023).

In the wet granulation method, the drying process determines the moisture content of the granules, which affects tablet hardness. The drier the tablet, the lower its porosity, resulting in stronger interparticle bonds. Longer drying times and higher temperatures increase water evaporation, producing drier granules, but this can also increase tablet friability (Maskuriah *et al.*, 2021).

METHODS

Materials

Paracetamol (active ingredient), polyvinylpyrrolidone PVP K-25 (binder), magnesium stearate and talcum (lubricants), corn starch (disintegrant), and distilled water (solvent).

Equipment

Glassware, analytical balance (OHAUS Pioneer PA213), mixer (Erweka AR-401), sieves No. 12 and 14 (Patraproduk), oven (MEMERT BE 500), moisture balance (Kern MLS 50-3C), flowability tester (Erweka AR 401), volumenometer (HY-100B), rotary tablet press (Shanghai Develop Machinery ZP-5B), caliper 0.05mm (NRT-PRO), attrition tester (ATMI), hardness tester (Pharmas Test PTB 302), and disintegration tester (Develop BJ-2).

Preparation of PVP K-25 solution

This study used one concentration as a binder, which was 3% w/w. The solution was made by dissolving PVP K-25 powder in aquadest as a solvent.

Wet granulation method

Paracetamol, lactose, magnesium stearate, talc, and corn starch are weighed according to the specified amounts (Table 1). Paracetamol and lactose powders are mixed using a mixer until homogeneous. The mixture is transferred to a plastic tray, and the PVP K-25 solution is gradually added as a wetting agent until a damp granulation mass is formed. Then, a small amount of corn starch is added to the granulation mixture and stirred until homogeneous. The granulation mass is sieved through a 12-mesh sieve, and the weight of the wet granules is measured. The granules are then dried at 40°C and 60°C for 3 and 6 hours using an oven.

Moisture content

After drying, the granules undergo moisture content testing using a moisture balance with a sample weighing 1 gram. Moisture content can be calculated using the following formula:

$$\%MC = \frac{\text{weight of wet granules} - \text{weight of dry granules}}{\text{weight of wet granules}} \times 100\%$$

The optimal moisture content for granules is 1%–5% (Mariyani *et al.*, 2016). The dried granules are then sieved again using a 14-mesh sieve.

Lubrication method

The dried granules are placed into a mixer, and the remaining corn starch is added and mixed. Then, magnesium stearate and talcum are incorporated into the mixture and blended until homogeneous.

Flow time

Granule flow time testing is performed using a flowability tester. A hundred grams of granules are placed into the funnel, and the flow time is measured until all the granules pass through. The result is recorded in seconds, with the standard ≤ 10 seconds (Noval *et al.*, 2021).

Angle of repose

The angle of repose test is conducted by weighing 100 grams of granules and placing them into a flow funnel. As the granules flow, the angle of repose is determined by measuring the steepness of the pile formed. The angle of repose is determined using a formula (United States Pharmacopeia Convention, 2024a) :

$$\tan \alpha = \frac{h}{r}$$

The angle of repose is classified into several levels: excellent (25–30°), good (31–35°), fair (36–40°), passable (41–45°), poor (46–55°), very poor (56–65°), and extremely poor (>66°) (Taylor and Aulton, 2022).

Table 1. Formulation, Temperature, and Granule Drying Time of Paracetamol Tablets

Ingredients	Formula (mg/tablet)			
	F1	Fa	Fb	Fab
Paracetamol	500	500	500	500
Lactose	320	320	320	320
PVP K-25 3%	3.6	3.6	3.6	3.6
Corn starch	20	20	20	20
Talc	6.5	6.5	6.5	6.5
Magnesium stearate	6.5	6.5	6.5	6.5
Temperature for drying (°C)	40	60	40	60
Drying time (hours)	3	3	6	6

Compressibility index

The compressibility index test for granules was conducted using a volumeter. First, 40 grams of granules were weighed and placed into a 100 mL graduated cylinder, and the initial volume of the granules was recorded. The graduated cylinder was then mounted on the volumeter and tapped 500 times, after which the final volume of the granules was recorded. A good compressibility index for granules ranges from 5-16% or <20% and an optimal Hausner ratio falls between 1.12-1.18. The compressibility index is calculated using the following equation:

$$\text{Compressibility Index} = 100 \times \left(\frac{V_1 - V_2}{V_1} \right)$$

Notes:

V1 : Initial volume

V2 : Final tapped volume

Tablet Manufacturing

The mixed granules are then compressed into tablets using a single-punch tablet machine. After compression, the physical properties of the tablets are tested to evaluate their quality.

Evaluation of Granule Characteristic Organoleptic test

Organoleptic testing is the initial step in achieving maximum objectivity in tablet evaluation. This test involves visually observing the tablets using human senses, including color, odor, shape, and surface texture (Syukri, 2018).

Weight Variation Test

Ten tablets are individually weighed with precision. The active ingredient content in each tablet is determined and expressed as a percentage of the labeled amount. The obtained assay results from each tablet are then used to calculate the acceptance value according to the established criteria. The acceptance value for weight uniformity is met if $L1 = 15.0$ and $L2 = 25.0$ (Direktorat Jenderal Kefarmasian dan Alat Kesehatan, 2020).

Tablet hardness test

Using a hardness tester, tablet hardness testing is used as a reference to determine whether tablet meets the required standards. Ten tablets are used as test samples, each placed in the center of the hardness tester in a vertical position. The device then applies pressure until the tablet breaks. The acceptance criterion for tablet hardness is 4-8 kg for oral tablets (Rori *et al.*, 2016).

Tablet friability test

Friability, a parameter used to measure the tablet's surface resistance to friction during packaging and transportation, is measured using a friabilator. Ten tablets, cleaned of dust, are weighed and recorded before being placed in the testing device, which operates at a speed of 25 rpm for 4 minutes. The tablets are then cleaned and reweighed to determine the weight loss. The acceptance criterion for tablet friability is less than 1% weight loss from the total tablet mass (United States Pharmacopeia Convention, 2024a). The acceptance results are calculated using the formula in the following equation:

$$F = \frac{W_1 - W_2}{W_1} \times 100\%$$

Notes:

F : Friability percentage

W1 : Initial tablet weight

W2 : Final tablet weight

Disintegration time test

The disintegration test is an essential tablet evaluation with specific requirements to ensure rapid therapeutic effects. The test is conducted using a disintegration tester, where six tablets are placed in each tube of the device. Water at a temperature of $37^\circ \pm 2^\circ\text{C}$ is used as the medium, and the testing duration is 15 minutes. At the end of the test, all tablets are observed to ensure complete disintegration, and the disintegration time is recorded. If one or two tablets fail to disintegrate completely, the test is repeated with an additional 12 tablets. The disintegration time requirement is met if at least 16 out of 18 tested tablets disintegrate. The acceptance criterion for uncoated tablets is a disintegration time of less than or equal to 15 minutes (Direktorat Jenderal Kefarmasian dan Alat Kesehatan, 2020).

Data Analysis

The data were analyzed using the Shapiro-Wilk test in SPSS 23 (IBM Corp., Chicago, USA). Based on the data, the factorial design equation was calculated using the formula ($Y = \beta_0 + \beta_1.X_A + \beta_2.X_B + \beta_{12}.X_A.X_B$), where Y represents the observed response, X_A and X_B represent the levels, and β_0 , β_1 , β_2 , and β_{12} are coefficients. The goal is to determine the optimal formulation of drying temperature and time for granules. Data with a normal distribution were then analyzed using Design Expert 13 (free trial) to obtain

ANOVA values and determine the optimal formulation.

RESULTS AND DISCUSSION

A total of 720 respondents participated in our Table 2 presents the results of the physical properties test of paracetamol granules. Analysis using Design Expert software with the factorial

design method indicates that the physical properties of paracetamol granules and tablets were obtained under optimal drying conditions. The contour plot in Figure 1 (a), (b), (c) and (d) illustrates the relationship between drying time and temperature on the physical properties of paracetamol granules.

Table 2. Results of Evaluation of Paracetamol Granule Characteristic

Granule Quality Control	F1	Fa	Fb	Fab
Moisture Content (%)	4.931 ± 0.003	2.430 ± 0.0045	3.267 ± 0.0030	1.107 ± 0.0046
Flow Time (s)	3.8 ± 0.7	2.43 ± 0.9	2.9 ± 0.7	1.56 ± 0.5
Angle of Repose (°)	34.71 ± 1.47	30.39 ± 1.07	31.40 ± 1.14	27.45 ± 1.80
Compressibility index	9.907 ± 3.214	7.508 ± 2.516	8.411 ± 2.516	6.911 ± 3.055

Note for temperature and drying time respectively:

Formula 1: 40°C and 3 hours.

Formula a: 60°C and 3 hours.

Formula b: 40°C and 6 hours.

Formula ab: 60°C and 6 hours.




Figure 1. Contour plot of granule test response (a) moisture content; (b) powder flow; (c) angle of repose; and (d) compressibility index.

Table 3. Effect of Temperature, Time, and Interaction on Moisture Content

Factor	Effect Value	%Contribution	<i>p</i> -value	<i>p</i> -value Equation
Temperature	-2.3305	70.6282	<0.0001	<0.0001
Time	-1.4935	28.9932	<0.0001	(Significant)
Interaction	0.1705	0.3780	<0.0001	

Table 4. Effect of Temperature, Time, and Interaction on Powder Flow

Factor	Effect Value	%Contribution	<i>p</i> -value	<i>p</i> -value Equation
Temperature	-1.35	45.935	0.0114	0.0293
Time	-0.88	19.666	0.0649	(Significant)
Interaction	0.015	0.007	0.9688	

The Effect of Temperature and Drying Time on the Physical Properties of Granule Moisture content

Table 2 shows that all formulations meet the moisture content requirement of 1-5% (Noval, dkk., 2021). Formula ab has the lowest moisture content (1.107%) compared to formula a, b and formula 1 (respectively 2.430%, 3.267% and 4.931%) due to the higher drying temperature of 60°C. Figure 1(a) supports this finding with a contour plot, where red shading indicates high moisture content and blue shading indicates low moisture content. Higher temperatures accelerate the evaporation of the binding liquid during the drying process. The factorial equation predicts moisture content using $Y = 12.279 - 0.142100X_A - 0.7818X_B + 0.00568X_AX_B$, where Y represents the moisture content response, X_A represents temperature, X_B represents time, and X_AX_B represents the interaction between temperature and time.

The analysis results show that all factors significantly influence the moisture content (p -value < 0.0001). Temperature is the most dominant factor with a contribution of 70.6282%, followed by drying time at 28.9932%. Although the interaction between temperature and drying time is statistically significant, its contribution is very small (0.3780%), so its influence can be practically ignored. The negative effect values for temperature and drying time indicate that increasing these two factors tends to decrease the moisture content (Table 3).

Flow time

The fastest flow time value was found in Fab (1.56±0.5 s), followed by Fa (2.43±0.9 s), then Fb (2.9±0.7 s), and the slowest in F1 (3.8±0.7 s). This shows that Fab granules have

the best flow properties, while F1 has the least good flow properties.

According to Table 4, the analysis results show that only the temperature factor has a significant effect on the powder flow (p -value = 0.0114 < 0.05), with a contribution of 45.935%. The time factor, although having a relatively moderate contribution (19.666%), does not show statistical significance (p -value = 0.0649), so its effect on the powder flow is inconsistent. Meanwhile, the interaction between temperature and drying time has no significant effect (p -value = 0.9688) with a very small contribution (0.007%). Thus, the powder flow variation is mainly controlled by temperature changes.

Angle of repose

The smallest angle of repose value was found in Fab (27.45±1.80°), followed by Fa (30.39±1.07°), then Fb (31.40±1.14°), and the largest in F1 (34.71±1.47°). This result indicates that Fab granules have the best flow (freer), while F1 tends to be more cohesive.


Based on Table 5, the analysis results showed that temperature (p = 0.0009) and drying time (p = 0.0049) significantly influenced the angle of repose, contributing 53.1982% and 30.3351%, respectively. Meanwhile, the interaction between temperature and drying time was not significant (p = 0.8254) and had a very small contribution (0.1063%). The negative effect values for both main factors indicated that increasing temperature and drying time tended to decrease the angle of repose value. Thus, the angle of repose variation was primarily controlled by temperature and drying time factors independently.

Table 5. Effect of Temperature, Time, and Interaction on Angle of Repose

Factor	Effect Value	%Contribution	<i>p</i> -value	<i>p</i> -value Equation
Temperature	-4.135	53.1982	0.0009	0.0016
Time	-3.125	30.3351	0.0049	(Significant)
Interaction	0.185	0.1063	0.8254	

Table 6. Effect of Temperature, Time, and Interaction on Compressibility Index

Factor	Effect Value	%Contribution	<i>p</i> -value	<i>p</i> -value Equation
Temperature	-1.949	74.564	<0.0001	<0.0001
Time	-1.047	21.491	<0.0001	(Significant)
Interaction	0.449	3.960	<0.0001	

**Figure 2.** Interaction of Drying Temperature and Time on (a) Moisture Content; (b) Powder Flow Response; (c) Angle of Repose Response; and (d) Compressibility Index Response.

Compressibility index

The lowest compressibility index value was found in Fab ($6.911 \pm 3.055\%$), followed by Fa ($7.508 \pm 2.516\%$), then Fb ($8.411 \pm 2.516\%$), and the highest was in F1 ($9.907 \pm 3.214\%$). This indicates that Fab granules have the best flow and compaction capabilities, while F1 has the least good.

According to Table 6, the analysis results showed that all factors significantly influenced the response (p -value < 0.0001). However, based on the percentage contribution, temperature was the most dominant factor in explaining response variability (74.564%), followed by time (21.491%), while the interaction effect was relatively small (3.960%). This indicates that

compressibility index is primarily controlled by temperature of drying.

Figure 2(a) illustrates the interaction between temperature and time by two non-parallel lines, one red and one black. The interaction is minimal since the lines do not intersect or touch. The graph indicates that higher temperatures and longer drying times reduce granule moisture content (Hiremath *et al.*, 2019 ; Elisabeth *et al.*, 2018). Based on the graph in Figure 2(b), the non-parallel red and black lines represent an interaction between temperature and drying time. Increasing temperature from 40°C to 60°C reduces granule flow time, indicating improved flow properties (United States Pharmacopeia Convention, 2024a; Elisabeth *et al.*, 2018). Figure 2(c) shows that as the drying temperature increases from 40°C to 60°C, the angle of repose of the granules tends to decrease, both at 3 hours (black line) and 6 hours (red line). This decrease in the angle of repose suggests that higher temperatures enhance granule flow properties, making the granules easier to flow at higher temperatures (Elisabeth *et al.*, 2018). Figure 2(d) demonstrates that increasing the temperature from 40°C to 60°C significantly reduces the compressibility index. The black line represents a drying time of 3 hours, while the red line represents a drying time of 6 hours. Granules are more sensitive to temperature at shorter drying times, whereas the effect of temperature becomes more moderate with longer drying times (Cahyani *et al.*, 2023).

The Effect of Temperature and Drying Time on the Physical Properties of Tablet

Tablet friability

All tablet formulas meet the friability requirements, which is less than 1% (United States Pharmacopeia Convention, 2024b). Tablets with fragility values respectively from low to high are formula ab (0.55%), a (0.66%), b (0.74%) and the highest is formula 1 (0.85%).

Based on Table 8, the analysis results showed that temperature ($p = 0.0001$) and drying time ($p = 0.0053$) significantly influenced the tablet friability, contributing 68.5911% and 20.1717%, respectively. Meanwhile, the interaction between temperature and drying time was not significant ($p = 0.9952$) and had a very small contribution ($\approx 0.00005\%$). The negative effect values for both main factors indicated that increasing temperature and drying time tended to decrease the tablet friability. Thus, the response variation was primarily controlled by temperature and time factors independently.

Tablet hardness

Table 7 indicates that all formulations meet the weight uniformity acceptance criteria within the range of 4-8 kg (Rori, dkk., 2016). Formula a and b exhibit the highest hardness, measuring 5.58 kg and 5.71 kg, respectively, while formula 1 and b show hardness values of 3.73 kg and 5.57 kg. The greater hardness observed in formula a and b is attributed to the drying temperature applied. Figure 3(a), (b), and (c) presents a contour plot of tablet friability, hardness and disintegrating time respectively, where red shading represents higher hardness, and blue shading represents lower hardness. Higher drying temperatures and longer drying times reduce the moisture content in granules, enhancing particle bonding during compression and leading to harder tablets. Conversely, excessive moisture can result in brittle tablets, while insufficient moisture reduces cohesion, making tablets prone to cracking and breaking (Maskuriah *et al.*, 2021). The factorial equation predicting tablet hardness is represented as: $Y = 5.90 + 0.191X_A + 1.98X_B - 0.0331X_AX_B$ where Y is the tablet hardness response, X_A denotes temperature, X_B represents drying time, and X_AX_B accounts for the interaction between temperature and time.


Table 7. Results of Evaluation of Paracetamol Tablet Characteristics

Parameter	F1	Fa	Fb	Fab
Weight Variation Test NP	2.93	6.66	7.03	7.94
Tablet Hardness (kg) $\bar{X} \pm SD$	3.73 $\pm$ 0.48	5.58 $\pm$ 0.40	5.71 $\pm$ 0.53	5.57 $\pm$ 0.60
Tablet Friability (%) $\bar{X} \pm SD$	0.85 $\pm$ 0.05	0.66 $\pm$ 0.03	0.74 $\pm$ 0.03	0.55 $\pm$ 0.05
Disintegration Time (min) $\bar{X} \pm SD$	2 $\pm$ 1	6.67 $\pm$ 1.52	3 $\pm$ 1	7.33 $\pm$ 1.52

SD, standard deviation.

Table 8. Effect of Temperature, Time, and Interaction on Tablet Friability

Factor	Effect Value	%Contribution	<i>p</i> -value	<i>p</i> -value Equation
Temperature	-0.1872	68.5911	0.0001	0.0004
Time	-0.1015	20.1717	0.0053	(Significant)
Interaction	0.0002	5.43886E-05	0.9952	

**Figure 3.** Contour Plot of Tablets Test Response (a) Hardness Tester; (b) Flowability Tester; and (c) Disintegration Time.



According to Table 9, temperature, drying time, and their interaction significantly affect tablet hardness, with *p*-values <0.05 for each factor. Temperature has an effect value of 0.085 and contributes 21.61% to tablet hardness, whereas drying time has an effect value of 0.98 with a contribution of 28.50%. Drying time has a more significant influence compared to temperature. These results indicate that the combination of both factors has a more significant impact than each factor individually.

Tablet disintegration time

Based on Table 7, tablets with disintegration time respectively from low to high are formula 1 (2minutes), b (3 minutes), a (6.67 minutes) and the highest is formula ab (7.33 minutes). Table 10 shows that temperature significantly affects tablet disintegration time, with a *p*-value of 0.0003 and a contribution of 79.67%. Drying time has a minor influence (2.73%) and is not statistically significant (*p*-value 0.2960). Meanwhile, the interaction between temperature and time contributes 0.11% with a *p*-value of 0.8287, which is not statistically significant. Therefore,

Table 10. Effect of Temperature, Time, and Interaction on Tablet Disintegration Time

Factor	Effect Value	%Contribution	p-value	p-value Equation
Temperature	4.5	79.6721	0.0003	0.0021
Time	0.8333	2.7322	0.2960	(Significant)
Interaction	-0.1666	0.1092	0.8287	

**Figure 4.** Interaction of Drying Temperature and Time on (a) Friability Tester; (b) Hardness Tester; and (c) Disintegration Time.**Figure 5.** Superimposed Contour Plot.

temperature is the primary factor influencing tablet disintegration time.

Figure 4(a) shows the interaction between temperature and drying time, depicted by black and red lines that are not parallel. Increasing in temperature from 40°C to 60°C reduces tablet friability, indicating improved flow properties. Figure 4(b) shows that the interaction between temperature and drying time is represented by black and red lines that are not parallel and intersect at high temperatures. At low temperatures (40°C), tablet hardness significantly increases with longer drying times. Conversely, tablet hardness decreases at high temperatures (60°C) when the drying time is extended to 6 hours, indicating an over-drying effect that makes granules brittle and increases the risk of cracking and chipping. The interaction between temperature and time is statistically significant. Figure 4(c) shows the interaction between temperature and drying time with black and red lines that are not parallel. The interaction is minimal, as the lines do not intersect or touch, indicating that both factors work independently, with temperature being the dominant factor. Lower temperatures accelerate tablet disintegration, while higher temperatures result in longer disintegration times. Figure 5 shows the superimposed contour plot with the optimal drying temperature marked by the yellow area indicating an optimal response for the physical properties of granules and tablets.

CONCLUSIONS

Optimization of drying temperature and time in the production of paracetamol tablets using wet granulation with Polyvinylpyrrolidone K-25 shows a significant effect on the physical properties of granules and tablets. There are optimal responses for the physical properties of granules and tablets that meet quality standards. The optimal formulation was achieved at 55°C for 4 hours. Formula A (40°C, 6 hours) and Formula B (60°C, 3 hours) demonstrated the best results in producing tablets with desirable physical properties.

CONFLICT OF INTEREST

The authors declared no conflict of interest.

REFERENCES

- Ardiansyah., Rahmawati, T.E., Nabila, N.A., Maliza, F.N., Antoni. F., 2022. Optimasi formula tablet parasetamol dengan metode granulasi basah (Wet Granulation). *Jurnal Farmasi*, 2: 29. <https://journal.aisyahuniversity.ac.id/index.php/IFA/article/view/ardianoptimasi>
- Abimanyu, A.R.P., Rahma, A.D., Putri, D.R., Ismayfatin, H., Audia, W.A., Fatmala, W., Khasanah, Y., Yuniarsih, N., 2023. Formulasi dan uji efektifitas terhadap tablet paracetamol dengan metode granulasi basah dan kempa langsung: Riview artikel. *Innovative: Journal of Social Science Research* 3(2): 7476–92. <https://j-innovative.org/index.php/Innovative/article/view/939>
- Cahyani, A.N., Susanto, A., Dewi, I.R., Nurkhikmah, I., 2023. Formulasi tablet parasetamol dengan kombinasi PVP dan Amilum Umbi Porang (*Amorphophallus onchophyllus*) sebagai bahan pengikat terhadap sifat fisik tablet. *Jurnal Ilmiah Jophus*, 4: 6-7. <https://doi.org/10.46772/jophus.v4i02.886>
- Direktorat Jenderal Kefarmasian dan Alat Kesehatan, 2020. Farmakope Indonesia, jilid VI. Kementerian Kesehatan Republik Indonesia, Jakarta.
- Elisabeth, V., YamLean, P.V.Y., Supriati, H.S., 2018. Formulasi sediaan granul dengan bahan pengikat pati kulit pisang Goroho (*Musa acuminata* L.) dan pengaruhnya pada sifat fisik granul. *Jurnal Ilmiah Farmasi*, 7: 1–11. [doi: https://doi.org/10.35799/pha.7.2018.21416](https://doi.org/10.35799/pha.7.2018.21416)
- Hiremath, P., Nuguru, K., Agrahari, V., 2019. *Handbook of Pharmaceutical Wet Granulation*. Academic Press, America.
- Kurakula, M., Rao, G.S.N.K., 2020. Pharmaceutical assessment of polyvinylpyrrolidone (PVP): As excipient from conventional to controlled delivery system with a spotlight on COVID-19 inhibition. *Journal of Drug Delivery Science and Technology*, 60: 102046. [doi: 10.1016/j.jddst.2020.102046](https://doi.org/10.1016/j.jddst.2020.102046)
- Mariyani, K. A., Arisanti, I. S. and Setyawan, E. I. (2016). Pengaruh Konsentrasi Amilum Jagung Pregelatinasi sebagai Bahan Penghancur Terhadap Sifat Fisik Tablet Vitamin E. *J. Farm. Udayana*, vol. 2, pp. 39–49
- Maskuriah, D.Z., Wicaksono, Y., Isnawati, N., 2021. Pengaruh suhu dan waktu pengeringan pada formulasi tablet ekstrak daun Kelor (*Moringa oleifera*) dengan metode granulasi basah menggunakan

- Fluid Bed Dryer. *Internasional Journal of Islamic Medicine*, 2: 62.
- Noval, Kuncahyo, I., Pratama, A., Nabillah, S., Hatmayana, R., 2021. Formulasi sediaan tablet effervescent dari ekstrak etanol tanaman Bundung (*Actionoscirpus grossus*) sebagai antioksidan [Formulation effervescent tablets of Bundung plants (*Actinoscirpus grossus*) ethanol extract as a antioxidant]. *Jurnal Surya Medika*, 7(1): 131. doi: <https://doi.org/10.33084/jsm.v7i1.2649>
- Rori, W. M., Yamlean, P. V., & Sudewi, S. (2016). Formulasi dan Evaluasi Sediaan Tablet Ekstrak Daun Gedi Hijau (*Abelmoschus manihot*) Dengan Metode Granulasi Basah. *Pharmacon*, 5(2): 243-50. <https://doi.org/10.35799/pha.5.2016.12212>.
- Sudarsono, A.P.P., Nur, M., & Febrianto, Y., 2021. Pengaruh perbedaan suhu pengeringan granul (40°C,50°C,60°C) terhadap sifat fisik tablet paracetamol. *Jurnal Farmasi & Sains Indonesia*, 4: 44-51. doi: [10.52216/jfsi.v4i1.72](https://doi.org/10.52216/jfsi.v4i1.72)
- Syukri, Y., 2018. *Teknologi Sediaan Obat Dalam Bentuk Solid*. Islam Indonesia University Press, Yogyakarta.
- Taylor K.M.G., and Aulton, M.E., 2022. Aultons's Pharmaceutics the Design and Manufacture of Medicines. 6th edition. Elsevier.
- United States Pharmacopeia Convention, 2024a. <1174> Powder Flow. United States Pharmacopeia National Formulary 46 NF-41. U.S. Pharmacopeia Convention, Inc., Rockville.
- United States Pharmacopeia Convention, 2024b. <1216> Tablet Friability. United States Pharmacopeia National Formulary 46 NF-41. U.S. Pharmacopeia Convention, Inc., Rockville.
- Zaman NN & Sopyan I. 2020. Metode Pembuatan dan Kerusakan Fisik Sediaan Tablet', *Majalah Farmasetika*, 5(2): 82-93. doi: <https://doi.org/10.24198/mfarmasetika.v5i2.26260>.



JURNAL FARMASI SAINS DAN KOMUNITAS (JOURNAL OF PHARMACEUTICAL SCIENCES AND COMMUNITY)

📍 [FAKULTAS FARMASI UNIVERSITAS SANATA DHARMA](#)

☀️ P-ISSN : 16935683 ↔ E-ISSN : 25277146 🌐 Subject Area : Health

 **1.4**
Impact

 **1919**
Google
Citations


 **Sinta 2**
Current
Accreditation

🔗 [Google Scholar](#) 🔗 [Garuda](#) 🌐 [Website](#) 🌐 [Editor URL](#)

History Accreditation

2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029

Citation Per Year By Google Scholar



Journal By Google Scholar

	All	Since 2021
Citation	1919	1372
h-index	21	17
i10-index	57	43

Garuda

[Google Scholar](#)

Search...

[Quality Assessment of Extemporaneous Capsule Preparation containing Methylprednisolone, CTM, and Theophylline using Spectrophotometrics and Chemometrics Techniques](#)

[Sanata Dharma University](#)  [Jurnal Farmasi Sains dan Komunitas \(Journal of Pharmaceutical Sciences and Community\) Vol 22, No 1 \(2025\) 151-163](#)

📅 2025 📄 DOI: [10.24071/jpsc.008772](#) 🏷️ [Accred : Sinta 2](#)

[Determination of Sun Protection Factor Value and Physical Stability of Wungu Leaf \(*Graptophyllum pictum* \(L.\) Griff\) Extract Spray Sunscreen](#)

[Sanata Dharma University](#)  [Jurnal Farmasi Sains dan Komunitas \(Journal of Pharmaceutical Sciences and Community\) Vol 22, No 1 \(2025\) 1-9](#)

📅 2025 📄 DOI: [10.24071/jpsc.006632](#) 🏷️ [Accred : Sinta 2](#)

[Health Related Quality of Life \(HRQoL\) in Type 2 Diabetes Mellitus Patients Based on Antidiabetic Prescription Patterns](#)

[Sanata Dharma University](#)  [Jurnal Farmasi Sains dan Komunitas \(Journal of Pharmaceutical Sciences and Community\) Vol 22, No 1 \(2025\) 141-150](#)

📅 2025 📄 DOI: [10.24071/jpsc.008444](#) 🏷️ [Accred : Sinta 2](#)

[Formulation of Lerak Extract \(*Sapindus rarak* DC\) and Aloe Vera Extract \(*Aloe vera* Linn\) as Paper Hand Soap: Application of Factorial Design](#)

Sanata Dharma University. [Jurnal Farmasi Sains dan Komunitas \(Journal of Pharmaceutical Sciences and Community\) Vol 22, No 1 \(2025\) 115-122](#)
2025 [DOI: 10.24071/jpsc.008106](#) [Accred : Sinta 2](#)

Effectiveness of 70% Ethanol Extract of Pomegranate (*Punica granatum* Linn) Against the Healing Process of Incisions and Burns in Mice

Sanata Dharma University. [Jurnal Farmasi Sains dan Komunitas \(Journal of Pharmaceutical Sciences and Community\) Vol 22, No 1 \(2025\) 10-17](#)
2025 [DOI: 10.24071/jpsc.006779](#) [Accred : Sinta 2](#)

Formulation of Nanoemulsion Sunscreen of *Centella asiatica* (L.) Extract and Its Effectivity Test

Sanata Dharma University. [Jurnal Farmasi Sains dan Komunitas \(Journal of Pharmaceutical Sciences and Community\) Vol 22, No 1 \(2025\) 170-181](#)
2025 [DOI: 10.24071/jpsc.009910](#) [Accred : Sinta 2](#)

Antimicrobial Activity Test of Silver Nanoparticle Gel from Green Synthesis using Kelor Leaf (*Moringa oleifera*) as a Bioreductant Against *Cutibacterium acne*

Sanata Dharma University. [Jurnal Farmasi Sains dan Komunitas \(Journal of Pharmaceutical Sciences and Community\) Vol 22, No 1 \(2025\) 195-201](#)
2025 [DOI: 10.24071/jpsc.008272](#) [Accred : Sinta 2](#)

Knowledge and Attitudes towards Antibiotic Use among Non-Health Undergraduate Students at Andalas University

Sanata Dharma University. [Jurnal Farmasi Sains dan Komunitas \(Journal of Pharmaceutical Sciences and Community\) Vol 22, No 1 \(2025\) 44-50](#)
2025 [DOI: 10.24071/jpsc.007052](#) [Accred : Sinta 2](#)

The Predisposing Factors to Communicate Effectively in Providing Drug Information

Sanata Dharma University. [Jurnal Farmasi Sains dan Komunitas \(Journal of Pharmaceutical Sciences and Community\) Vol 22, No 1 \(2025\) 18-29](#)
2025 [DOI: 10.24071/jpsc.007028](#) [Accred : Sinta 2](#)

Routine Stock Opname in Enhancing Service of Halodoc's Order and Increasing Transactions

Sanata Dharma University. [Jurnal Farmasi Sains dan Komunitas \(Journal of Pharmaceutical Sciences and Community\) Vol 22, No 1 \(2025\) 134-140](#)
2025 [DOI: 10.24071/jpsc.008323](#) [Accred : Sinta 2](#)